



Department of Agriculture
STATE OF HAWAII

LICENSED

PERIOD **2008-2010** LIC. NO.

3-15/16"



Department of Agriculture
STATE OF HAWAII

LICENSED

PERIOD **2011-2013** LIC. NO.

4-3/16"

Hartz®

ULTRA
GUARD™

Flea & Tick
Spray for
Dogs

Kills:



Fleas



Ticks



KEEP OUT OF
REACH OF CHILDREN

CAUTION: SEE BACK
PANEL FOR INGREDIENTS AND
ADDITIONAL PRECAUTIONS.

16 fl. oz. (473ml)

RM105863

4-3/16"

H I N G E

DIRECTIONS FOR USE: It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

SHAKE WELL BEFORE USE

1. Hold bottle upright 6 inches from pet. Direct spray towards pet and spray entire coat, pressing dispenser with quick short strokes. Move bottle to get even coverage of coat (until tips of hair are moist).
2. Apply lightly and rub into animal's coat. For longhaired dogs, ruffle hair for spray to reach skin.
3. Recommended dosage: Spray 25-30 strokes for a small dog (e.g. Terrier). Spray 30-40 strokes for a medium dog (e.g. Spaniel). Spray 40-70 strokes for a large dog (e.g. Labrador). More spray may be needed for longhaired dogs.
4. Ticks are tough—spray directly.
5. After 10 minutes, dry dog with towel.
6. Comb and brush coat.
7. Repeat once a week. Use on bedding and other areas as needed.

STORAGE AND DISPOSAL

Storage: Store at room temperature.

Disposal: If empty: Do not re-use this container. Place in trash or offer for recycling if available. **If partly filled:** Call your solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

BACK OF
TOP PLY
(Page 2)

4-3/16"

1/4"

3-15/16"



Department of Agriculture
STATE OF HAWAII

LICENSED

PERIOD **2014-2016** LIC. NO.

9235.422

TOP PLY
(Page 1)

4-3/16"

↑ For full
precautions and
directions for use,
peel back here.

Hartz®
UltraGuard™
Flea & Tick
Spray for Dogs

When your dog has fleas
and ticks, your family and pet
may be exposed to:

- Lyme disease
- Tapeworm infestation
- Flea bite anemia
- Flea allergy dermatitis

To learn more, visit www.hartz.com

For more information, call our flea and tick experts
at 1-800-275-1414

ACTIVE INGREDIENT:

Tetrachlorvinphos 1.08%

OTHER INGREDIENTS: 98.92%

TOTAL 100.00%

Hartz® and other trademarks are trademarks of The Hartz Mountain Corporation.

Made by ©The Hartz Mountain Corporation,
400 Plaza Drive, Secaucus, NJ 07094

EPA Reg. No. 2596-125
EPA Est. No. 2596-NJ-1
Patented Formula
U.S. Patent No. 5,595,749



RM105864

3-15/16"

4-3/16"

READ ENTIRE
LABEL BEFORE
EACH USE. USE
ONLY ON DOGS

**PRECAUTIONARY
STATEMENTS
HAZARDS TO
HUMANS AND
DOMESTIC ANIMALS**

CAUTION: Causes moderate eye irritation. Avoid contact with eyes or clothing. Wash hands and exposed skin thoroughly with soap and water after handling. Do not use on puppies under 6 weeks old. Consult a veterinarian before using this product on debilitated, aged, or medicated animals. Sensitivity may occur after using ANY pesticide product for pets. If signs of sensitivity occur, bathe your pet with mild soap and rinse with large amounts of water. If signs continue, consult a veterinarian immediately. Do not use near birds, fish or foodstuffs. If used on pregnant or nursing dogs, do not retreat until the puppies are at least 6 weeks old. Do not use this product simultaneously or within 30 days before or after treatment with or exposure to cholinesterase inhibiting drugs, pesticides or chemicals. However, flea and tick collars may be immediately replaced after use.

FIRST AID: Contains an organophosphate that inhibits cholinesterase.

If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a Poison Control Center or doctor for treatment advice. Have the product container or label with you when calling a Poison Control Center or doctor, or going for treatment.

NOTE TO PHYSICIANS AND VETERINARIANS: This product contains an organophosphate and may cause cholinesterase inhibition. Cholinergic symptoms may include salivation, miosis, incoordination, muscle fasciculation and/or weakness, vomiting and diarrhea. Atropine is antidotal. 2-PAM may be effective as an adjunct to atropine. Call 1-800-275-1414 or your local Poison Control Center for further information.

BASE PLY
(Page 3)

H I N G E

4-3/16"

15/16"

1/4"

11/16"