

Longlife[®] 90 Day[™] Collar
by Hartz[®] FOR CATS

Longlife[®] 90 Day[™] Collar
by Hartz[®] FOR CATS

KILLS FLEAS • KILLS TICKS

Kills Deer Ticks which may carry the organism which causes Lyme Disease



ACTIVE INGREDIENT:	BY WEIGHT
Tetrachlorvinphos.....	14.55%
OTHER INGREDIENTS:	85.45%
TOTAL	100.00%

Do not let children play with this collar

CAUTION

See back panel for Precautionary Statements and Directions for Use

CONTENTS: 1 COLLAR, NET WT. 0.4 oz 11g

Longlife[®] 90 Day[™] Collar
by Hartz[®] FOR CATS

MONEY-BACK GUARANTEE BY Longlife[®] by Hartz[®]



Department of Agriculture
STATE OF HAWAII

Department of Agriculture
STATE OF HAWAII

LICENSED

LICENSED

PERIOD 2011-2013 LIC. NO. PERIOD 2014-2016 LIC. NO.

9235.462

READ ENTIRE LABEL BEFORE EACH USE. USE ONLY ON CATS PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION: Harmful if swallowed or absorbed through skin. Causes moderate eye irritation. Avoid contact with eyes, skin or clothing. Wash hands thoroughly with soap and water after handling.

FIRST AID - Contains an organophosphate that inhibits cholinesterase.

IF SWALLOWED: Call a Poison Control Center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a Poison Control Center or doctor. Do not give anything by mouth to an unconscious person.

IF ON SKIN OR CLOTHING: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a Poison Control Center or doctor for treatment advice.

IF IN EYES: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a Poison Control Center or doctor for treatment advice.

Have the product container or label with you when calling a Poison Control Center or doctor, or going for treatment.

NOTE TO PHYSICIANS AND VETERINARIANS

This product contains an organophosphate and may cause cholinesterase inhibition. Cholinergic symptoms may include salivation, miosis, incoordination, muscle fasciculation and/or weakness, vomiting, and diarrhea. Atropine is antidotal. 2-PAM may be effective as an adjunct to atropine. Call 1-800-275-1414 or your local Poison Control Center for further information.

DIRECTIONS FOR USE: It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

REMOVE COLLAR FROM PACKAGE, UNROLL AND STRETCH TO ACTIVATE INSECTICIDE GENERATOR.

Do not use on kittens under 12 weeks of age. Do not unroll collar until ready to use. Place the Longlife[®] by Hartz[®] 90 Day[™] Brand Collar for Cats around the cat's neck, adjust for proper fit, and buckle in place. The collar must be worn loosely to allow for growth of the cat and to permit the collar to move about the neck. Generally, a properly fitted collar is one that when fastened will snugly slide over the cat's head. Leave 2 or 3 inches on the collar for extra adjustment



and cut off and dispose of the extra length. Consult a veterinarian before using this product on debilitated, aged, medicated, pregnant or nursing animals. Sensitivity may occur after using ANY pesticide product for pets.

Some animals may become irritated by any collar if it is applied too tightly. If this occurs, loosen the collar. If irritation continues, remove the collar from the animal. If signs of sensitivity occur, remove the collar and bathe your pet with mild soap and rinse with large amounts of water. If signs continue, consult a veterinarian immediately. Do not use this product on animals simultaneously or within 30 days before or after treatment with or exposure to cholinesterase inhibiting drugs, pesticides, or chemicals. However, flea & tick collars may be immediately replaced with new collars. The collar is intended for use only as an insecticide generator.

For continuous flea protection under normal conditions, replace the collar every 3 months. Under conditions where cats are exposed to severe flea infestation, it may be necessary to replace the collar every 2 months. Ticks may be an occasional problem for cats. The collar will kill ticks, including the Deer Tick, which may carry Lyme Disease, for 60 days and continue to aid in their control an additional 30 days. The cat should be examined from time to time and the collar replaced when mature tick or flea populations begin to appear. Wetting will not impair the collar's effectiveness or pet's protection. If the cat is out in the rain, it is not necessary to remove the collar. The Longlife[®] by Hartz[®] 90 Day[™] Brand Collar for Cats may be worn with a regular collar.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage and disposal.


Pesticide Storage: Store this product in a cool, dry place that is inaccessible to children.

Pesticide Disposal: If empty: Nonrefillable container. Do not reuse or refill this container. Offer for recycling, if available. If partly filled: Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

For more information call our flea & tick experts at 1-800-275-1414 weekdays, 9 am - 5 pm E.S.T.
EPA Reg. No. 2596-49 EPA Est. No. 2596-OH-1
The Hartz Mountain Corporation Secaucus, NJ 07094
Hartz[®] and other trademarks are trademarks of The Hartz Mountain Corporation.

Made and printed in U.S.A.

RM109472



To activate collar hold at each end and STRETCH

