

Hartz **ADVANCED™**
Flea Treatment for **CATS**

For use ONLY on cats
8 weeks and older
and weighing
over 9 lbs.

**CONTAINS
THE SAME ACTIVE
INGREDIENTS USED IN
ADVANTAGE® II* PRODUCTS**

Hartz®

ADVANCED™
Flea Treatment for **CATS**

For use ONLY on cats
8 weeks and older
and weighing
over 9 lbs.



kills flea eggs



kills fleas



kills life stages

Kills fleas for 30 days

**4 Monthly
Treatments**

ACTIVE INGREDIENTS:

Imidacloprid 9.10%
Pyriproxyfen 0.46%
Other Ingredients 90.44%
TOTAL 100.00%

**KEEP OUT OF REACH OF CHILDREN
CAUTION:**

See back panel for Precautionary
Statements and Directions for Use.

NET CONTENTS:
0.108 fl. oz. (3.2 ml),
Contains 4 tubes,
each 0.027 fl. oz. (0.8 ml)

*Hartz Advanced™ Flea Treatment for Cats
is not manufactured by or distributed
by Bayer. Advantage® II is a registered
trademark of Bayer Healthcare, LLC.



LICENSED

PERIOD **2015-2017** LIC. NO. **9235.522**

Hartz **ADVANCED™** Flea Treatment for **CATS**

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS. CAUTION:

Harmful if swallowed. Causes moderate eye
irritation. Avoid contact with eyes
or clothing. Wash hands thoroughly with soap
and warm water after handling. Keep out of
reach of children. Do not contaminate feed or food.

HAZARDS TO DOMESTIC ANIMALS

For external use only. Do not apply to cats or kittens
under 8 weeks of age or weighing less than 9 lbs. As
with any product, consult your veterinarian before
using this product on debilitated, aged, pregnant or
nursing cats. Individual sensitivities, while rare, may
occur after using ANY pesticide product for cats. If
signs persist, or become more severe, consult a
veterinarian immediately. If your cat is on medication,
consult your veterinarian before using this or any
other product.

FIRST AID

Have the product container or label with you
when calling a poison control center or doctor, or
going for treatment. For medical emergencies or
consumer questions call 1-800-275-1414.


IF SWALLOWED: Call a poison control center or
doctor immediately for treatment advice. Have
person sip a glass of water if able to swallow.
Do not induce vomiting unless told to do so by a
poison control center or doctor. Do not give
anything by mouth to an unconscious person.

IF IN EYES: Hold eye open and rinse slowly and
gently with water for 15-20 minutes. Remove
contact lenses, if present, after the first 5
minutes, then continue rinsing eye. Call a poison
control center or doctor for treatment advice.

IF ON SKIN OR CLOTHING: Take off
contaminated clothing. Rinse skin immediately
with plenty of water for 15 - 20 minutes. Call a
poison control center or doctor for treatment or
advice.

For more information call our experts at
1-800-275-1414 (weekdays, 9 am - 5 pm E.S.T.)
EPA Reg. No. 2596-181 EPA Est. No. 2596-OH-1
Hartz® and other trademarks are trademarks of
The Hartz Mountain Corporation.

Made by The Hartz Mountain Corporation,
400 Plaza Drive, Secaucus, NJ 07094 (USA)



0 32700 14737 2 RM112584

For the Prevention and Treatment of Flea Infestations on Cats

DIRECTIONS FOR USE:

It is a violation of Federal law to use this product in
a manner inconsistent with its labeling.
Do not allow children to apply product.

READ ENTIRE LABEL BEFORE EACH USE

RESTRICTIONS

**USE ONLY ON CATS OR KITTENS
8 WEEKS AND OLDER WEIGHING
OVER 9 lbs.**

**DO NOT USE ON OTHER ANIMALS.
DO NOT EXCEED LABELED DOSAGE
AMOUNT FOR CATS.**

**DO NOT APPLY MORE
THAN ONE (1) TUBE PER
TREATMENT ON ANY CAT.**

**DO NOT HAVE CONTACT
OR ALLOW CHILDREN TO
HAVE CONTACT WITH
TREATED AREA UNTIL
COMPLETELY DRY.**

Side Effects (Cats): Monitor your cat after application. Side
effects, although very rare, may include signs of skin irritation such as
redness, scratching, or other signs of discomfort. Gastrointestinal
signs such as hypersalivation, vomiting or diarrhea have also been
reported. If these or other side effects (such as lethargy) occur,
consult your veterinarian or call, 1-800-275-1414.


HOW TO APPLY



1) Remove one
applicator tube
from the
package.



2) Hold applicator tube in an
upright position **facing away from
you and your pet's face and eyes.**



3) Twist
dispensing tip
clockwise
about 1/2 turn
while pushing
down to break the tube's seal. Do
not remove the dispensing tip.

4) Part the hair on the neck at the base of the
skull until the skin is visible. Place the tip of
the tube on the skin and squeeze the tube to
expel the entire contents directly on the skin.
**Do not get this product in your pet's eyes,
or allow your pet to ingest this product.**
The product is bitter tasting and salivation
may occur for a short time if the pet licks
the product immediately after treatment.

Treatment at the base of the skull will
minimize the opportunity for the cat to lick the product. Do not allow the
product to run off.

5) Discard empty tube as described in the Storage and Disposal section.

6) Under normal conditions this product is effective for a month. However, in
case of severe flea infestation, retreatment may be necessary earlier than
four (4) weeks. On cats, do not retreat more often than once every seven (7)
days. After flea control is attained, return to a monthly treatment schedule.

STORAGE AND DISPOSAL

PESTICIDE STORAGE: Store in a cool, dry place inaccessible to children and pets.

PESTICIDE DISPOSAL AND CONTAINER HANDLING:

Non-refillable container. If empty: Do not reuse or refill this container. Place in
trash or offer for recycling if available. If partly filled: Call your local solid waste
agency or 1-800-275-1414 for disposal instructions. Never place unused product
down any indoor or outdoor drain.



112584

