

# RONSTAR® 0.95% Plus Fertilizer



**NOT FOR USE** on plants being grown for sale or other commercial use, or for commercial seed production, or for research purposes. For use on plants intended for aesthetic purposes or climatic modification and being grown in ornamental gardens or parks, or on golf courses or lawns and grounds.

## NOT FOR USE ON HOME LAWNS

Contains LESCO® Poly Plus® Sulfur Coated Urea to provide a uniform growth with extended nitrogen feeding.

### ACTIVE INGREDIENT:

Oxadiazon; 2-tert-butyl-4-(2,4-dichloro-5-isopropoxyphenyl)  
 $\Delta$ 1,3,4-oxadiazolin-5-one ..... 0.95%

**OTHER INGREDIENTS:** ..... 99.05%  
 Total ..... 100.00%

### SUGGESTED SPREADER SETTINGS

| Spreader            | Ground Speed | Recommended Rates in lb per 1,000 sq ft |         |         |        |
|---------------------|--------------|---|---------|---------|--------|
|                     |              | 4.8 lb                                  | 7.25 lb | 8.25 lb | 9.6 lb |
| LESCO               | 3 mph        | 14                                      | 17      | 18      | 19     |
| Cyclone® or Spyker® | 3 mph        | 4                                       | 4½      | 4¾      | 5      |
| Vicon®              | 4.5 mph      | 22                                      | 32      | 34      | 36     |
| Lely®               | 4.5 mph      | 4 II                                    | 4½ II   | 5 II    | 5½ II  |

\*IMPORTANT: These settings are only approximate. Age, condition of spreader, and operator speed can cause wide variation. Be sure to calibrate your spreader with each application using the LESCO Calibrating Gauge #009952.

### WHEN TO APPLY

Appropriate Times to Apply This Pre-emergent Herbicide:

| Jan. | Feb. | Mar.     | Apr.      | May       | June | July     | Aug. | Sep.      | Oct.     | Nov. | Dec. |
|------|------|----------|-----------|-----------|------|----------|------|-----------|----------|------|------|
|      |      | 1. _____ |           | 2. _____  |      | 3. _____ |      |           |          |      |      |
|      |      |          | 1a. _____ |           |      |          |      | 3a. _____ |          |      |      |
|      |      |          |           | 2a. _____ |      |          |      |           | 4. _____ |      |      |

1. For Crabgrass and other Grasses and Broadleaf Weeds germinating in the Spring in Northern Turf.

1a. For Crabgrass in Southern Turf.  
 2. For Goosegrass and other Late Germinating Annual Grasses and Broadleaf Weeds such as Prostrate Spurge.

2a. Goosegrass in Southern Turf.  
 3. For Poa annua in Northern Turf  
 3a. For Poa annua in Southern Turf.

4. For Fall-Winter-Spring control of Annual Weeds in Southern Turf.

### PRECAUTIONS FOR USING

Do not use on desirable Red Fescue, Bentgrass varieties, Dichondra, or Centipedegrass since damage will occur. Do not use on putting greens or "tees" due to varying cultural practices used. Use only the rate of 50 lb to treat 6,900 sq ft (7.25 lb/1,000 sq ft) on St. Augustinegrass since temporary discoloration will occur from heavier rates. Avoid contact with flowers and shrubs except as recommended on the Ornamental Use Supplement Labeling available for this product. For newly seeded lawns, do not apply this product until after the second or third cutting.

### RESEEDING TREATED AREAS

Delay new seeding until 4 months after treatment.

### MINI TURF FERTILIZER

# 19-2-19

### GUARANTEED ANALYSIS

|  |        |
|--|--------|
| TOTAL NITROGEN (N) .....   | 19.00% |
| 0.66% Ammoniacal Nitrogen*   |        |
| 18.34% Urea Nitrogen*  |        |
| AVAILABLE PHOSPHATE (P <sub>2</sub> O <sub>5</sub> ) .....                                   | 2.00%  |
| SOLUBLE POTASH (K <sub>2</sub> O) .....  | 19.00% |
| SULFUR (S) Total .....   | 16.21% |
| 9.26% Free Sulfur (S)  |        |
| 6.95% Combined Sulfur (S)  |        |
| DERIVED FROM: Polymer Coated Sulfur Coated Urea, Urea, Ammonium Phosphate, Sulfate of Potash |        |
| *16.20% Slowly available Urea Nitrogen from Polymer Coated Sulfur Coated Urea.               |        |

### CONDITION OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

**NOTICE:** Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product must be followed carefully and completely. It is impossible to eliminate all risks inherently associated with the use of this product. To the extent consistent with applicable law, Buyer and/or User assume all risks of ineffectiveness or other unintended consequences or damages that may result from conditions outside or beyond the control of LESCO, Inc. including but not limited to, such factors as manner of use or application, weather or weather conditions outside the range considered normal at the application site or for the time period in which the product is applied, the presence of other materials, incompatible products, or other influencing factors which are beyond the control of LESCO, Inc. To the extent consistent with applicable law, all such risks shall be assumed by Buyer and/or User, and Buyer and/or User agrees to hold LESCO, Inc. harmless for any claims relating to such factors.

LESCO, Inc. warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with the Directions for Use under normal use conditions. To the extent consistent with applicable law, this warranty does not extend to the use of this product contrary to label instructions, or under abnormal conditions or under conditions not reasonably foreseeable to or beyond the control of LESCO, Inc. and Buyer and/or User assumes the risk of any such use. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, LESCO, INC. MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE BUYER AND/OR USER AND THE EXCLUSIVE LIABILITY OF LESCO, INC. FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF LESCO, INC. REPLACEMENT OF THE PRODUCT, OR IF NOT ACQUIRED BY PURCHASE, REPLACEMENT OF SUCH QUANTITY. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO EVENT SHALL LESCO, INC., BE LIABLE FOR ANY INCIDENTAL, CONSEQUENTIAL OR SPECIAL DAMAGES.

LESCO, Inc. offers this product, and Buyer and/or User accepts it, subject to the foregoing Conditions of Sale and Limitation of Warranty and Liability, which may not be modified except by written agreement signed by a duly authorized representative of LESCO, Inc.

Information concerning the raw materials composing this product can be obtained by writing to: LESCO, Inc., Attn: RA Dept., 1301 East 9th Street, Suite 1300, Cleveland, Ohio 44114-1849, referring to the Item number found on this bag.

Information regarding the contents and levels of metals in this product is available on the Internet at <http://www.regulatory-info.com>

Contains RONSTAR Herbicide, a registered trademark of Bayer. LESCO and Poly Plus are registered trademarks and the sweeping design is a trademark of LESCO Technologies, LLC. Spyker and Cyclone are registered trademarks of Spyker Spreaders LLC. Lely is a registered trademark of C Van Der Lely N.V. Vicon is a registered trademark of Ransomes Cushman-Ryan Company.

GAREGULWPIERIEVIEWAprvd Lbl - Word089270.doc  
 Created: 3/23/07 (6/07/02) Rev.: 3/27/07 DB



Net Weight: 50 lb (22.7 kg)



## KEEP OUT OF REACH OF CHILDREN WARNING

### FIRST AID

|                               |  |
|-------------------------------|--|
| <b>If inhaled</b>             | <ul style="list-style-type: none"> <li>Move person to fresh air.</li> <li>If person is not breathing, call 911 or ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible.</li> <li>Call a poison control center or doctor for further treatment advice.</li> </ul>   |
| <b>If on skin or clothing</b> | <ul style="list-style-type: none"> <li>Take off contaminated clothing.</li> <li>Rinse skin immediately with plenty of water for 15-20 minutes.</li> <li>Call a poison control center or doctor for treatment advice.</li> </ul>  |
| <b>If in eyes</b>             | <ul style="list-style-type: none"> <li>Hold eye open and rinse slowly and gently with water for 15-20 minutes.</li> <li>Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li> <li>Call a poison control center or doctor for treatment advice.</li> </ul>  |
| <b>If swallowed</b>           | <ul style="list-style-type: none"> <li>Call poison control center or doctor immediately for treatment advice.</li> <li>Have person sip a glass of water if able to swallow.</li> <li>Do not induce vomiting unless told to do so by the poison control center or doctor.</li> <li>Do not give anything by mouth to an unconscious person.</li> </ul> |

### HOT LINE NUMBER

Have the product container or label with you when calling a poison control center or doctor, or going to treatment. You may also contact 1-(800) 424-9300 for emergency treatment information.

### PRECAUTIONARY STATEMENTS

#### HAZARDS TO HUMANS AND DOMESTIC ANIMALS

**WARNING:** Causes eye and skin irritation. Avoid contact with skin, eyes, or clothing. Thoroughly wash face and hands with soap and water after exposure and before eating or smoking. Wear protective clothing including long-sleeved shirt and gloves when handling. Launder exposed clothing before reuse. Do not allow domestic animals to graze treated areas. Avoid contamination of feed and foodstuffs.

#### ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish. Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not apply when weather conditions favor drift from treated areas. Do not apply where runoff is likely to occur. Do not contaminate water when disposing of equipment washwaters. Apply only as specified on this label.

#### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

### STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.  
**Storage:** Store product in its original bag, in a cool, dry, locked place out of the reach of children. In case of spill, sweep up, and use according to directions or dispose of as follows:  
**Pesticide Disposal:** Pesticide wastes are toxic. Improper disposal of excess pesticide or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the hazardous waste representative at the nearest EPA regional office for guidance.  
**Container Disposal:** Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

### FOR LAWNS AND ORNAMENTAL TURF

When applied according to directions, this product can be used safely on the following established turfgrass varieties: Bermudagrass, Perennial Bluegrass, Perennial Ryegrass, St. Augustinegrass, Tall Fescue, and Zoysiagrass lawns and ornamental turf. It may be used for parks, golf courses, athletic fields, cemeteries, and all other ornamental and recreational turfgrass areas.

### CONTROLS GRASSY WEEDS AND BROADLEAF WEEDS

Annual Bluegrass (*Poa annua*), Barnyardgrass, Crabgrass, Fall Panicum, Florida Pusley, Goosegrass, Green Foxtail, Oxalis, Stinging Nettle, Carpetgrass, Pigweed.

### HOW TO APPLY

Measure the area to be treated and uniformly apply amount using a fertilizer spreader (see Spreader Settings chart). Do not apply to turf when it is wet. Do not overlap spreader application on St. Augustinegrass since discoloration "streaking" may temporarily occur from "double rate". After application, if rain is not expected within 24 hours, sprinkle the treated turf with 1/4" to 1/2" of water to get maximum weed control.

### HOW MUCH TO USE

**For Crabgrass:** Apply 50 lb/10,400 sq ft (4.8 lb/1,000) when turf has been in a yearly maintenance program (equal to 2 lb A.I. herbicide per acre). Apply 50 lb to treat 6,900 sq ft (7.25 lb/1,000) of lawn if it has been heavily infested with crabgrass (equal to 3 lb A.I. herbicide per acre). Apply in early spring 1 to 2 weeks prior to weed seed germination. This product is a pre-emergent herbicide. It will not control emerged and growing weeds; only germinating weed seeds. Late applications will control only those weed seeds that germinate after the application. See "When to Apply" chart.

**Goosegrass and Prostrate Spurge:** These weeds do not usually germinate until late spring or early summer. Apply 50 lb to treat 5,200 sq ft of lawn (9.6 lb/1,000 sq ft; 4 lb A.I./acre) and apply 1 to 2 weeks prior to seed germination. See chart for when to apply. Observe maximum rate for St. Augustinegrass: 50 lb/6,900 sq ft.

**For Annual Bluegrass (*Poa annua*):** Apply 50 lb per 5,200 to 6,050 sq ft of lawn (8.25-9.6 lb/1,000 sq ft; equal to 3.5 to 4 lb A.I. herbicide per acre). Apply in late summer to fall or in early spring 1 to 2 weeks before expected weed seed germination. Will not control Annual Bluegrass present at time of application, only germinating seeds. Refer to chart for proper timing in your locale. Annual Barnyardgrass and Prostrate Spurge may also be controlled by this same rate. Observe maximum rate for St. Augustinegrass: 50 lb/6,900 sq ft.

EPA Reg. No. 10404-63

EPA Est. No. 82757-OH-001 (M), 82757-FL-001 (S)

First letter of lot code indicates manufacturing site.

# #089270

F1560

Distributed by: LESCO, Inc.

1301 East 9th Street

Cleveland, OH 44114-1849