Description:	18lb Lilly Miller Ultra Green Weed & Feed 28-2-3
UPC #:	70624-16326
Consumer Unit Size:	D: 2.25 IN. W: 11 IN H: 24.75 IN.
Program:	Adobe Illustrator CS4
Printer:	Printpack
Creation Date:	05.03.2011
Rev. Date:	09.00.2011

Ink Colors: Black, Reflex Blue, Yellow, 2nd Black, Magenta, Cyan, PMS 350, PMS 116, PMS 354

UPC Check | Proofread/QC

Initial Date (Schawk Minneapolis) | Proofreader Initial Date

Numerical Scan Only: Pass Fail | Quality Control Initial Date

SCHAWK! 2626 2nd Street NE, Minneapolis, MN 55418 USA 612.789.8514 schawk.com

The color on this copy will not match the final printed package.
Schawk Minneapolis has checked this artwork for accuracy. Final approval is the client's responsibility.
Please double check for errors before reproduction.

28.000

2.250 | 11.000 | 2.250 | 11.000

24.750

Department of Agriculture STATE OF HAWAII
LICENSED
PERIOD 2014-2016 LIC. NO.
9317.323

Department of Agriculture STATE OF HAWAII
LICENSED
PERIOD 2011-2013 LIC. NO.

Quick Green That Lasts Up to Two Months

COVERS UP TO 5,000 SQ. FT.

Lilly Miller®

ULTRAGREEN®

WEED & FEED with TRIMEC® herbicide

28•2•3

- Controls over 200 weeds!*
- Kills even the tough weeds like Dandelion, Plantain and Clover
- Builds thick, green lawns from the roots up

SPRING	SUMMER	FALL	WINTER
GREEN CYCLE™	CRABGRASS CONTROL & LAWN FOOD	WEED & FEED	LAWN FOOD	FALL & WINTER LAWN FOOD

Lilly Miller® ULTRAGREEN® feeds and protects all year long!

- Reverdecimiento rápido que dura hasta dos meses
- Cubre hasta 5,000 pies cuadrados
- ¡Controla más de 200 malezas!
- Elimina incluso las malezas difíciles de controlar como el diente de león, lantén y trébol
- Desarrolla céspedes tupidos y verdes desde las raíces hasta arriba

ACTIVE INGREDIENTS:
Dimethylamine salt of 2,4-dichlorophenoxyacetic acid 0.65%
Dimethylamine salt of (+)-(R)-2-(2-methyl-4-chlorophenoxy)propionic acid 0.16%
Dimethylamine salt of dicamba (3,6-dichloro-o-anisic acid)... 0.03%
INERT INGREDIENTS 99.13%
TOTAL 100.00%

THIS PRODUCT CONTAINS:
18.00 lbs. 2,4-dichlorophenoxyacetic acid equivalent per ton or 0.54%
2.6 lbs. (+)-(R)-2-(2-methyl-4-chlorophenoxy) propionic acid equivalent per ton or 0.13%
1.2 lbs. 3,6-dichloro-o-anisic acid equivalent per ton or 0.06%
Isomer Specific by AOAC Methods

TRIMEC® is a registered trademark of PBI/GORDON CORPORATION.

KEEP OUT OF REACH OF CHILDREN
CAUTION
SEE BACK PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS & FIRST AID
MANTÉNGASE FUERA DEL ALCANCE DE LOS NIÑOS
PRECAUCIÓN
LEA EL PANEL POSTERIOR PARA MÁS DECLARACIONES DE PRECAUCIÓN Y PRIMEROS AUXILIOS

NET WT./PESO NETO 18 LBS. (8.1 kg)

Lilly Miller®
ULTRAGREEN® WEED & FEED with TRIMEC® herbicide
28•2•3
COVERS UP TO 5,000 SQ. FT.

Lilly Miller® ULTRAGREEN® WEED & FEED with TRIMEC® herbicide 28•2•3 COVERS UP TO 5,000 SQ. FT.

Lilly Miller® ULTRAGREEN® WEED & FEED with TRIMEC® herbicide 28•2•3

GUARANTEED ANALYSIS

Total Nitrogen (N)	28%
9.5% Ammoniacal Nitrogen	
18.5% Urea Nitrogen*	
Available Phosphate (P_2O_5)	2%
Soluble Potash (K_2O)	3%
Sulfur (S)	9%
9.0% Combined Sulfur (S)	
Iron (Fe)	2%

Derived from Urea, Ammonium Sulfate, Polymer Coated Urea, Ammonium Phosphate, Muriate of Potash, Iron Sucrate.

*7.0% Slowly available urea nitrogen from polymer coated urea.

Information regarding the contents and levels of metals in this product is available on the internet at http://www.aapfco.org/metals.htm

Filled by weight not by volume. Product may settle after packaging.

This product may stain concrete and stone. Where spilling occurs, sweep off immediately, then flush with water.

Llenado por peso y no por volumen. El producto puede asentarse después de empaquetar.

Es posible que este producto manche el concreto y la piedra. Donde sucedan derrames, quite inmediatamente con una escoba y luego lave con agua.

READ THE ENTIRE LABEL FIRST.
OBSERVE ALL PRECAUTIONS AND FOLLOW DIRECTIONS CAREFULLY.

STOP

LEA PRIMERO TODA LA ETIQUETA. RESPETE TODAS LAS PRECAUCIONES Y SIGA LAS INSTRUCCIONES CON CUIDADO.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

STORAGE & DISPOSAL
Do not contaminate water, food, or feed by storage and disposal.
PESTICIDE STORAGE: Store in original container in a locked storage area inaccessible to children and domestic animals. Do not store near food or feedstuffs.
PESTICIDE DISPOSAL & CONTAINER HANDLING: Nonrefillable container. Do not reuse this container. *If Empty*—Place in trash or offer for recycling if available. *If Partly Filled*—Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

GENERAL PRECAUTIONS AND RESTRICTIONS:
- Do not apply this product in a way that will contact any person or pet, either directly or through drift. Keep people and pets out of the area during application. Do not allow people or pets to enter the treated area until dusts have settled.
- Do not use this product where desirable clovers are present. Apply when the air is calm. Do not apply to newly seeded lawns. During application, close spreader openings when turning.
- Do not use this product on or near desirable plants, and avoid spreading granules onto exposed root systems or adventitious shoots (rhizomes) within the drip line of desirable trees and shrubs, since injury may result.
- Do not apply directly to or near water, storm drains, gutters, sewers, or drainage ditches. Do not apply within 25 feet of rivers, fish ponds, lakes, streams, reservoirs, marshes, estuaries, bays, and oceans. Do not apply when windy. Apply this product directly to your lawn, and sweep any product landing on the driveway, sidewalk, gutter, or street back onto the treated area. To prevent product run-off, do not over water the treated area(s) to the point of runoff or apply when raining or when rain is expected that day.

INSTRUCTIONS
Lilly Miller® ULTRAGREEN® Weed & Feed controls a wide range of lawn weeds including dandelion, chickweed, knotweed, plantain, henbit, spurge and many others commonly found in home lawns. This product may be applied to established lawns of bermudagrass, bentgrass, bahiagrass, centipedegrass, Kentucky bluegrass, fescues, perennial ryegrass, zoysiagrass, and St. Augustinegrass. Do not use on dichondra or carpetgrass. Do not apply this product to 'Floratam' St. Augustinegrass in Florida. In addition, a 28-2-3 fertilizer provides plant nutrients.

Possible retreatment may be necessary if (a) lawn is heavily overrun by weeds, or, (b) if adverse low moisture conditions prevail and weeds are in a state of poor growth. A second application may be made in approximately 30 days.

1. Mow lawn to normal height 1 to 2 days before application.
2. Water lawn thoroughly at least 1 day before application to sustain moisture until the next watering (see step 4).
3. Apply when weeds are young and growing actively, preferably in the morning when dew is on the grass. If grass is not moist at time of application, sprinkle lightly with water to allow the granules to adhere and to remain on the leaf surface of the weeds.
4. Do not wash from weed leaves for 1 to 2 days after application. At this time a thorough watering should be made.

Apply at the rate of 1 lb. of product per 275 sq. ft.

The maximum broadcast application rate is 4 pounds of product per 1,000 sq. ft. per application (0.94 lb. 2,4-D ae, 0.23 lb. MCPP-p ae and 0.10 lb. dicamba ae per acre per application). The maximum number of broadcast applications is limited to 2 per year with a minimum of 30 days between applications. The maximum seasonal rate is 8 pounds of product per 1,000 sq. ft. (1.88 lbs. 2,4-D ae, 0.46 lb. MCPP-p ae and 0.20 lb. dicamba ae per acre).

SETTING YOUR LAWN SPREADER
THESE SETTINGS ARE A GUIDE. They were determined by landscape professionals using new spreaders. The condition of your spreader, your walking speed, the roughness of the lawn, and even the humidity can affect how fast the lawn food comes out. Watch the flow rate. If fertilizer appears to be coming out either slower or faster than you are covering the lawn, adjust the setting. Close the spreader when filling, turning or stopping.

INSTRUCCIONES

Usar este producto de modo inconsistente con las indicaciones de su etiqueta constituye una violación de la ley federal.

ALMACENAMIENTO Y ELIMINACIÓN
Al almacenarlo y eliminarlo no contamine el agua, comida o alimento para animales.
ALMACENAMIENTO DEL PESTICIDA: Almacénelo en el envase original en un área de almacenamiento bajo llave inaccesible para niños y animales domésticos. No lo almacene cerca de comida o alimento para animales.
ELIMINACIÓN DEL PESTICIDA Y MANEJO DEL ENVASE: Envase que no se puede rellenar. No vuelva a usar este envase. *Si está vacío*—tírelo a la basura o, si está disponible, ofrézcalo para reciclaje. *Si está parcialmente lleno*—llame a su agencia local de desechos sólidos para obtener instrucciones sobre cómo eliminarlo. Nunca elimine el producto sin usar en ningún drenaje interior o exterior.

PRECAUCIONES GENERALES Y RESTRICCIONES:
- No aplique este producto de modo que entre en contacto con alguna persona o mascota, ya sea directamente o por diseminación. Durante la aplicación mantenga fuera del área a las personas y mascotas. No permita que los niños o las mascotas entren en el área tratada hasta que el polvo se haya fijado.
- No use este producto cuando haya tréboles que desee conservar. Aplíquelo cuando el aire esté tranquilo. No lo aplique a los céspedes recién sembrados. Durante la aplicación, al girar cierre las aberturas del esparcidor.
- No use este producto en plantas que desee conservar o cerca de ellas, y evite esparcir gránulos en los sistemas de raíces expuestas o con yemas (rizomas) dentro de la línea de goteo de árboles y arbustos que desee conservar pues puede dañarlos.
- No lo aplique directamente o cerca de agua, drenajes pluviales, alcantarillas o zanjas de drenaje. No lo aplique a menos de 25 pies de ríos, estanques de peces, lagos, arroyos, represas, pantanos, esteros, bahías y océanos. No lo aplique cuando haya viento. Aplique este producto directamente a su césped y barra cualquier producto que caiga en el camino de acceso, aceras, cuneta o la calle de regreso al área tratada. Para prevenir la escorrentía, no riegue sobre las áreas tratadas al punto en que haya escorrentía ni lo aplique cuando esté lloviendo o se espere lluvia durante ese día.

INSTRUCCIONES
Lilly Miller® ULTRAGREEN® Weed & Feed controla una amplia gama de malas hierbas de gramas; por ejemplo el diente de león, estalaria, sanguinaria mayor, llantén, ortiga mansa, euforbia y varias otras que comúnmente se encuentran en el césped de los hogares. Se puede aplicar este producto al césped establecido de grama Bermuda, grama Bentgrass, grama Bahia, grama ciempiés, pasto azul de Kentucky, cañuela, raigrás perenne, grama Zoysia y grama de San Agustín. No use en oreja de ratón o grama alfombra. No aplique este producto en grama de San Agustín "Floratam" en Florida. Además, un fertilizante 28-2-3 ofrece a las plantas los nutrientes o el alimento.

Puede ser necesario el posible retratamiento si (a) el césped está sumamente poblado de malas hierbas, o (b) si prevalecen las condiciones adversas de baja humedad y las malas hierbas se encuentra en un estado de poco crecimiento. Se puede realizar una segunda aplicación en aproximadamente 30 días.

1. Corte el césped a una altura normal 1 o 2 días antes de la aplicación.
2. Riegue el césped completamente por lo menos 1 día antes de la aplicación para mantener la humedad hasta el siguiente riego (Vea el paso 4).
3. Aplique cuando las malas hierbas estén jóvenes y crezca activamente, de preferencia en la mañana cuando existe rocío en el césped. Si la grama no está húmeda al momento de la aplicación, rocíela ligeramente con agua para permitir que se adhieran los gránulos y que permanezcan en la superficie de las hojas de las malezas.
4. No lave las hojas de las malas hierbas durante 1 o 2 días después de la aplicación. En este momento, se debe hacer un riego completo.

Aplique en una proporción de 1 lb. de producto por cada 275 pies cuadrados.

La dosis máxima de aplicación al voleo es 4 libras de producto por cada 1,000 pies cuadrados por aplicación (0.94 lb. 2,4-D ae, 0.23 lb. MCPP-p ae y 0.10 lb. dicamba ae por acre por aplicación). La cantidad máxima de aplicaciones al voleo está limitada a 2 por año con un mínimo de 30 días entre aplicaciones. La dosis máxima por temporada es 8 libras de producto por cada 1,000 pies cuadrados (1.88 lb. 2,4-D ae, 0.46 lb. MCPP-p ae y 0.20 lb. dicamba ae por acre).

AJUSTE DE SU ESPARCIDOR PARA CÉSPED
ESTOS AJUSTES SON PAUTAS. Paisajistas profesionales que usan los nuevos esparcidores los han determinado. La condición de su esparcidor, su velocidad al caminar, el mantenimiento del césped e incluso la humedad pueden afectar la velocidad en que surge el abono para plantas. Controle la tasa de flujo. Si el fertilizante parece que sale más lento o más rápido de lo que usted cubre el césped, calibre el ajuste. Cierre el esparcidor cuando llene, gire o se detenga.

(1.8 lbs. for 500 sq. ft.) / *(1.8 libras para 500 pies cuadrados)*

Drop Spreader Models / *Modelos de esparcidor de goteo*	Setting / *Ajuste*
• Scotts® AccuGreen® 1000	8.0
• Scotts® AccuGreen® 2000	8.0
• Scotts® AccuGreen® 3000	8.0
• Earthway®	13.0
• Precision Red Devil™	4.0
• Vigoro®	4.0

Rotary Spreader Models / *Modelos de esparcidor giratorio*	Setting / *Ajuste*
• Scotts® SpeedyGreen® 1000	8.0
• Scotts® SpeedyGreen® 2000	8.0
• Scotts® SpeedyGreen® 3000	8.0
• Earthway®	15.0
• Precision Red Devil™	11.0
• Vigoro®	7.5

WEEDS*

Annual yellow sweetclover	Buckhorn plantain	Clover, white	Hawkweed	Mustard	Prostrate spurge	Spiny cocklebur	Wild aster
Aster	Bulbous buttercup	Cockle	Healall	Narrowleaf plantain	Prostrate vervain	Spiny sowthistle	Wild buckwheat
Austrian fieldcress	Bull thistle	Cocklebur	Heartleaf drymary	Narrowleaf vetch	Puncturevine	Spotted catsear	Wild carrot
Bedstraw	Bullnettle	Common mullein	Heath aster	Nettle	Purslane, common	Spotted spurge	Wild four-o'clock
Beggarticks	Burclover	Creeping jenny	Hedge bindweed	Orange hawkweed	Ragweed	Spurweed	Wild garlic
Betony, Florida	Burdock	Cudweed	Hedge mustard	Oriental cocklebur	Red sorrel	Stinging nettle	Wild geranium
Bindweed, field	Burning nettle	Curly dock	Henbit	Oxalis	Redroot pigweed	Strawberry, India mock	Wild lettuce
Bird vetch	Burweed	Daisy, English	Hoary cress	Parsley-piert	Redstem filaree	Tall nettle	Wild marigold
Bitter wintercress	Buttercup	Daisy fleabane	Hoary plantain	Parsnip	Rough cinquefoil	Tall vervain	Wild mustard
Bittercress, hairy	Buttonweed	Daisy, oxeye	Hoary vervain	Pearlwort	Rough fleabane	Tansy ragwort	Wild onion
Bittersweet	Canada thistle	Dandelion	Horsenettle	Pennycress	Russian pigweed	Tansy mustard	Wild parsnip
Black-eyed Susan	Carolina geranium	Dichondra	Jimsonweed	Pepperweed	Russian thistle	Tanweed	Wild radish
Black medic	Carpetweed	Dogfennel	Knawel	Pigweed	Scarlet pimpernel	Thistle	Wild rape
Black mustard	Catchweed bedstraw	Dollarweed	Knotweed	Pineywoods bedstraw	Scotch thistle	Trailing crownvetch	Wild strawberry
Blackseed plantain	Catnip	False dandelion	Kochia	Plains coreopsis	Sheep sorrel	Tumble mustard	Wild sweet potato
Blessed thistle	Catsear	Falseflax	Lambsquarters	Plantain	Shepherdspurse	Tumble pigweed	Wild vetch
Bloodflower milkweed	Chickweed, common	False sunflower	Lespedeza	Poison ivy	Slender plantain	Velvetleaf	Woodsorrel
Blue lettuce	Chickweed, mouseear	Fiddleneck	Mallow	Poison oak	Smallflower galinsoga	Venice mallow	Woolly croton
Blue vervain	Chicory	Florida pusley	Matchweed	Pokeweed	Smooth dock	Virginia buttonweed	Woolly morningglory
Bracted plantain	Cinquefoil	Frenchweed	Mexicanweed	Poorjoe	Smooth pigweed	Virginia creeper	Woolly plantain
Bristly oxtongue	Clover, crimson	Galinsoga	Milk vetch	Prairie sunflower	Sorrel	Virginia pepperweed	Wormseed
Broadleaf dock	Clover, hop	Goldenrod	Morningglory	Prickly lettuce	Sowthistle	Wavyleaf bulthistle	Yarrow
Broadleaf plantain	Clover, red	Ground ivy	Mousear hawkweed	Prickly sida	Spanishneedles	Western clematis	Yellow rocket
Broomweed	Clover, strawberry	Gumweed	Mugwort	Prostrate knotweed	Speedwell	Western salsify	Yellowflower pepperweed
Buckhorn	Clover, sweet	Hairy fleabane	Musk thistle	Prostrate pigweed	Spiny amaranth	White mustard	and other broadleaf weeds

PRECAUTIONARY STATEMENTS

Hazards to Humans & Domestic Animals
CAUTION: Harmful if swallowed. Harmful if absorbed through skin. Causes moderate eye irritation. Avoid contact with eyes, skin or clothing. Wear long-sleeved shirt, long pants, shoes, socks and chemical-resistant gloves made of any waterproof material. Prolonged or frequently repeated skin contact may cause allergic reaction in some individuals. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet.

First Aid
If in eyes: Hold eye open and rinse slowly and gently with water for 15 - 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.
If on skin or clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15 - 20 minutes. Call a poison control center or doctor for treatment advice.
If swallowed: Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.
Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also call toll-free, 1-877-800-5556 for emergency medical treatment information.

ENVIRONMENTAL HAZARDS:
This pesticide is toxic to fish and aquatic invertebrates and may adversely affect non-target plants. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Sweeping any product that lands on a driveway, sidewalk, or street, back onto the treated area of the lawn or garden will help to prevent run off to water bodies or drainage systems.

LIMITED WARRANTY AND DISCLAIMER
FOR USE ONLY AS DIRECTED. To the extent consistent with applicable law, the manufacturer neither makes nor intends any express or implied warranties, including any warranty of merchantability or fitness for a particular purpose, which are hereby expressly disclaimed. To the extent consistent with applicable law, in no case shall the manufacturer be liable for incidental, consequential, or special damages resulting from the use or handling of this product. If these terms are not acceptable, return this product unopened immediately to the point of purchase, and the purchase price will be returned in full. The terms of this LIMITED WARRANTY AND DISCLAIMER cannot be varied by any written or verbal statements or agreements at the point of sale or elsewhere.

NOTICE – This product contains Cadmium, Chromium, Cobalt, Lead, and Nickel, chemicals known to the State of California to cause cancer and birth defects or other reproductive harm.

Lilly Miller and ULTRAGREEN are registered trademarks of Central Garden & Pet.
AccuGreen® Scotts® SpeedyGreen® are registered trademarks of OMS Investments, Inc. CENTRAL Garden & Pet USA

DECLARACIONES DE PRECAUCIÓN

Dañino para los seres humanos y animales domésticos
PRECAUCIÓN: Dañino si se traga. Es dañino si se absorbe a través de la piel. Causa irritación moderada en los ojos. Evite el contacto con ojos, piel o ropa. Use pantalones largos, camisa de manga larga, zapatos, calcetines y guantes resistentes a químicos hechos de material a prueba de agua. El contacto repetido en la piel, prolongado o frecuente, puede causar reacción alérgica en algunas personas. Lave completamente con agua y jabón después de usarlo y antes de comer, beber, consumir goma de mascar o tabaco o usar el baño.

Primeros auxilios
Si ocurre contacto con los ojos: Mantenga el ojo abierto y enjuague lenta y suavemente con agua durante 15 ó 20 minutos. Retire los lentes de contacto, si hubiera, después de los primeros 5 minutos, luego continúe enjuagando los ojos. Llame a un centro de control de envenenamiento o a un médico para recibir consejos de tratamiento.
Si ocurre contacto con la piel o ropa: Quítese la ropa contaminada. Enjuague la piel inmediatamente con abundante agua durante 15 ó 20 minutos. Llame a un centro de control de envenenamiento o a un médico para recibir consejos de tratamiento.
Si se ingiere: Llame de inmediato a un centro de control de envenenamiento o a un médico para recibir consejos de tratamiento. Haga que la persona beba tragos de agua si puede tragar. No provoque el vómito a menos que el centro de control de envenenamiento o médico le indique hacerlo. No dé nada por la boca a una persona inconsciente.
Tenga consigo el envase o etiqueta del producto cuando llame al centro de control de envenenamiento o al médico, o acuda al tratamiento. También puede llamar gratuitamente al 1-877-800-5556 para obtener información sobre el tratamiento médico.

RIESGOS AMBIENTALES:
Este pesticida es tóxico para los peces y los invertebrados acuáticos y puede afectar de forma negativa a las plantas que no son el objetivo. Para proteger el ambiente, no permita que el pesticida entre o se escurra en drenajes pluviales, zanjas de drenaje, cunetas o aguas superficiales. Aplicar este producto en clima tranquilo cuando no haya pronóstico de lluvia para las siguientes 24 horas ayudará a asegurarse de que el viento o la lluvia no soplen o laven el pesticida fuera del área de tratamiento. Barrer el producto que cae en un camino de acceso, aceras o en la calle de regreso al área tratada del césped o jardín ayudará a evitar que se escurra a los cuerpos de agua o sistemas de drenaje.

RENUNCIA Y GARANTÍA LIMITADA
PARA USARLO SOLAMENTE COMO SE INDICA. Hasta donde sea congruente con la ley que aplica, el fabricante no establece ni tiene la intención de establecer ninguna garantía expresa o implícita, incluyendo alguna garantía de comercialibilidad o idoneidad para un propósito en particular, a las que se renuncien por este medio. Hasta donde sea congruente con la ley que aplica, en ningún caso será responsable el fabricante por daños incidentales, consiguientes o especiales que resulten del uso o manejo de este producto. Si no son aceptables estos términos, regrese inmediatamente el producto sin abrir al punto de compra y se le devolverá el precio de compra en su totalidad. Los términos de esta RENUNCIA Y GARANTÍA LIMITADA no pueden variarse por ninguna declaración o acuerdo verbal o escrito en el punto de venta o en ningún otro lado.
ADVERTENCIA: este producto contiene cadmio, cromo, cobalto, plomo y níquel, químicos que el Estado de California conoce como causantes de cáncer y defectos de nacimiento u otros daños reproductivos.

SATISFACTION GUARANTEED
If for any reason you are not satisfied with this product, please send proof of purchase to the address shown and we will gladly refund your purchase price.

0 70624 16326 9

Distributed by:
LILLY MILLER BRANDS
2000 Powers Ferry Rd., Suite 270
Atlanta, GA 30339

EPA Reg. No. 2217-559-33116
EPA Est. 35480-OR-1

www.lillymiller.com

300029131-5/11

SATISFACCIÓN GARANTIZADA
Si por alguna razón usted no se encuentra satisfecho con este producto, sírvase enviar el comprobante de compra a la dirección mostrada y gustosamente le reembolsaremos el precio de su compra.

Cylinder

Web