



STATE OF HAWAII
Department of Agriculture

ACCEPTED

LICENSE NO. _____

**Fungicide
Insecticide**

**Dust or
Spray**

**Lilly
Miller**

9317.344

SULFUR DUST

*For use on roses, grapes,
citrus, fruits, berries, nuts,
vegetables, flowers,
shrubs, trees*

*Controls powdery mildew,
rust, scab, mites*



POWDERY MILDEW



RUST



MITES

ACTIVE INGREDIENT:

Sulfur 90%

INERT INGREDIENTS 10%

Keep out of reach of children

CAUTION

See back booklet for additional
Precautionary Statements and First Aid.

NET WT. 1 LB (453 g)

100524478-12/15
LM50 v.4



SULFUR DUST

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. For household use only.

USE PRECAUTIONS

Do not apply LILLY MILLER® SULFUR DUST dusts or sprays when shade temperatures during the day are likely to exceed 85 - 90 degrees F. Injury to foliage and fruit may result. Do not use LILLY MILLER® SULFUR DUST within 30 days of an oil spray (60 days on citrus). Do not combine with an oil spray.

HOW TO USE

Please read the entire label. Use only as directed, noting all use precautions.

AS A DUST: Before dusting, shake duster vigorously to mix air with the sulfur. Dust lightly. Frequent light dustings are more effective than single heavy dustings.

AS A SPRAY: Mix the required amount of LILLY MILLER® SULFUR DUST with sufficient water to make a thin, creamy paste before diluting with the balance of the required amount of water.

FRUITS AND NUTS

ALMONDS: Brown Rot, Scab. Dust or spray (3 tablespoons per gallon of water) before blossoms open, during bloom, and after bloom. Apply when disease threatens during rainy periods and repeat at 7 to 10 day intervals.

APPLES, PEARS: Scab, Powdery Mildew. Spray (3 tablespoons per gallon of water) when buds begin to swell and thereafter every 10 days until 2 weeks after the petals fall. Treat Delicious Apples and Anjou Pears only before bloom or injury may occur.

BERRIES (BLACKBERRIES, BOYSENBERRIES, GOOSEBERRIES, LOGANBERRIES, RASPBERRIES, STRAWBERRIES): Powdery Mildew. Dust or spray (1 tablespoon per gallon of water) when disease first appears and repeat at 7 to 10 day intervals or as necessary to maintain control.

LILLY MILLER BRANDS

1000 Parkwood Circle, Suite 700

Atlanta, GA 30339

1-800-265-0761

gardendivision@central.com

www.lillymiller.com

Printed in U.S.A.

EPA Reg. No. 802-16

EPA Est. 67572-GA-1, 67572-GA-2

Subscript used is first letter
of Run Code on Container

Lilly Miller, Lilly Miller with design, Central Garden & Pet and Central Garden & Pet with design are trademarks of Central Garden & Pet Company.

100524478 - 12/15

LM v.4



0 70624 50609 7

CHERRIES, NECTARINES, PEACHES, PLUMS, PRUNES: Powdery Mildew, Silver Mite, Brown Rot, Cherry Leaf Spot, Rust. Dust or spray (3 tablespoons per gallon of water) before blossoms open, during bloom, and after bloom. Apply when disease threatens during rainy periods and repeat at 7 to 10 day intervals. The first three applications will also provide control of Brown Rot Blossom Blight on Cherries and Peaches. Weekly applications from 5 weeks before and continuing up to the day of harvest will provide control of Brown Rot of Fruit.

CITRUS: Citrus Thrips, Citrus Flat Mite, Yuma Spider Mite, Sixspotted Mite, Citrus Rust Mite. Dust or spray (2 tablespoons per gallon of water) when pests first appear and repeat as necessary. Avoid application during or preceding high temperatures. Do not apply within two months of an oil application.

GRAPES: Powdery Mildew. Dust or spray (1 tablespoon per gallon of water) during the growing season when new shoot growth is 6 to 10 inches long and repeat at 7 to 10 day intervals. Concord and other Labrusca type grapes may be injured by LILLY MILLER® SULFUR DUST.

PISTACHIOS: Citrus Flat Mite. Dust or spray (3 tablespoons per gallon of water) when mites first appear and repeat as necessary.

VEGETABLES

BEANS: Powdery Mildew, Rust, Thrips, Spider Mites. Dust or spray (3 tablespoons per gallon of water) when pests are first noticed, repeat as necessary. For Rust control, start applications in early spring before any rust is noticed. Repeat applications at 7 day intervals until bloom.

CARROTS, ONIONS: Petrobia Mite. Dust when mites first appear. Repeat as necessary.

PEAS, TURNIPS, RUTABAGAS: Powdery Mildew. Dust when disease first appears and repeat at 7 to 14 day intervals.

TOMATOES: Tomato Russet Mite. Dust or spray (3 tablespoons per gallon of water) when pests first appear.

ORNAMENTAL FLOWERS, SHRUBS, TREES: Rust, Powdery Mildew. Dust or spray (3 tablespoons per gallon of water) to completely cover foliage. Repeat at 5 to 10 day intervals as needed. Follow directions given for specific plants.

CHRYSANTHEMUMS, CLEMATIS, RHODODENDRONS: Powdery Mildew. Dust or spray (3 tablespoons per gallon of water) at the first sign of mildew. Repeat at 4 to 5 day intervals until mildew is under control. Do not apply to open Chrysanthemum flowers as bleaching may result.

HOLLYHOCKS, SNAPDRAGONS: Rust. Dust or spray (1 tablespoon per gallon of water) in early spring. Repeat as needed until disease is under control - applications twice a week may be necessary.

ROSES: Powdery Mildew. Dust or spray (8 tablespoons per gallon of water) in spring when young growth first appears. Repeat at 4 to 5 day intervals until mildew is under control. See USE PRECAUTIONS.

STORAGE AND DISPOSAL

PESTICIDE STORAGE: Store this product in its original container and keep out of reach of children and domestic animals.

PESTICIDE DISPOSAL: *If partly filled:* Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

CONTAINER DISPOSAL: *If empty:* Nonrefillable container. Do not reuse or refill this container. Place in trash or offer for recycling if available.

Filled by weight not by volume. Product may settle after packaging.

NOTICE: To the extent consistent with applicable law, seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. To the extent consistent with applicable law, buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Causes moderate eye irritation. Harmful if inhaled, absorbed through skin, or swallowed. Avoid breathing dust. Avoid contact with skin, eyes, or clothing. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash clothing before reuse.

FIRST AID

Have product container or label with you when calling a poison control center or doctor, or going for treatment.

If swallowed:

- Call a poison control center or doctor immediately for treatment advice.
- Have person sip a glass of water if able to swallow.
- Do not induce vomiting unless told to do so by a poison control center or doctor.
- Do not give anything by mouth to an unconscious person.

If inhaled:

- Move person to fresh air.
- If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible.
- Call a poison control center or doctor for further treatment advice.

If on skin or on clothing:

- Take off contaminated clothing.
- Rinse skin immediately with plenty of water for 15-20 minutes.
- Call a poison control center or doctor for treatment advice.

If in eyes:

- Hold eye open and rinse slowly and gently with water for 15-20 minutes.
- Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.
- Call a poison control center or doctor for treatment advice.

ENVIRONMENTAL HAZARDS

Do not apply directly to water. Do not contaminate water when disposing of rinsate or equipment washwater.

LILLY MILLER BRANDS

1000 Parkwood Circle, Suite 700
Atlanta, GA 30339
1-800-265-0761
gardendivision@central.com
www.lillymiller.com

EPA Reg. No. 802-16
EPA Est. 67572-GA-1, 67572-GA-2₃
Subscript used is first letter of Run Code on Container
Printed in U.S.A.
100524478 - 12/15
LM50 v.4

