



Department of Agriculture  
STATE OF HAWAII

**LICENSED**

PERIOD **2015-2017** LIC. NO.

**9322.164**



Department of Agriculture  
STATE OF HAWAII

**LICENSED**

PERIOD **2012-2014** LIC. NO.

OPEN HERE ➤

# ACCELERON<sup>®</sup> IC-609

## Insecticide Seed Treatment For Corn

Acceleron<sup>®</sup> IC-609 Insecticide Seed Treatment for Corn is a systemic insecticide seed treatment for use on corn for the control of listed insect pests.

All seed treated with this product must be conspicuously coloured at the time of treatment.

**Active Ingredient:**

Clothianidin..... 48.0%

**Other Ingredients:** ..... 52.0%

*Total:* ..... 100.0%

Contains 5 lbs. active ingredient per gallon.

EPA Reg. No. 264-789-524

### KEEP OUT OF REACH OF CHILDREN CAUTION

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.  
(If you do not understand the label, find someone to explain it to you in detail.)

**FOR ADDITIONAL PRECAUTIONARY STATEMENTS: See label booklet.**

For **MEDICAL** And **TRANSPORTATION** Emergencies **ONLY**  
Call 24 Hours A Day **1-314-694-4000**

For **PRODUCT USE** Information Call Toll-Free **1-800-332-3111**

**Net Contents: 2.5 gallons**

100120L3-1

Packaged for: Monsanto Company • 800 North Lindbergh Blvd.  
St. Louis, MO 63167 • 1-314-694-4000

## FIRST AID

<b>IF SWALLOWED:</b>	<ul style="list-style-type: none"><li>• Immediately call a poison control center or doctor for treatment advice.</li><li>• Do not induce vomiting unless told to do so by a poison control center or doctor.</li><li>• Have person sip a glass of water if able to swallow.</li><li>• Do not give anything by mouth to an unconscious person.</li></ul>
<b>IF IN EYES:</b>	<ul style="list-style-type: none"><li>• Hold eye open and rinse slowly and gently with water for 15-20 minutes.</li><li>• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing.</li><li>• Call a poison control center or doctor for treatment advice.</li></ul>
<b>IF INHALED:</b>	<ul style="list-style-type: none"><li>• Move person to fresh air.</li><li>• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible.</li><li>• Call a poison control center or doctor for further treatment advice.</li></ul>
<b>IF ON SKIN OR CLOTHING:</b>	<ul style="list-style-type: none"><li>• Take off contaminated clothing.</li><li>• Rinse skin immediately with plenty of water for 15-20 minutes.</li><li>• Call a poison control center or doctor for treatment advice.</li></ul>

**For MEDICAL Emergencies Call 24 Hours A Day 1-314-694-4000.**

**Have the product container or label with you when calling a poison control center or doctor or going for treatment.**

**NOTE TO PHYSICIAN:** No specific antidote is available. Treat the patient symptomatically.

### Precautionary Statements

#### CAUTION

##### Hazards to Humans and Domestic Animals

Harmful if swallowed or absorbed through the skin. Avoid contact with skin or clothing. Wash hands thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse.

##### Personal Protective Equipment (PPE)

Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for category C on an EPA chemical resistance category selection chart.

**Applicators and other handlers must wear:** Long-sleeved shirt and long pants, chemical-resistant gloves (such as nitrile, butyl, neoprene, barrier laminate, polyvinyl chloride or Viton) and shoes plus socks. Follow manufacturer's instructions for cleaning/maintaining PPE. If no instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

### User Safety Recommendations

**Users should:** Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. Remove Personal Protective Equipment immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

### Environmental Hazards

This product is toxic to aquatic invertebrates. Contain any product spills or equipment leaks and dispose of wastes according to disposal instructions on this label. Do not contaminate water when disposing of equipment washwaters.

### Directions for Use

**It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Read entire label before using this product.**

For use by commercial treaters only. Not for use in agricultural establishments in hopper-box, slurry-box, or similar on farm seed treatment applicators used at planting. This product is to be used in commercial liquid or slurry treaters. Mix thoroughly before use or use entire container at one time.

All seed treated with this product must be conspicuously coloured at the time of treatment. This product contains no colorant. An appropriate colorant must be added when this product is applied to seed to distinguish and prevent subsequent inadvertent use as a food for man or feed for animals. Regulations pertaining to coloration of treated seed enforced by *40 CFR 153.155* must be strictly adhered to when using this product.

## Storage and Disposal

### PESTICIDE STORAGE

Do not contaminate water, food or feed by storage or disposal. Store in a cool, dry secured storage area.

### PESTICIDE DISPOSAL

Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

### CONTAINER DISPOSAL

Non-refillable container. Do not reuse or refill this container. Offer for recycling, if available. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Dispose of rinsed empty container in a sanitary landfill or by incineration, or if allowed by state and local authorities, by burning. If burned stay out of smoke.

### SEED QUALITY AND STORAGE OF TREATED SEED

Treatment of highly mechanically damaged seed, or seed of known low vigor or poor quality, may result in reduced germination and/or reduction of seed and seedling vigor. Treat and conduct germination tests on a small portion of seed before committing the total seed lot to a selected chemical treatment. Due to seed quality conditions beyond the control of Monsanto Company, no claims are made to guarantee germination of carry-over seed.

### LABELING OF TREATED SEED

The *Federal Seed Act* requires that the container of seed treated with Acceleron® IC-609 Insecticide Seed Treatment for Corn must be labeled with the following statements:

- This seed has been treated with Acceleron® IC-609 Insecticide Seed Treatment for Corn, which contains clothianidin.
- Do not use treated seed for food, feed or oil processing.

In addition, the US Environmental Protection Agency requires the following statements on the container of seed treated with Acceleron® IC-609 Insecticide Seed Treatment for Corn:

- Store away from feeds and foodstuffs.
- Wear long-sleeved shirt, long pants, shoes, socks, and chemical-resistant gloves when handling treated seed.
- Dispose of all excess treated seed. Left over treated seed may be double sown around the headland or

buried away from water sources in accordance with local requirements. Do not contaminate water bodies when disposing of planting equipment washwaters.

- Dispose of seed packaging in accordance with local requirements.
- This compound is toxic to birds and mammals. Treated seeds exposed on soil surface may be hazardous to birds and mammals. Cover or collect treated seeds spilled during loading.
- For Field Corn: Immediate Plant-back: Cereal grains, Field corn, popcorn, sweet corn, rapeseed, canola, soybeans, root and tuber vegetable, 30-Day Plant-back: Grasses, non-grass animal feeds, dry beans, 8-Month Plant-back: Sugar cane, 12-Month Plant-back: Any crops without an earlier plant-back interval.

Rotational crop plant-back intervals				
Crop Planted	Immediate Plant-back	30-Day Plant-back	8-Month Plant-back	12-Month Plant-back
Field corn	Cereal grains, Field corn, popcorn, sweet corn, rapeseed, canola, soybeans, root and tuber vegetables	Grasses, non-grass animal feeds, dry beans,	Sugar cane	Any crops without an earlier plant-back interval

### USE INFORMATION

Acceleron® IC-609 Insecticide Seed Treatment for Corn is a systemic insecticide seed treatment for use on corn for the control of listed insects.

For use in commercially available equipment designed for seed treatment only. Not for use in hopper-box, slurry-box, or similar seed treatment applications used at planting. This product is to be used in liquid or slurry treaters. Mix thoroughly before use or use entire container at one time. Pre-test all tank mixes to determine physical compatibility between formulations. Observe all cautions and limitations on labeling of all products used in mixtures.

### For Early Season Protection Against Listed Insects:

Acceleron® IC-609 Insecticide Seed Treatment for Corn will aid in the protection of seeds and seedlings against injury by listed early season insects.

### APPLICATION RATES

CROP	INSECT PESTS	RATE Acceleron® IC-609 Insecticide Seed Treatment for Corn	
		mg a.i./ Kernel	80,000 Seed Unit
Field corn	Corn rootworm (including Northern, Western, Southern and Mexican) Southern corn billbug <sup>1</sup>	1.25	5.64 fl oz or 166.7 mL
Field corn	Chinch bug Corn flea beetle Corn leaf aphid Cutworm, black Grape colaspis Seedcorn maggot Southern corn leaf beetle Southern corn rootworm Southern green stinkbug Storage pests <sup>2</sup> (including Indian meal moth, Lesser grain borer, Red flour beetle, Rice weevil, Rusty grain beetle) Sugarcane beetle White grub (including European, Chafer larvae, May/June beetle larvae, Japanese beetle larvae) Thrips Wireworm	0.25 to 0.50	1.13 fl oz or 33.3 mL to 2.26 fl oz or 66.6 mL
Field corn	Corn flea beetle (through first true seed leaf) Seedcorn maggot <sup>3</sup> Storage pests <sup>2</sup> (including Indian meal moth, Lesser grain borer, Red flour beetle, Rice weevil, Rusty grain beetle) Wireworm <sup>3</sup> White grub <sup>3</sup> (including European, Chafer larvae, May/June beetle larvae, Japanese beetle larvae) Wireworm <sup>3</sup>	0.125	0.57 fl oz or 16.7 mL

<sup>1</sup> Follow-up foliar sprays of a locally registered insecticide may be needed under heavy pest pressure. Consult your local Extension office for recommendations.

<sup>2</sup> It is recommended that seed with existing populations of storage pests be fumigated prior to treating and bagging seed.

<sup>3</sup> For seed protection only.

#### IMPORTANT: READ BEFORE USE

Read the entire Directions for Use, Conditions, Disclaimer of Warranties and Limitations of Liability before using this product. If terms are not acceptable, return the unopened product container at once.

Treatment of highly mechanically damaged seed, or seed of known low vigor and poor quality, may result in reduced germination and/or reduction of seed and seedling vigor. Treat and conduct germination tests on a small portion of seed before committing the total seed lot to a selected chemical treatment. Due to seed quality conditions beyond the control of Monsanto Company no claims are made to guarantee germination of carry-over seed.

By using this product, user or buyer accepts the following Conditions, Disclaimer of Warranties and Limitations of Liability.

**CONDITIONS:** The directions for use of this product are believed to be adequate and must be followed carefully. However, it is impossible to eliminate all risks associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of Monsanto Company. All such risks shall be assumed by the user or buyer.

**DISCLAIMER OF WARRANTIES:** TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, MONSANTO COMPANY MAKES NO OTHER WARRANTIES, EXPRESS OR IMPLIED, OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE OR OTHERWISE, THAT EXTEND BEYOND THE STATEMENTS MADE ON THIS LABEL. No agent of Monsanto Company is authorized to make any warranties beyond those contained herein or to modify the warranties contained herein. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, Monsanto Company disclaims any liability whatsoever for special, incidental or consequential damages resulting from the use or handling of this product.

**LIMITATIONS OF LIABILITY:** TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER FOR ANY AND ALL LOSSES, INJURIES OR DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER IN CONTRACT, WARRANTY, TORT, NEGLIGENCE, STRICT LIABILITY OR OTHERWISE, SHALL NOT EXCEED THE PURCHASE PRICE PAID, OR AT MONSANTO COMPANY'S ELECTION, THE REPLACEMENT OF PRODUCT.

**Acceleron** is a trademark of Monsanto Technology LLC.

Packaged For:  
Monsanto Company  
800 N. Lindbergh Blvd.  
St. Louis, MO 63167  
1-314-694-4000  
© 2013

# ACCELERON<sup>®</sup> IC-609

Acceleron<sup>®</sup> IC-609 Insecticide Seed Treatment for Corn is a systemic insecticide seed treatment for use on corn for the control of listed insect pests.

All seed treated with this product must be conspicuously coloured at the time of treatment.

**Active Ingredient:**

Clothianidin..... 48.0%

**Other Ingredients:** ..... 52.0%

*Total:* ..... 100.0%

Contains 5 lbs. active ingredient per gallon.

EPA Reg. No. 264-789-524

## KEEP OUT OF REACH OF CHILDREN CAUTION

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.  
(If you do not understand the label, find someone to explain it to you in detail.)

**FIRST AID: IF SWALLOWED:** Immediately call a poison control center or doctor for treatment advice. Do not induce vomiting unless told to do so by a poison control center or doctor. Have person sip a glass of water if able to swallow. Do not give anything by mouth to an unconscious person. **IF IN EYES:** Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing. Call a poison control center or doctor for treatment advice. **IF INHALED:** Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. Call a poison control center or doctor for further treatment advice. **IF ON SKIN OR CLOTHING:** Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

**For MEDICAL Emergencies Call 24 Hours A Day 1-314-694-4000. Have the product container or label with you when calling a poison control center or doctor or going for treatment. Precautionary Statements: CAUTION. Hazards to Humans and Domestic Animals.** Harmful if swallowed or absorbed through the skin. Avoid contact with skin or clothing. Wash hands thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse. **Environmental Hazards:** This product is toxic to aquatic invertebrates. Contain any product spills or equipment leaks and dispose of wastes according to disposal instructions on this label. Do not contaminate water when disposing of equipment washwaters. **Storage and Disposal: PESTICIDE STORAGE.** Do not contaminate water, food or feed by storage or disposal. Store in a cool, dry secured storage area. **PESTICIDE DISPOSAL.** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility. **CONTAINER DISPOSAL.** Non-refillable container. Do not reuse or refill this container. Offer for recycling, if available. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Dispose of rinsed empty container in a sanitary landfill or by incineration, or if allowed by state and local authorities, by burning. If burned stay out of smoke.

For **MEDICAL** And **TRANSPORTATION** Emergencies **ONLY** Call 24 Hours A Day **1-314-694-4000**

For **PRODUCT USE** Information Call Toll-Free **1-800-332-3111**

**REFER TO LABEL BOOKLET FOR COMPLETE DIRECTIONS FOR USE.**

**Net Contents: 2.5 gallons**

100120L3-1

**Packaged for: Monsanto Company • 800 North Lindbergh Blvd.  
St. Louis, MO 63167 • 1-314-694-4000**