Department of Agriculture
STATE OF HAWAII
LICENSED
PERIOD 2005-2007 LIC. NO.

Department of Agriculture
STATE OF HAWAII
LICENSED
PERIOD 2008-2010 LIC. NO.

Department of Agriculture
STATE OF HAWAII
LICENSED
PERIOD 2011-2013 LIC. NO.

Apply in Fall
When Weeds are Growing!

Expert GARDENER®

Winterizer Weed & Feed

HERBICIDA Y ABONO

Department of Agriculture
STATE OF HAWAII
LICENSED DISC.
PERIOD 2014-2016 LIC. NO.
9387.184

Kills Fall Weeds and Greens the Lawn!
¡Elimina Las Malezas y Enverdece el Césped!

- ***Kills 200+ Weeds While Providing Winter Lawn Protection and Early Spring Green-Up!***
- *Formulated with Scotts® Poly-S® Technology for Longer Lasting Greening and Feeding*

Net Wt 16.00 lb (7.25 kg)

22-3-11

CAUTION / PRECAUCIÓN:
KEEP OUT OF REACH OF CHILDREN.
MANTENGA FUERA DEL ALCANCE DE LOS NIÑOS.
See back for additional precautionary statements.

Do not use on Dichondra, Carpetgrass, or Floratam, a type of St. Augustinegrass.

ACTIVE INGREDIENTS:
2, 4-D, dimethylamine salt 0.723%
Mecoprop-P, dimethylamine salt 0.148%
Dichlorprop-P, dimethylamine salt 0.088%
INERT INGREDIENTS: 99.041%
TOTAL 100.000%

THIS PRODUCT CONTAINS:
2,4-dichlorophenoxyacetic acid equivalent, 0.601%.
(+)-(R)-2-(2-methyl-4-chlorophenoxy) propionic acid equivalent, 0.122%.
(+)-(R)-2-(2,4-dichlorophenoxy) propionic acid equivalent, 0.074%.
Isomer Specific by AOAC Methods. TRIMEC® is a registered trademark of PBI/Gordon Corporation.

Treats & Feeds 5,000 sq ft
(1/8 acre = 5,445 sq ft)

Kills Fall Weeds and Greens the Lawn! Apply in Fall!
Winterizer Weed & Feed
37105
1

Kills Fall Weeds and Greens the Lawn! Apply in Fall!
Winterizer Weed & Feed
37105
1

Expert Gardener® Fertilizer Program

For Best Results!

Use the Expert Gardener® Fertilizer Program!



	Early Spring	Mid/Late Spring	Summer	Fall	Grass Type
Northern	Crabgrass Preventer	Weed & Feed	Lawn Food	Winterizer or Winterizer Weed & Feed	Bluegrass, Rye, Fescues, Bermuda, Zoysia
Southern	Crabgrass Preventer	Weed & Feed	Lawn Food	Winterizer or Winterizer Weed & Feed	Bermuda, Fescues, Bahia
	Nothing until grass starts to green up	St. Augustine Weed & Feed or Lawn Food	Lawn Food	Winterizer or St. Augustine Weed & Feed	St. Augustine, Centipede, Zoysia

Nitrogen (N)
For Deep Green Color

Phosphorus (P)
For Strong Roots

Potassium (K)
For Overall Plant Health

Controls 200+ Weeds! Including:

Annual Ylw Sweetclover
Aster
Austrian Fieldcress
Bedstraw
Beggarticks
Betony, Florida
Bindweed, Field
Bird Vetch
Bitter Wintercress
Bittercress, Hairy
Bitterweed
Black-eyed Susan
Black Medic
Black Mustard
Blackseed Plantain
Blessed Thistle
Bloodflower Milkweed
Blue Lettuce
Blue Vervain
Bracted Plantain
Brassbuttons
Bristly Oxtongue
Broadleaf Dock
Broadleaf Plantain
Broomweed
Buckhorn
Buckhorn Plantain
Bulbous Buttercup
Bull Thistle
Bullnettle
Burclover
Burdock
Burning Nettle
Burweed
Buttercup
Buttonweed
Canada Thistle
Carolina Geranium
Carpetweed
Catchweed Bedstraw
Catnip
Catsear
Chickweed, Common
Chickweed, Mouseear
Chicory
Cinquefoil
Clover, Crimson
Clover, Hop
Clover, Red
Clover, Strawberry
Clover, Sweet & White
Cockle
Cocklebur
Common Mullein
Creeping Jenny
Cudweed
Curly Dock
Daisy, English
Daisy Fleabane
Daisy, Oxeye
Dandelion
Dichondra
Dogbane
Dogfennel
Dollarweed
Elderberry
False Dandelion
Falseflax
False Sunflower
Fiddleneck
Florida Pusley
Frenchweed
Galinsoga
Goathead
Goldenrod
Ground Ivy
Gumweed
Hairy Fleabane
Hawkweed
Healall
Heartleaf Drymary
Heath Aster
Hedge Bindweed
Hedge Mustard
Hemp
Henbit
Hoary Cress
Hoary Plantain
Hoary Vervain
Horsenettle
Jimsonweed
Knawel
Knotweed
Kochia
Lambsquarters
Lespedeza
Mallow
Matchweed
Mexicanweed
Milk Vetch
Morningglory
Mouseear Hawkweed
Mugwort
Musk Thistle
Mustard
Narrowleaf Plantain
Narrowleaf Vetch
Nettle
Orange Hawkweed
Oriental Cocklebur
Oxalis
Parsley-piert
Parsnip
Pearlwort
Pennycress
Pennywort
Pepperweed
Pigweed
Pineywoods Bedstraw
Plains Coreopsis (tickseed)
Plantain
Poison Ivy
Poison Oak
Pokeweed
Poorjoe
Prairie Sunflower
Prickly Lettuce
Prickly Sida
Prostrate Knotweed
Prostrate Pigweed
Prostrate Spurge
Prostrate Vervain
Puncturevine
Purslane, Common
Ragweed
Red Sorrel
Redroot Pigweed
Redstem Filaree
Rough Cinquefoil
Rough Fleabane
Russian Pigweed
Russian Thistle
Scarlet Pimpernel
Scotch Thistle
Sheep Sorrel
Shepherdspurse
Slender Plantain
Smallflower Galinsoga
Smooth Dock
Smooth Pigweed
Sorrel
Sowthistle
Spanishneedles
Speedwell
Spiny Amaranth
Spiny Cocklebur
Spiny Sowthistle
Spotted Catsear
Spotted Spurge
Spurweed
Stinging Nettle
Strawberry, India Mock
Tall Nettle
Tall Vervain
Tansy Ragwort
Tansy Mustard
Tanweed
Thistle
Trailing Crownvetch
Tumble Mustard
Tumble Pigweed
Velvetleaf
Venice Mallow
Virginia Buttonweed
Virginia Creeper
Virginia Pepperweed
Wavyleaf Bullthistle
Western Clematis
Western Salsify
White Mustard
Wild Aster
Wild Buckwheat
Wild Carrot
Wild Four-o'clock
Wild Garlic
Wild Geranium
Wild Lettuce
Wild Marigold
Wild Mustard
Wild Onion
Wild Parsnip
Wild Radish
Wild Rape
Wild Strawberry
Wild Sweet Potato
Wild Vetch
Woodsorrel
Woolly Croton
Woolly Morningglory
Woolly Plantain
Wormseed
Yarrow
Yellow Rocket
Yellowflower Pepperweed
And Other Broadleaf Weeds

When to Apply: Apply when broadleaf weeds are actively growing in the early fall.

STOP READ THE ENTIRE LABEL FIRST. OBSERVE ALL PRECAUTIONS AND FOLLOW DIRECTIONS CAREFULLY.

Directions for Use: It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Do not allow people (other than applicator) or pets on treatment area during application. Do not enter treatment areas until dust has settled. Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.

Where to Use: Expert Gardener® Winterizer Weed & Feed controls a wide range of lawn weeds including dandelion, chickweed, knotweed, plantain, henbit, spurge and many others commonly found in home lawns. In addition, the fertilizer provides a supply of plant food. Supplemental nitrogen, phosphorous, and potash should be applied according to soil tests, or State Extension Service recommendations or professional advice. Expert Gardener® Winterizer Weed & Feed controls broadleaf weeds commonly found in cool season and warm season turfgrass established in residential areas. Established turf of Kentucky bluegrass, fescues, and perennial ryegrass may be treated. Established turf of bermudagrass, bahiagrass, buffalograss, zoysiagrass, St. Augustinegrass and centipedegrass may be treated. Residential areas include turfgrass established around residences, apartment complexes, condominiums, vacant lots, or areas associated with households.

Use Precautions: Do not apply this product to carpetgrass, dichondra, ornamentals, or lawns containing desirable clovers. Choose a calm day to ensure uniform coverage and avoid spreading granules onto flowers, vegetables, and ornamental shrubbery. Do not exceed the application rate for any area and use care in the application of this product within the dripline of trees, shrubs, and other ornamentals. If overseeding wait 4 weeks after grass seeds sprout before applying this product.

Application:

1. Mow lawn to normal height 1 to 2 days before application.
2. Water lawn thoroughly at least 1 day before application to sustain moisture until the next watering (see step 4).
3. Apply when weeds are young and growing actively, preferably in the morning when dew is on the grass. If grass is not moist at time of application, sprinkle lightly with water to allow the granules to adhere and to remain on the leaf surface of the weeds.
4. Do not irrigate or sprinkle for 1 to 2 days after application. Irrigate, sprinkle, or water the lawn thoroughly at 2 to 3 days after application.

How Much To Apply: For Kentucky bluegrass, fescues, perennial ryegrass, bermudagrass, bahiagrass, buffalograss, zoysiagrass, St. Augustinegrass and centipedegrass: Apply 3.2 pounds of product per 1,000 square feet of turfgrass. A second or follow-up application may be required for dense infestations of weeds or for adverse seasonal conditions. The second application of 3.2 pounds of product per 1,000 square feet should be used 30 days after the first application. Cultivars vary in tolerance to this product. Do not apply this product to 'Floratam' St. Augustinegrass. The maximum application rate to turf is 1.1 pounds 2,4-D acid equivalent per acre per application per site. The maximum number of broadcast applications per treatment site is 2 per year.

Storage: Store in original container and keep any bags containing unused product in a locked storage area inaccessible to children and domestic animals.

Container Disposal: If empty: Do not reuse this container. Place in trash or offer for recycling if available. **If partly filled:** Call your local solid waste agency or 1-800-CLEANUP for disposal instructions. Never place unused product down any indoor or outdoor drain.

Distribution and Coverage: A uniform distribution of product on the treated area is essential. Choose a calm day to ensure uniform coverage and avoid spreading granules onto flowers, vegetables, and ornamental shrubbery. Do not exceed the application rate for any area and use care in the application of this product within the dripline of trees, shrubs, and other ornamentals.

Spreading Tips:

Apply header strips. Then apply product in longest direction of lawn. For best results, use a Scotts® spreader. For additional spreader settings, please call 1-800-543-TURF (8873). Calibrate your spreader before beginning your application and adjust the settings if necessary.

Spreader Settings	Covers 5,000 sq. ft.
Scotts Broadcast/Rotary Spreaders Each pass should be within 5 ft. of each other to ensure an overlap. **Scotts Basic™, Standard™, Deluxe™ EdgeGuard®, Lawn Pro™ or SpeedyGreen®**	4 ½
Scotts EasyGreen®	27 ½
Scotts Drop Spreaders Make sure spreader wheel completely overlaps previous wheel marks. **Scotts AccuGreen®**	7 ¾

Coverage: To determine how much you need to apply, multiply the length of your lawn by its width. This equals your total square footage. Deduct any non-lawn areas (house, garden, etc.)

What To Expect: Weed stems and leaves will begin to twist and curl in a few days, but a month or more may be required for complete control. This product will not prevent weed seeds from producing new plants. **Important:** It is important to follow directions as indicated because over application may cause injury to your lawn. Spreader settings are approximate. Differences may occur due to condition of spreader, speed at which applied and roughness of area treated.

Expert Gardener® Guarantee

When you choose Expert Gardener® products, you can be assured of high quality and performance in your lawn and garden. If for any reason you are not completely satisfied with this product, we will refund your money. Please mail your name, address and proof of purchase (dated receipt and package UPC/barcode) to the address below. Also, to ensure quality control, please write us a brief note explaining your experience with our product and reason for seeking a refund.

Distributed by The Scotts Company, 14111 Scottslawn Rd., Marysville, OH 43041

Contact Us

Consumer Helpline
1-866-369-2533

Information regarding the contents and levels of metals in this product is available on the Internet at:
http://www.regulatory-info-sc.com

Cuándo Aplicar: Aplique cuando las malezas de hoja ancha crezcan activamente a inicios de otoño.

STOP LEA PRIMERO TODA LA ETIQUETA. RESPETE TODAS LAS PRECAUCIONES Y SIGA MINUCIOSAMENTE LAS INSTRUCCIONES.

Instrucciones: Utilizar este producto de modo inconsistente con las declaraciones de su etiqueta constituye una violación de las leyes federales. No permita el ingreso de otras personas (distintas al aplicador) o mascotas a la zona de tratamiento durante la aplicación. No ingrese a las zonas de tratamiento hasta que se haya asentado el polvo. No se debe usar en tepe cultivado para la venta u otro uso comercial como cobertura, o para la producción comercial de semillas u otros fines de investigación.

Dónde Usar: El producto contra la mala hierba y fertilizante para el invierno Expert Gardener® controla una amplia variedad de malezas de césped, incluyendo el diente de león, estalaria, sanguinaria, llantén, ortiga mansa, lechetrezna y varias más que se encuentran comúnmente en el césped doméstico. Además, ofrece fertilizante a las plantas. El suplemento de nitrógeno, fósforo y potasa debe aplicarse de acuerdo con las pruebas del suelo, o las recomendaciones o consejo profesional del Servicio de Extensión del Estado. El producto contra la mala hierba y fertilizante para el invierno Expert Gardener® controla las malezas de hoja ancha que se encuentran comúnmente en el tepe de las estaciones fría y cálida establecido en las zonas residenciales. Se puede tratar el tepe establecido de pasto azul de Kentucky, festucas y raigrás perenne. Se puede tratar el tepe establecido de grama Bermuda, grama Bahía, grama búfalo, grama Zoysia, grama San Agustín y grama ciempiés. Las zonas residenciales incluyen al tepe establecido alrededor de las residencias, complejos de apartamentos, condominios, lotes vacantes o zonas asociadas con viviendas.

Precauciones De Uso: No aplique este producto a la grama alfombra, oreja de ratón, plantas ornamentales o césped que contenga trébol deseable. Elija un día calmado para asegurar la cobertura uniforme y evitar la diseminación de los gránulos en las flores, vegetales y arbustos ornamentales. No exceda la tasa de aplicación para ninguna zona y tenga cuidado con la aplicación de este producto dentro de la zona de goteo de los árboles, arbustos y otras plantas ornamentales. Si se excede en el sembrado, espere 4 semanas después de la germinación de las semillas de césped para aplicar este producto.

Aplicación:

1. Pode el césped a una altura normal 1 ó 2 días antes de la aplicación.
2. Riegue el césped totalmente por lo menos 1 día antes de la aplicación para sostener la humedad hasta el siguiente riego (véase el paso 4).
3. Aplique cuando las hierbas se encuentren jóvenes y crezcan activamente, de preferencia en la mañana cuando el rocío se encuentre en el césped. Si el césped no se encuentra húmedo en el momento de la aplicación, rocíe ligeramente agua para permitir que los gránulos se adhieran y permanezcan en la superficie de las hojas de las malezas.
4. No riegue ni rocíe nada durante 1 ó 2 días después de la aplicación. Riegue, rocíe o vierta agua al césped completamente a los 2 ó 3 días después de la aplicación.

Cómo Aplicar: Para el pasto azul de Kentucky, festucas, raigrás perenne, grama Bermuda, grama Bahía, grama Búfalo, grama Zoysia, grama San Agustín y grama ciempiés. Aplique 3.2 libras de producto por 1,000 pies cuadrados de tepe. Se puede necesitar una segunda aplicación para las infestaciones densas de malezas o en caso de condiciones adversas del clima. La segunda aplicación de 3.2 libras de producto por 1,000 pies cuadrados debe utilizarse 30 días después de la primera aplicación. Las variedades cultivadas varían su tolerancia a este producto. No aplique este producto en grama San Agustín "floratam". La tasa de aplicación máxima en el tepe es 1.1 libras de equivalente de ácido 2,4-D por acre por aplicación por zona. El número máximo de aplicaciones de siembra a voleo por zona de tratamiento es 2 al año.

Almacenamiento: Guarde en el envase original y conserve las bolsas que contengan el producto no utilizado en una zona de almacenamiento con cerradura inaccesible para los niños y animales domésticos.

Eliminación Del Envase: Si está vacío: No vuelva a utilizar este envase. Coloque en la basura u ofrezca para el reciclado si existe. **Si se encuentra parcialmente lleno:** Llame a su agencia local de desechos sólidos o al 1-800-CLEANUP para obtener instrucciones de eliminación. Nunca coloque el producto sin usar debajo de ningún drenaje interior o exterior.

Distribución Y Cobertura: Es esencial la distribución uniforme del producto en la zona tratada. Elija un día calmado para asegurar la cobertura uniforme y evitar la diseminación de los gránulos en las flores, vegetales y arbustos ornamentales. No exceda la tasa de aplicación para ninguna zona y tenga cuidado con la aplicación de este producto dentro de la zona de goteo de los árboles, arbustos y otras plantas ornamentales.

Cobertura: Para determinar cuánto necesita aplicar, multiplique la longitud y el ancho del jardín. Esto es igual al área total. Deduzca las zonas sin césped (casa, huerta, etc.)

Qué Esperar: Los tallos y las hojas de las malezas comenzarán a torcerse y curvarse en unos días, pero se puede necesitar un mes o más para obtener el control completo. Este producto no evitará que las semillas de malezas originen nuevas plantas. **IMPORTANTE:** Es importante seguir las instrucciones según se indiquen porque la aplicación en exceso puede dañar su césped. Los ajustes del esparcidor son aproximados. Pueden ocurrir diferencias debido a las condiciones del esparcidor, velocidad en la que se aplica y escabrosidad del área tratada.

CAUTION:
KEEP OUT OF REACH OF CHILDREN.
PRECAUTIONARY STATEMENTS: HAZARDS TO HUMANS & DOMESTIC ANIMALS.

CAUTION: Causes moderate eye irritation. Harmful if inhaled, or if absorbed through skin, or if swallowed. Avoid breathing dust. Avoid contact with eyes or clothing. When using this product, wear long-sleeved shirt, long pants, socks and shoes. After using this product, remove clothing and launder separately before reuse, and promptly and thoroughly wash hands and exposed skin with soap and water.

FIRST AID: ***If in eyes:*** Hold eye open and rinse slowly and gently with water for 15 - 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice. ***If on skin or clothing:*** Take off contaminated clothing. Rinse skin immediately with plenty of water for 15 - 20 minutes. Call a poison control center or doctor for treatment advice. ***If swallowed:*** Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor. Do not give anything by mouth to an unconscious person. Have the product container or label with you when calling a poison control center or doctor or going for treatment.

ENVIRONMENTAL HAZARDS: This product is toxic to aquatic invertebrates. Drift or runoff may adversely affect nontarget plants. Do not apply directly to water. Do not contaminate water when disposing of equipment washwater.

PRECAUCIÓN:
MANTENGA FUERA DEL ALCANCE DE LOS NIÑOS.
AVISOS DE PRECAUCIÓN: RIESGOS PARA LOS SERES HUMANOS Y ANIMALES DOMÉSTICOS.

PRECAUCIÓN: Causa irritación moderada a los ojos. Perjudicial si se inhala, ingiere o absorbe a través de la piel. Evite respirar el polvo. Evite el contacto con los ojos o la ropa. Cuando utilice este producto, use camisa de mangas largas, pantalones largos, medias y zapatos. Después de usar este producto, quítese la ropa y lávela por separado antes de volver a usarla, lávese de inmediato las manos y la piel expuesta completamente con agua y jabón.

PRIMEROS AUXILIOS: ***Si ocurre contacto con los ojos:*** Mantenga el ojo abierto y enjuague lenta y suavemente con agua durante 15 ó 20 minutos. Retire los lentes de contacto, si hubiera, después de los primeros 5 minutos, luego continúe enjuagando los ojos. Llame a un centro de control de envenenamiento o a un médico para recibir consejos de tratamiento. ***Si ocurre contacto con la piel o ropa;*** Quítese la ropa contaminada. Enjuague la piel inmediatamente con abundante agua durante 15 ó 20 minutos. Llame a un centro de control de envenenamiento o a un médico para recibir consejos de tratamiento. ***Si se ingiere:*** Llame de inmediato a un centro de control de envenenamiento o a un médico para recibir consejos de tratamiento. Haga que la persona beba un vaso de agua si puede tragar. No provoque el vómito a menos que el centro de control de envenenamiento o médico le indique hacerlo. No dé nada por la boca a una persona inconsciente. Tenga consigo el envase o etiqueta del producto cuando llame al centro de control de envenenamiento o al médico o acuda al tratamiento.

RIESGOS AMBIENTALES: Este producto es tóxico para los invertebrados acuáticos. La deriva y el escurrimiento pueden afectar adversamente a plantas que no se desee eliminar. No aplique directamente al agua. No contamine el agua con las aguas de lavado del equipo.

GUARANTEED ANALYSIS

Expert Gardener® Winterizer Weed & Feed 22-3-11 **F 643**

Total Nitrogen (N) 22%
1.2% ammoniacal nitrogen
20.8% urea nitrogen*
Available phosphate (P_2O_5) 3%
Soluble potash (K_2O) 11%
Sulfur (S) 1.9%
1.9% free sulfur (S)

Derived from: polymer-coated, sulfur-coated urea; urea, ammonium phosphate, potassium chloride.
* A portion of the urea has been coated to provide 4.9% coated slow-release nitrogen.

EPA REG. No. 2217–827–538
EPA Est. Nos. 9198–OH–1M, 8378–-IN–1K ,35512–FL–2H, 538–OH–1S.
Superscript used is first letter of run code on end of container.
CAS Registry Numbers: 2,4-D, DMA salt CAS#2008-39-1
MCPP-p, DMA salt CAS#66423-09-4
2,4-DP-p, DMA salt CAS#104786-87-0

Scotts® & Poly-S®, SpeedyGreen®, EdgeGuard®, EasyGreen®, AccuGreen® and PrecisionGreen® are registered trademarks of OMS Investments, Inc.
Basic™, Standard™ and Deluxe™ EdgeGuard® are trademarks of OMS Investments, Inc.

120%
12-digit UPC
(non-suppressed)
0 32247 37105 6

125050617

7-15-05

Expert Gardener® is Marketed Exclusively by Wal-Mart Stores, Inc., Bentonville, AR 72716
Shop @ www.walmart.com

© 2005

Product of U.S.A

U.S. Patent Nos.
5,219,465; 5,300,135; 5,405,426; 5,466,274