



SIPHOTROL® Plus II PREMISE SPRAY

with PRECOR® Insect Growth Regulator (IGR)

Siphotrol® Plus II Premise Spray (Siphotrol® Plus II) kills adult fleas, hatching flea eggs, flea larvae and ticks. PRECOR®, the unique ingredient in Siphotrol® Plus II, provides 30 weeks' protection by preventing eggs from ever developing into biting fleas. For best results, use Siphotrol® Plus II with Vet Kem® on-animal products for complete control of fleas and ticks.



9426.515

LICENSED

LICENSED

PERIOD 2014-2016 LIC.NO. PERIOD 2011-2013 LIC.NO.

Siphotrol®

Plus II

Department of Agriculture
STATE OF HAWAII

PREMISE SPRAY LICENSED

Contains PRECOR® Insect Growth Regulator

Plus adulticides

PERIOD 2008-2010 LIC.NO.

FAST-ACTING FORMULA

100% knock down for adult fleas in 10 minutes

For use in the home and garage to control fleas and ticks

Kills both adult and immature fleas & ticks

Treats 2,000 square feet

Leaves no lingering odor, no stains, no sticky mess

Prevents reinfestation and flea build-up for 30 weeks

Easy to apply water-based aerosol

ACTIVE INGREDIENTS:

(S)-Methoprene (CAS #65733-16-6)	0.085%
Permethrin (CAS #52645-53-1)	0.350%
Phenothrin (CAS #26002-80-2)	0.300%
N-octyl bicycloheptene dicarboximide (CAS #113-48-4)	2.000%
Piperonyl butoxide, Technical* (CAS #51-03-6)	1.400%
OTHER INGREDIENTS**:	95.865%
TOTAL	100.000%

* consists of 1.120% (butylcarbityl)

(6-propylpiperonyl) ether and 0.280% related compounds

** contains petroleum distillates

KEEP OUT OF REACH OF CHILDREN

CAUTION

See Back Panel for
Additional Precautionary Statements

NET CONTENTS: 16 OZ (454 g)

READ ENTIRE LABEL BEFORE USE. NOT FOR USE ON HUMANS OR PETS.

DIRECTIONS FOR USE: It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

SHAKE WELL BEFORE USING

FOR USE IN THE HOME/GARAGE/KENNEL

1. **Identify treatment areas**, especially those where pets frequent, such as in and around pet's sleeping and resting areas. These are the primary areas where fleas, their eggs, and ticks are found. Before treatment, make a test application to upholstery, drapery, light-colored, delicate fabrics, or wood surfaces in an inconspicuous place.
2. **Prepare treatment areas**. Cover all food processing surfaces and utensils during treatment or thoroughly wash after product use. Cover or remove exposed food. Remove pets and cover and unplug aquariums before spraying. Remove motor vehicles before using in garages. For best results with Surface/Carpet treatments, vacuum thoroughly before spraying and discard vacuum bag in outside trash. This product may also be used on infested surfaces inside motor vehicles.
3. **Apply treatment**. For flea/tick infestations, a single can treats 2,000 square feet. Hold container upside down 2-3 feet from surface, point actuator towards the surface, and push button. Using a sweeping motion, apply a light uniform spray to all surfaces of furniture, rugs, carpets, drapes, and around all pet resting areas. Avoid thoroughly wetting surfaces. Mist treated areas only until 'slightly damp'. Do not over-treat. Reapply in 14 days, if necessary. **IMPORTANT:** Only complete and proper application to carpets, upholstery, drapes, and other fabrics will provide thorough flea/tick elimination. Keep people and pets out of areas during treatment. Vacate room after treatment and ventilate until surfaces are dry before reoccupying.
4. For best results, eliminate fleas/ticks on your pet with appropriate Vet Kem® on-animal products.

STORAGE and DISPOSAL: Storage: Store in a cool, dry area. Do not transport or store below 32° F. Disposal: Do Not Puncture or Incinerate! If empty: Place in trash or offer for recycling if available. If partly filled: Call your local solid waste agency for disposal instructions.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS - CAUTION: Avoid contact with skin or clothing. Wash hands thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals.

FIRST AID:	
IF ON SKIN OR CLOTHING	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.
IF SWALLOWED	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Do not induce vomiting unless told to do so by a poison control center or doctor. • Do not give ANY liquid to the person. • Do not give anything by mouth to an unconscious person.
NOTE TO PHYSICIAN/VETERINARIAN: Contains petroleum distillates – vomiting may cause aspiration pneumonia.	
In case of emergency, call 1-800-766-7661 or a poison control center. Have the product container or label with you when calling a poison control center or doctor or going for treatment.	

PHYSICAL OR CHEMICAL HAZARDS: Contents under pressure. Do not use or store near heat or open flame. Do not puncture or incinerate container. Exposure to temperatures above 130° F may cause bursting.

Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and handling of this material when such use and handling are contrary to label instructions.

For product use information or in case of emergency, call 1-800-766-7661.

EPA Reg. No. 2724-490

© EPA Est. No. 11715-TN-1 © EPA Est. No. 11623-GA-1

Circled letter before EPA Est. No. corresponds to the first letter of the lot number on the bottom of the container.

Manufactured for:

Wellmark International
1501 East Woodfield Road 200W
Schaumburg, Illinois 60173

Precor, Siphotrol, VetKem and the VetKem logo are registered trademarks of Wellmark International.

Product Code: 3006302 RM#: 99.000020.00 08-1936

©2005-2008 Wellmark International Made in USA

www.vet-kem.com

