



Department of Agriculture
STATE OF HAWAII

LICENSED

PERIOD 2008-2010 LIC. NO.



Department of Agriculture
STATE OF HAWAII

LICENSED

PERIOD 2011-2013 LIC. NO.



Department of Agriculture
STATE OF HAWAII

LICENSED

PERIOD 2014-2016 LIC. NO.

9426.527



Siphotrol®

Contains PRECOR®
Insect Growth Regulator
Plus Permethrin,
an adulticide

Plus

FOGGER

Kills adult fleas and ticks

Stops hatching eggs and larvae from
developing into adult fleas for 30 weeks

Leaves no lingering odor, no stains,
no sticky mess

Prevents flea reinestation and flea
build-up for 30 weeks

Treats 6,000 cubic feet

**KEEP OUT OF
REACH OF CHILDREN
CAUTION**

See Back Panel for
Additional Precautionary
Statements

ACTIVE INGREDIENTS:	
(S)-Methoprene (CAS #65733-16-6)	0.09%
Permethrin (CAS #52645-53-1)	0.58%
OTHER INGREDIENTS:	99.33%
TOTAL	100.00%

NET CONTENTS: 6 OZ (170 g)

Siphotrol® *Plus* Fogger



Siphotrol® *Plus* Fogger kills adult fleas, hatching flea eggs, and ticks. PRECOR®, the ingredient in the Siphotrol® *Plus* Fogger, provides 30 weeks' protection by preventing eggs from ever developing into biting fleas. This product enables you to fill an entire room with insect-killing fog while you are away. For best results, use Siphotrol® *Plus* Fogger together with other Vet-Kem® products for complete control of fleas and ticks.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

In the home, all food processing surfaces and utensils should be covered during treatment, or thoroughly washed before use. Exposed food should be covered or removed.

SHAKE WELL BEFORE USING.

- Vacuum thoroughly before fogging. To ensure unobstructed flow of fogger mist, open inside doors and cabinets. Remove or cover exposed food, dishes, utensils, food preparation equipment, and surfaces. Remove pets and birds and remove or cover (and turn off) aquariums. Close outside doors and windows. **TURN OFF ALL IGNITION SOURCES SUCH AS PILOT LIGHTS (SHUT OFF GAS VALVES), OTHER OPEN FLAMES, OR RUNNING ELECTRICAL APPLIANCES THAT CYCLE OFF AND ON (i.e., REFRIGERATORS, THERMOSTATS, etc.). Call your gas utility or management company if you need assistance with your pilot lights.**
- Use one container for each 6000 cubic feet of area (a room 30' X 25' with an 8' ceiling). **DO NOT USE MORE THAN 1 FOGGER PER ROOM. DO NOT USE IN SMALL ENCLOSED SPACES SUCH AS CLOSETS, CABINETS, OR UNDER COUNTERS OR TABLES. DO NOT USE IN A ROOM 5' X 5' OR SMALLER; INSTEAD, ALLOW FOG TO ENTER FROM OTHER ROOMS.**
- Remove all motor vehicles before fogging garage.
- Elevate fogger by placing on stand or table covered by newspaper in center of room to be fogged. Newspapers should be spread on the floor for several feet around area of release.
- Before activating, tilt can away from face (container sprays straight up). Press down firmly on edge of actuator tab until it locks into place.
- After activating fogger, leave room or house at once. Keep treated area closed for at least 2 hours, then open all doors and windows and allow treated area to air for 1 hour before re-occupancy.
- For best results, spot treat with a Vet-Kem® premise product under beds, furniture, or obstructed areas where fogger mist may not penetrate.

STORAGE AND DISPOSAL: STORAGE: Store in a cool, dry area away from heat or open flame. **DISPOSAL:** Do Not Puncture or Incinerate! **Empty:** Place in trash or offer for recycling if available. **If partly filled:** Call your local solid waste agency for disposal instructions.

PRECAUTIONARY STATEMENTS - HAZARDS TO HUMANS AND DOMESTIC ANIMALS - CAUTION: USE ONLY WHEN AREA TO BE TREATED IS VACATED BY HUMANS AND PETS.

Harmful if inhaled, swallowed, or absorbed through the skin. Avoid breathing vapors or spray mist. May cause eye irritation. Avoid contact with eyes, skin, and clothing. Wash thoroughly with soap and water after using. Avoid contamination of food, foodstuffs, and food preparation areas. Leave room or house at once after activating fogger. If pet is exposed to wet surface, wash fur with soap and water.

FIRST AID: Call a poison control center or doctor immediately for treatment advice.

If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.

If inhaled: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible.

If swallowed: Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor. Do not give anything by mouth to an unconscious person.

If on skin or clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-766-7661 for emergency medical treatment information. Contact veterinarian if pets are exposed during application.

Note to Physician/Veterinarian: May present aspiration hazard.

Highly Flammable
Ingredient  Ingrediente Altamente
Inflamable

PHYSICAL OR CHEMICAL HAZARDS: Extremely Flammable. Contents under pressure. Keep away from heat, sparks, and open flame. Do not puncture or incinerate container. Exposure to temperatures above 130°F may cause bursting. **THIS PRODUCT CONTAINS A HIGHLY FLAMMABLE INGREDIENT. IT MAY CAUSE A FIRE OR EXPLOSION IF NOT USED PROPERLY. FOLLOW THE "DIRECTIONS FOR USE" ON THIS LABEL VERY CAREFULLY.**

In case of emergency or for product use information, call 1-800-766-7661.

Wellmark International
1501 East Woodfield Road 200W
Schramburg, Illinois 60173

Precor, Siphotrol, Vet-Kem and the Vet-Kem logo
are registered trademarks of Wellmark International.
©2003, 2008 Wellmark International

www.vet-kem.com

EPA Reg. No. 2724-454
EPA Est. No. 11623-GA-1

Made in USA
RM# 300502296 08-0988



0 4 1535 60460 8