STATE OF HAWAII
Department of Agriculture
ACCEPTED
LICENSE NO.

9426.548

DISC.

Do Not Varnish This Area

ZODIAC®
Flea & Tick Collar
for Cats & Kittens

READ ENTIRE LABEL BEFORE EACH USE. USE ONLY ON CATS AND KITTENS 12 WEEKS OF AGE AND OLDER.

DIRECTIONS FOR USE:
It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.
1. Buckle collar around cat's neck. Fit collar snug enough to prevent placement of the cat's leg or lower jaw through the collar, yet loose enough to be rotated about the neck. **2.** Leave 1-2 extra inches on the length of the collar for adjustment, cut off excess, and dispose of in trash. **3.** Check periodically and adjust fit if necessary, especially when kittens are rapidly growing. **4.** Replace in 8 months for continuous protection from adult fleas and flea egg hatch. Replace in 6 months for continuous protection from ticks. Collar does not need replacement if cat becomes wet.
To minimize flea infestation, use appropriate products to: Treat the pet; Treat the indoor environment, especially where pets spend time resting; Treat the outdoor environment when necessary.
STORAGE and DISPOSAL: Collar should be left in place once fitted on cat. If collar is temporarily removed, do not set it on wood or other surfaces. **Storage:** Store in original unopened package away from children. **Disposal:** Do not reuse container or reuse collar. Dispose of in trash.
WHAT TO EXPECT:
For Fleas: 1. This ZODIAC® collar is designed to kill adult fleas and flea eggs. **2.** This collar begins working to kill fleas and flea eggs as soon as it is placed around the cat's neck. **3.** Protection against fleas and flea eggs will be achieved within 48 hours, providing continuous protection for 8 months. **4.** Replace collar every 8 months to help control adult flea infestations on your pet.
For Ticks: 1. Full protection against adult and immature ticks will be achieved within a few days, providing continuous protection for up to 6 months. **2.** Adult ticks will be killed over the entire pet. Dead ticks will fall off or may be easily removed. **3.** Replace collar every 6 months for continuous protection from ticks.
PRECAUTIONARY STATEMENTS - HAZARDS TO HUMANS AND DOMESTIC ANIMALS - CAUTION: Do not get dust or collar in mouth. Harmful if swallowed or absorbed through skin. Do not get dust in eyes, will cause temporary constriction of the pupils*. The dust released by this collar is a cholinesterase inhibitor. Avoid contact with skin, eyes, or clothing.
HUMANS: Do not open package until ready to use. Do not allow children to play with this collar. Pesticidal dust will form on this collar during storage. Wash hands thoroughly with soap and water after handling collar. **ANIMALS:** Consult a veterinarian before using this product on debilitated, aged, medicated, pregnant, or nursing cats. Do not use on kittens under 12 weeks of age. Do not use this product on cats simultaneously or within 30 days before or after treatment with or exposure to cholinesterase inhibiting** drugs or pesticides (e.g. carbaryl, propoxur, etc.). However, flea and tick collars may be immediately replaced. When collar is first worn, observe neck area every few days for irritation. Remove collar at the first sign of irritation or adverse reaction. Sensitivities may occur after using ANY pesticide product for pets. If signs of sensitivity occur, remove collar and bathe your cat with mild soap and rinse with large amounts of water.

FIRST AID	
If swallowed	• Call poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by the poison control center or doctor. • Do not give anything by mouth to an unconscious person.
If on skin or clothing	• Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes • Call poison control center or doctor for treatment advice.
If in eyes	• Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call poison control center or doctor for treatment advice.
IN CASE OF EMERGENCY, CALL 1-800-950-4783 OR A POISON CONTROL CENTER. Have the product container or label with you when calling a poison control center or doctor or going for treatment.	
NOTE TO PHYSICIAN/VETERINARIAN: *Atropine is antidotal only if symptoms of cholinesterase inhibition are present.	

Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risks of use and handling of this material when such use and handling are contrary to label instructions.

Satisfaction guaranteed. For product use information, call 1-800-950-4783 or visit our Web site: **www.zodiacpet.com**.

** For information on cholinesterase inhibition, please, call our 800 number or consult your veterinarian.

ZODIAC® is a registered trademark of Wellmark International.
©2003 Wellmark International Made in USA

EPA Reg. No. 2724-491
EPA Est. No. 2724-TX-1
46207

Wellmark
Schaumburg, Illinois 60173

Do Not Varnish This Area

ZODIAC® Flea & Tick Collar *for Cats & Kittens*

Do Not Varnish This Area

ZODIAC®
Flea & Tick Collar
for Cats & Kittens

- Kills Fleas & Ticks *(including those carrying Lyme disease)*
- Kills Flea Eggs *(to prevent reinfestation)*

DO NOT ALLOW CHILDREN TO PLAY WITH THIS COLLAR
CAUTION
See Precautions on Back
NET WEIGHT: .37 OZ (10.5g)

ACTIVE INGREDIENTS:
Propoxur (CAS #114-26-1).....10.0%
(S)-Methoprene (CAS # 65733-16-6).....2.1%
OTHER INGREDIENTS:.....87.9%
TOTAL:.....100.0%

ZODIAC® Flea & Tick Collar *for Cats & Kittens*

Do Not Varnish This Area

0 41535 46200 0

Do Not Varnish This Area
Lot Code Prints Here

Do Not Varnish This Area