



STATE OF HAWAII
Department of Agriculture

ACCEPTED

LICENSE NO.

9426.557

Paramite® L.A.

INSECTICIDAL SPRAY & BACKRUBBER FOR CATTLE & SWINE

For use on cattle and swine only.
NOT for residential use on companion animals.

- Controls horn flies, lice, sarcoptic mange, and ticks on cattle
- Controls lice and sarcoptic mange on swine

ACTIVE INGREDIENT:

Phosmet (CAS# 732-11-6) 11.75%

OTHER INGREDIENTS* 88.25%

TOTAL 100.00%

* Contains petroleum distillates

Contains 1 lb active ingredient per gallon

KEEP OUT OF REACH OF CHILDREN

WARNING

SEE OTHER PRECAUTIONS/FIRST AID ON BACK PANELS.

READ ALL DIRECTIONS BEFORE USING.

1 QT (32 FL OZ) 946 ml

Product Code: 61830 RM#: 300504510 09-1074

2.5625"	2.5625"	2.6875"																															
<div style="border: 1px solid black; padding: 5px;"> <p style="text-align:center;">CATTLE BACKRUBBER</p> <table border="1" style="width:100%; border-collapse: collapse;"> <thead> <tr> <th rowspan="2"></th> <th colspan="2">DILUTION RATE</th> </tr> <tr> <th>Paramite® L.A.</th> <th>Fuel Oil or Other Suitable Carrier*</th> </tr> </thead> <tbody> <tr> <td>To Control Horn Flies Use fuel oil or other suitable carrier*</td> <td>1 gallon</td> <td>50.00 gallons</td> </tr> <tr> <td>Charge backrubber device or sack or cloth, as required.</td> <td>1/2 gallon</td> <td>25.00 gallons</td> </tr> <tr> <td>Retreat backrubber, as needed.</td> <td>1 quart</td> <td>12.50 gallons</td> </tr> <tr> <td>*Other Suitable Carriers – Diesel oil, Kerosene</td> <td>1 pint</td> <td>6.25 gallons</td> </tr> <tr> <td></td> <td>1 cup</td> <td>3.10 gallons</td> </tr> </tbody> </table> <p>CATTLE/WARNINGS:</p> <ol style="list-style-type: none"> 1. Paramite® L.A. is a cholinesterase inhibitor. Do not use this product on animals simultaneously or within a few days before or after treatment with or exposure to cholinesterase inhibiting drugs, pesticides, or chemicals. Atropine is antidotal. Consult veterinarian at first sign of adverse reaction. 2. Do not treat sick, convalescent, stressed, or animals less than 3 months old. 3. CATTLE MAY BE SLAUGHTERED 3 DAYS AFTER TREATMENT. 4. Do not treat non-lactating dairy cattle within 28 days of freshening. If freshening should occur within the 28 day period after treatment, that milk must not be used as human food. 5. IMPORTANT: In cattle, repeat treatment as necessary, but not more often than every 7 to 10 days. Treatment for lice, ticks, sarcoptic mange, and horn flies may be made any time of the year except when cattle grub larvae are in the gullet or spinal canal. Consult your veterinarian, extension livestock specialist, or extension entomologist regarding timing of treatment. </div>		DILUTION RATE		Paramite® L.A.	Fuel Oil or Other Suitable Carrier*	To Control Horn Flies Use fuel oil or other suitable carrier*	1 gallon	50.00 gallons	Charge backrubber device or sack or cloth, as required.	1/2 gallon	25.00 gallons	Retreat backrubber, as needed.	1 quart	12.50 gallons	*Other Suitable Carriers – Diesel oil, Kerosene	1 pint	6.25 gallons		1 cup	3.10 gallons	<div style="border: 1px solid black; padding: 5px;"> <p style="text-align:center;">SWINE SPRAY</p> <table border="1" style="width:100%; border-collapse: collapse;"> <thead> <tr> <th rowspan="2"></th> <th colspan="3">DILUTION RATE</th> </tr> <tr> <th>1 gallon of Paramite® L.A. in 100 gallons of water</th> <th>1 quart of Paramite® L.A. in 25 gallons of water</th> <th>1 cup of Paramite® L.A. in 6.25 gallons of water</th> </tr> </thead> <tbody> <tr> <td>To Control Lice, Sarcoptic Mange</td> <td></td> <td></td> <td></td> </tr> </tbody> </table> <p>SWINE/WARNINGS:</p> <ol style="list-style-type: none"> 1. Paramite® L.A. is a cholinesterase inhibitor. Do not use this product on animals simultaneously or within a few days before or after treatment with or exposure to cholinesterase inhibiting drugs, pesticides, or chemicals. Atropine is antidotal. Consult veterinarian at first sign of adverse reaction. 2. Do not treat sick, convalescent, or stressed animals. Do not apply directly to suckling pigs. 3. SWINE MAY BE SLAUGHTERED 1 DAY AFTER TREATMENT. 4. In swine, single applications for lice and sarcoptic mange control are usually effective; however, should a second application be necessary, it may be made 14 days following first treatment. </div>		DILUTION RATE			1 gallon of Paramite® L.A. in 100 gallons of water	1 quart of Paramite® L.A. in 25 gallons of water	1 cup of Paramite® L.A. in 6.25 gallons of water	To Control Lice, Sarcoptic Mange				<div style="border: 1px solid black; padding: 5px;"> <p style="text-align:center;">VET-KEM®</p> <p style="text-align:right;">PEEL BACK</p> <h1 style="text-align:center;">Paramite® L.A.</h1> <p style="text-align:center;">INSECTICIDAL SPRAY & BACKRUBBER FOR CATTLE & SWINE (Paramite® L.A.)</p> <p>PRECAUTIONARY STATEMENTS – HAZARDS TO HUMANS AND DOMESTIC ANIMALS – WARNING: May be fatal if swallowed. Causes substantial but temporary eye injury. Harmful if absorbed through skin. Do not get in eyes or on clothing. Avoid contact with skin. Wear protective eyewear (goggles, face shield, or safety glasses), long-sleeved shirt, long pants, shoes and socks, and chemical-resistant gloves (barrier laminate or viton, selection category G).</p> <div style="text-align:right;">  <p>0 4 15 35 1 6 18 30 8</p> </div> <p>Wellmark International 1501 East Woodfield Road 200W Schaumburg, Illinois 60173</p> <p><small>Paramite, Vet-Kem and the Vet-Kem logo are registered trademarks of Wellmark International. Central Garden & Pet and its logo are trademarks of Central Garden & Pet Co.</small></p> <p><small>Made in USA ©2009 Wellmark International Product Code: 61830 RM#: 300504511 09-1074</small></p> <p style="text-align:right;">CENTRAL Garden & Pet</p> </div>
		DILUTION RATE																															
	Paramite® L.A.	Fuel Oil or Other Suitable Carrier*																															
To Control Horn Flies Use fuel oil or other suitable carrier*	1 gallon	50.00 gallons																															
Charge backrubber device or sack or cloth, as required.	1/2 gallon	25.00 gallons																															
Retreat backrubber, as needed.	1 quart	12.50 gallons																															
*Other Suitable Carriers – Diesel oil, Kerosene	1 pint	6.25 gallons																															
	1 cup	3.10 gallons																															
	DILUTION RATE																																
	1 gallon of Paramite® L.A. in 100 gallons of water	1 quart of Paramite® L.A. in 25 gallons of water	1 cup of Paramite® L.A. in 6.25 gallons of water																														
To Control Lice, Sarcoptic Mange																																	
Page 5	Page 6	Page 1																															
2.8125"	2.8125"	2.9375"																															
2.6875"	2.5625"	2.5625"																															

2.8125"	2.8125"	2.9375"
2.6875"	2.5625"	2.5625"

<div style="border: 1px solid black; padding: 5px;"> <p>FIRST AID</p> <p>If swallowed • Immediately call a poison control center or doctor. • Do not induce vomiting unless told to do so by a poison control center or doctor. • Do not give any liquid to the person. Do not give anything by mouth to an unconscious person.</p> <p>If in eyes • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eyes. • Call a poison control center or doctor for treatment advice.</p> <p>If on skin or clothing • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.</p> <p>Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-766-7661 for emergency medical treatment information.</p> <p>NOTE TO PHYSICIAN/VETERINARIAN: Probable mucosal damage may contraindicate the use of gastric lavage. Paramite® L.A. is an organophosphate insecticide and a cholinesterase inhibitor. If signs of cholinesterase inhibition are present, atropine is antidotal. 2-PAM is also antidotal and may be administered in conjunction with atropine. If ingested, do not induce vomiting. May present aspiration hazard. Usual symptoms of organophosphate poisoning in humans include: headache, blurred vision, weakness, nausea, discomfort in the chest, vomiting, abdominal cramps, diarrhea, salivation, sweating, and pin-point pupils. Usual symptoms of poisoning in animals include: salivation, labored breathing, gastrointestinal disturbance, tremors, staggering, and pin-point pupils.</p> </div>	<div style="border: 1px solid black; padding: 5px;"> <p>User Safety Recommendations:</p> <p>Users should:</p> <ul style="list-style-type: none"> • Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet. • Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. • Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing. <p>ENVIRONMENTAL HAZARDS: This pesticide is toxic to fish and aquatic invertebrates. Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water by cleaning of equipment or disposal of equipment washwaters or rinsate. Drift and runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. This chemical can contaminate surface water. Under some conditions, it may have a high potential for runoff into surface water after application. These include poorly draining or wet soils with readily visible slopes toward adjacent surface waters, frequently flooded areas, areas overlying extremely shallow groundwater, areas with in-field canals or ditches that drain to surface water, areas not separated from adjacent surface waters with vegetated filter strips, and areas overlying tile drainage systems that drain to surface water. Do not allow spray to drift from the application site and contact people, structures people occupy at any time and the associated property, parks and recreation areas, nontarget crops, aquatic and wetland areas, woodlands, pastures, rangelands, or</p> </div>	<div style="border: 1px solid black; padding: 5px;"> <p>animals. This product is highly toxic to bees exposed directly to residues. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area. Protective information may be obtained from your cooperative Agricultural Extension Service.</p> <p>PHYSICAL OR CHEMICAL HAZARDS: Do not use or store near heat or open flame. Protect from temperatures below 20° F.</p> <p>DIRECTIONS FOR USE: It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.</p> <p>Stir thoroughly. Apply fresh mixture as a coarse spray, taking care to wet skin, not just hair. Do not apply to point of run-off. Do not allow spray to drift.</p> <p>Dilute as shown in tables below.</p> <table border="1" style="width:100%; border-collapse: collapse;"> <thead> <tr> <th rowspan="2"></th> <th colspan="3">DILUTION RATE</th> </tr> <tr> <th>1 gallon of Paramite® L.A. in 100 gallons of water</th> <th>1 quart of Paramite® L.A. in 25 gallons of water</th> <th>1 cup of Paramite® L.A. in 6.25 gallons of water</th> </tr> </thead> <tbody> <tr> <td>To Control</td> <td></td> <td></td> <td></td> </tr> <tr> <td>Cattle Ticks, Southern Cattle Ticks</td> <td>240</td> <td>60</td> <td>15.0</td> </tr> <tr> <td>Horn Flies</td> <td>200</td> <td>50</td> <td>12.5</td> </tr> <tr> <td>Lice</td> <td>150</td> <td>38</td> <td>9.5</td> </tr> <tr> <td>Sarcoptic Mange, Winter Ticks, Lone Star Ticks, Gulf Coast Ear Ticks</td> <td>100</td> <td>25</td> <td>6.2</td> </tr> </tbody> </table> <p style="text-align:right;"><small>(continued on next page)</small></p> </div>		DILUTION RATE			1 gallon of Paramite® L.A. in 100 gallons of water	1 quart of Paramite® L.A. in 25 gallons of water	1 cup of Paramite® L.A. in 6.25 gallons of water	To Control				Cattle Ticks, Southern Cattle Ticks	240	60	15.0	Horn Flies	200	50	12.5	Lice	150	38	9.5	Sarcoptic Mange, Winter Ticks, Lone Star Ticks, Gulf Coast Ear Ticks	100	25	6.2
	DILUTION RATE																												
	1 gallon of Paramite® L.A. in 100 gallons of water	1 quart of Paramite® L.A. in 25 gallons of water	1 cup of Paramite® L.A. in 6.25 gallons of water																										
To Control																													
Cattle Ticks, Southern Cattle Ticks	240	60	15.0																										
Horn Flies	200	50	12.5																										
Lice	150	38	9.5																										
Sarcoptic Mange, Winter Ticks, Lone Star Ticks, Gulf Coast Ear Ticks	100	25	6.2																										
Page 2	Page 3	Page 4																											
2.9375"	2.8125"	2.8125"																											

STORAGE and DISPOSAL: Do not contaminate water, food, or feed by storage or disposal.

Pesticide Storage: Store in a cool, dry place. Always store pesticides in the original container. If a leaky container must be contained within another, mark the outer container to identify the contents. The storage areas must be dry, well-lit, and well-ventilated. Keep pesticide storage areas clean. Clean up spills promptly. Store pesticides away from food, pet food, feed, seeds, fertilizers, and veterinary supplies.

Pesticide Disposal - Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency or the Hazardous Waste Representative at the nearest EPA Regional Office for guidance.

Container Handling: - Nonrefillable container. Do not reuse or refill this container. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

To the extent consistent with applicable law, seller makes no warranty, express or implied, concerning the use of this product other than indicated on the label. To the extent consistent with applicable law, buyer assumes all risk of use and handling of this material when such use and handling are contrary to label instructions. For information, or in case of an emergency, call 1-800-766-7661 or visit our Web site: www.vet-kem.com

VET-KEM



Paramite® L.A.

INSECTICIDAL SPRAY & BACKRUBBER FOR CATTLE & SWINE

For use on cattle and swine only. NOT for residential use on companion animals.

- Controls horn flies, lice, sarcoptic mange, and ticks on cattle
- Controls lice and sarcoptic mange on swine

ACTIVE INGREDIENT:
Phosmet (CAS# 732-11-6) 11.75%
OTHER INGREDIENTS* 88.25%
TOTAL 100.00%

* Contains petroleum distillates
Contains 1 lb active ingredient per gallon

KEEP OUT OF REACH OF CHILDREN

WARNING

SEE OTHER PRECAUTIONS/FIRST AID ON BACK PANELS.
READ ALL DIRECTIONS BEFORE USING.

1 QT (32 FL OZ) 946 ml
Product Code: 61830 RM#: 300504511 09-1074

Back Page
3.4375"