



# FLYS-X<sup>®</sup>

The Horse World's Most Trusted Name<sup>®</sup>

READ ENTIRE LABEL BEFORE EACH USE

For use on Horses, Beef Cattle, Dairy Cattle, Pigs, Goats, Sheep, Donkeys, Llamas, Poultry, and Dogs

EFFECTIVE, FAST-ACTING, READY TO USE INSECTICIDE

## DIRECTIONS FOR USE:

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

**Entry Restrictions:**  
- Except when applying to dogs, foals, horses and livestock, do not allow others to enter treated area until sprays have dried.

**Application Restrictions:**  
- Except when applying to dogs, foals, horses and livestock, do not apply this product in a way that will contact workers or other persons either directly or through drift. Only protected handlers may be in the area during application.

- Do not apply using a high pressure hand-wand or hand-held fogger in enclosed spaces.

- Not for use in or around floor drains or sewers.

- Remove or cover exposed food and drinking water before application.

- Remove or cover dishes, utensils, food processing equipment and food preparation surfaces or wash them before use.

- Do not apply more than 1 time per day.

- Remove pets, birds and cover fish aquarium before surface spraying.

- Do not apply to animals less than 12 weeks of age.

- Do not apply this water-based spray on or around electrical equipment due to the possibility of shock hazard.

## STORAGE AND DISPOSAL

**Pesticide Storage:** Store in a cool area above 32°F and protect from freezing. Store in original container only in a locked storage area.

**Pesticide Disposal and Container Handling:** Refillable container. Refill this container with pesticide only. Do not reuse this container for any other purpose. If empty: Place in trash or offer for recycling if available. If partly filled: Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

Directions For Use continued on side panels

FIRST AID	
<b>If in eyes:</b>	<ul style="list-style-type: none"> <li>• Hold eye open and rinse slowly and gently with water for 15-20 minutes.</li> <li>• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-628-9653 for emergency medical treatment information.

## PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

**CAUTION:** Causes moderate eye irritation. Avoid contact with eyes or clothing. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals.

**Personal Protective Equipment (PPE)**  
Mixers, loaders, applicators, and other handlers must wear the following: - long-sleeve shirt - long pants - shoes plus socks

**User Safety Requirements**  
Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry. Discard clothing and other absorbent material that have been drenched or heavily contaminated with this product. Do not reuse them.

- User Safety Recommendations:**
- Users should wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
  - Users should remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
  - Users should remove PPE immediately after handling this product. As soon as possible, wash thoroughly and change into clean clothing.



**HORSE AND LIVESTOCK SPRAY:** Effective, fast-acting, ready to use insecticide. Apply with conventional sprayers or a cloth. To protect beef cattle, dairy cattle, horses, pigs, goats, sheep, donkeys, llamas, and other listed livestock from horn flies, house flies, mosquitos, and gnats, apply a light mist sufficient to wet the surface of the hair. To kill stable flies, horse flies and deer flies, apply at a rate of 2-4 ounces per adult animal sufficient to wet the hair thoroughly. Repeat treatment daily or at intervals necessary to give continued protection. To kill face flies, apply with a cloth dampened with the spray. To kill blood sucking lice, apply to the infested areas of the animal. Use a stiff brush to get the spray to the base of the hair. Repeat every 2-3 weeks as required.

**POULTRY INSECTICIDE USE:** To control poultry lice and exposed stages of Darkling Beetles, thoroughly spray roosts, walls, nests, and cages, followed by spraying the birds with a fine mist.

**DOG AND FOAL INSECTICIDE USE:** To kill flea and brown dog ticks on premises and on dogs. Effective, fast-acting insecticide.

**Surface spraying:** Remove dogs prior to treatment. Thoroughly spray infested areas, pet beds, resting quarters, nearby cracks and crevices, along and behind baseboards, moldings, windows and door frames and entire areas of floor covering. Fresh bedding should be placed in animal quarters following treatment.

**On Dogs and Foals:** Start spraying at the tail, moving the dispenser rapidly and making sure the animal's entire body is covered including the legs and under body. While spraying, fluff the hair so spray will penetrate to the skin. Do not spray into eyes or face. Avoid contact with eyes and genitalia. Sensitivities may occur after using ANY pesticide product on pets. If signs of sensitivity occur, bathe your pet with mild soap and rinse with large amounts of water. If signs continue, consult a veterinarian immediately. Consult a veterinarian before using this product on debilitated, aged, pregnant or nursing animals or animals on medication. During the flea season, retreat weekly if necessary.



# FLYS-X<sup>®</sup>

## READY TO USE INSECTICIDE (AQUEOUS)



## FOR MULTI-SPECIES USE\*\* KILLS: HORN FLIES, STABLE FLIES, FACE FLIES, DEER FLIES, MOSQUITOES, GNATS, AND BLOOD SUCKING LICE ON LIVESTOCK

\*\* See back label for specific species

**ACTIVE INGREDIENTS:**

Pyrethrins.....	0.1%
Piperonyl Butoxide*.....	1.0%
<b>OTHER INGREDIENTS:</b> .....	98.9%
<b>TOTAL:</b> .....	100.0%

\*(butylcarbonyl) (6-propylpiperonyl) ether and related compounds

**KEEP OUT OF REACH OF CHILDREN  
CAUTION**  
SEE BACK PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS

Net Contents:  
32 fl oz. (946 mL)



STATE OF HAWAII  
Department of Agriculture

**ACCEPTED**

LICENSE NO. 9464.13

**HOUSEHOLD SPRAY:** Effective, fast-acting ready to use spray for use in homes. Leaves no oily or objectionable residues. Use of this product in kitchens should be limited to periods when not in use.

**Indoor surface spraying:** For maximum results, a combination of spot treatment and surface spray is recommended. To kill cockroaches, palmetto bugs, ants, silverfish, spiders, crickets, clover mites, cheese mites, granary weevils, confused flour beetles, spider beetles, cigarette beetles, drugstore beetles, meal worms, grain mites and cadelles. Adjust spray to a coarse wet spray. Direct the spray into hiding places, cracks and crevices, around containers of stored foods, behind shelves and drawers. For silverfish, spray bookcases. For ants, spray trails, nests and points of entry. If surface application only is to be used, spray floors and other surfaces at the rate of 1 gallon to 750 square feet of area. Repeat application weekly if necessary.



### FOR USE ON DOGS

Distributed by:  
W.F. Young, Inc.  
302 Benton Drive  
E. Longmeadow, MA 01028  
©WFO  
RM 364460  
Printed in U.S.A.  
[www.absorbine.com](http://www.absorbine.com)  
EPA Est. No. 7401-TX-01  
EPA Reg. No. 1543-10