

NO COPY AREA: FOLD LINES: SAFE COPY AREA: TRIM DIMENSIONS ARE IN MILLIMETERS

Customer: Amvac
Schematic #: 3230Schem45
Description: 510 (face) x 150 (gusset) x 310 (cut-off)
Machine(s): pch 2
Date Created: 03/23/15

3005 State Road
Telford, PA 18969
215-721-4600

System USA, Inc.
Fres-co

Customer: Amvac
Schematic #: 3230Schem45
Description: WISDOM 25 LB
Bag Style: Special Information: H/V



WISDOM[®] LAWN GRANULAR

INSECTICIDE

For use to control Ants (including Imported Fire Ants), Mole Crickets and other insect pests on lawns in landscaped areas and perimeters around residential, institutional, public commercial and industrial buildings, parks, recreational areas, and athletic fields.

ACTIVE INGREDIENT: 0.2%
By Weight
Inert Ingredients: 99.8%
TOTAL: 100.0%

*Cis isomers 87% minimum; trans isomers 13% maximum.

KEEP OUT OF REACH OF CHILDREN
CAUTION

FIRST AID

If on skin: Wash with plenty of water for 15-20 minutes.
If in eyes: Flush with water for 15-20 minutes.
If swallowed: Call a poison control center or doctor for treatment advice.

EMERGENCY INFORMATION

Have the product container or label with you when calling a poison control center or doctor for treatment advice.

FOR THE FOLLOWING EMERGENCIES, PHONE 24 HOURS A DAY:
For Medical Emergencies, including toll-free, 1-800-421-4281
For Poisoning Emergencies, including toll-free, 1-800-421-4281
For Product Use Information, including toll-free, 1-800-421-4281

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Harmful if absorbed through skin. Avoid contact with skin, eyes, or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Remove and wash contaminated clothing before reuse.

ENVIRONMENTAL HAZARDS

This pesticide is extremely toxic to fish and aquatic invertebrates. Do not apply when weather conditions favor drift on treated areas. Drift and run-off from treated areas may be hazardous to aquatic organisms in neighboring areas.

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Harmful if absorbed through skin. Avoid contact with skin, eyes, or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Remove and wash contaminated clothing before reuse.

ENVIRONMENTAL HAZARDS

This pesticide is extremely toxic to fish and aquatic invertebrates. Do not apply when weather conditions favor drift on treated areas. Drift and run-off from treated areas may be hazardous to aquatic organisms in neighboring areas.

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Harmful if absorbed through skin. Avoid contact with skin, eyes, or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Remove and wash contaminated clothing before reuse.

ENVIRONMENTAL HAZARDS

This pesticide is extremely toxic to fish and aquatic invertebrates. Do not apply when weather conditions favor drift on treated areas. Drift and run-off from treated areas may be hazardous to aquatic organisms in neighboring areas.

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Harmful if absorbed through skin. Avoid contact with skin, eyes, or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Remove and wash contaminated clothing before reuse.

ENVIRONMENTAL HAZARDS

This pesticide is extremely toxic to fish and aquatic invertebrates. Do not apply when weather conditions favor drift on treated areas. Drift and run-off from treated areas may be hazardous to aquatic organisms in neighboring areas.

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Harmful if absorbed through skin. Avoid contact with skin, eyes, or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Remove and wash contaminated clothing before reuse.

ENVIRONMENTAL HAZARDS

This pesticide is extremely toxic to fish and aquatic invertebrates. Do not apply when weather conditions favor drift on treated areas. Drift and run-off from treated areas may be hazardous to aquatic organisms in neighboring areas.

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Harmful if absorbed through skin. Avoid contact with skin, eyes, or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Remove and wash contaminated clothing before reuse.

ENVIRONMENTAL HAZARDS

This pesticide is extremely toxic to fish and aquatic invertebrates. Do not apply when weather conditions favor drift on treated areas. Drift and run-off from treated areas may be hazardous to aquatic organisms in neighboring areas.

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Harmful if absorbed through skin. Avoid contact with skin, eyes, or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Remove and wash contaminated clothing before reuse.

ENVIRONMENTAL HAZARDS

This pesticide is extremely toxic to fish and aquatic inve...

INSECTICIDE

In New York State, do make a single repeat application of Wisdom Lawn Granular Insecticide. If there are signs of repeated insect activity, do not re-apply until two weeks after the first application.

COMMENTS

Antennae, Cutworms, and Sod Webworms: To ensure optimum control of these insects, apply 0.25 lb. of insect immediately after application to a large release from the granules. Applications should be limited to control adult weevils as they leave their overwintering sites and move into grass. The treatment generally begins when Forsythia is in full bloom, and concludes when flowering dogwood is in full bloom. Consult your State Cooperative Extension Service for more specific information regarding application timing.

INDUSTRIAL NON-AGRICULTURAL APPLICATIONS
(Including Industrial Sites, Parks, Ball Fields and Cemeteries)

BROADCAST SPOT APPLICATIONS: For the control of actively foraging ants (including Imported Fire Ants) broadcast uniformly 2.5 to 4.0 lbs. WISDOM LAWN GRANULAR INSECTICIDE per 1000 sq. ft. (2.2 to 0.4 lb. a.i./A). See comments section under Lawn & Perimeter Applications for additional rate directions.

CHINCH BUGS: Chinch bugs infest the base of grass plants and are other than the insecticide. Chinch bugs can be one of the most difficult pests to control in grasses and the higher application rates may be required to control populations that contain both nymphs and adults during the summer.

PAD MOUNTED ELECTRICAL SITES: To reduce or prevent ant mound building activity on electrical pads, uniformly distribute 12 to 15 lbs. of WISDOM LAWN GRANULAR INSECTICIDE per 1000 sq. ft. of mounding pads. When existing mounds are completely or partially filled with soil excavated by resident ants, remove any excess fill and level the remaining soil prior to application.

CHINCH BUGS: Chinch bugs infest the base of grass plants and are other than the insecticide. Chinch bugs can be one of the most difficult pests to control in grasses and the higher application rates may be required to control populations that contain both nymphs and adults during the summer.

NOTE: Chemical requirements appear to be more attractive to ants during the cooler months. For best results, treat in the Fall or early Spring.

CHINCH BUGS: Chinch bugs infest the base of grass plants and are other than the insecticide. Chinch bugs can be one of the most difficult pests to control in grasses and the higher application rates may be required to control populations that contain both nymphs and adults during the summer.

NOTE: These settings should be used as starting points when Broadcast Application.

CHINCH BUGS: Chinch bugs infest the base of grass plants and are other than the insecticide. Chinch bugs can be one of the most difficult pests to control in grasses and the higher application rates may be required to control populations that contain both nymphs and adults during the summer.

NOTE: These settings should be used as starting points when Broadcast Application.

CHINCH BUGS: Chinch bugs infest the base of grass plants and are other than the insecticide. Chinch bugs can be one of the most difficult pests to control in grasses and the higher application rates may be required to control populations that contain both nymphs and adults during the summer.

NOTE: These settings should be used as starting points when Broadcast Application.

CHINCH BUGS: Chinch bugs infest the base of grass plants and are other than the insecticide. Chinch bugs can be one of the most difficult pests to control in grasses and the higher application rates may be required to control populations that contain both nymphs and adults during the summer.

NOTE: These settings should be used as starting points when Broadcast Application.

CHINCH BUGS: Chinch bugs infest the base of grass plants and are other than the insecticide. Chinch bugs can be one of the most difficult pests to control in grasses and the higher application rates may be required to control populations that contain both nymphs and adults during the summer.

NOTE: These settings should be used as starting points when Broadcast Application.

CHINCH BUGS: Chinch bugs infest the base of grass plants and are other than the insecticide. Chinch bugs can be one of the most difficult pests to control in grasses and the higher application rates may be required to control populations that contain both nymphs and adults during the summer.

NOTE: These settings should be used as starting points when Broadcast Application.

CHINCH BUGS: Chinch bugs infest the base of grass plants and are other than the insecticide. Chinch bugs can be one of the most difficult pests to control in grasses and the higher application rates may be required to control populations that contain both nymphs and adults during the summer.

NOTE: These settings should be used as starting points when Broadcast Application.

CHINCH BUGS: Chinch bugs infest the base of grass plants and are other than the insecticide. Chinch bugs can be one of the most difficult pests to control in grasses and the higher application rates may be required to control populations that contain both nymphs and adults during the summer.

NOTE: These settings should be used as starting points when Broadcast Application.

CHINCH BUGS: Chinch bugs infest the base of grass plants and are other than the insecticide. Chinch bugs can be one of the most difficult pests to control in grasses and the higher application rates may be required to control populations that contain both nymphs and adults during the summer.

NOTE: These settings should be used as starting points when Broadcast Application.

CHINCH BUGS: Chinch bugs infest the base of grass plants and are other than the insecticide. Chinch bugs can be one of the most difficult pests to control in grasses and the higher application rates may be required to control populations that contain both nymphs and adults during the summer.

NOTE: These settings should be used as starting points when Broadcast Application.

CHINCH BUGS: Chinch bugs infest the base of grass plants and are other than the insecticide. Chinch bugs can be one of the most difficult pests to control in grasses and the higher application rates may be required to control populations that contain both nymphs and adults during the summer.

NOTE: These settings should be used as starting points when Broadcast Application.

CHINCH BUGS: Chinch bugs infest the base of grass plants and are other than the insecticide. Chinch bugs can be one of the most difficult pests to control in grasses and the higher application rates may be required to control populations that contain both nymphs and adults during the summer.

NOTE: These settings should be used as starting points when Broadcast Application.

CHINCH BUGS: Chinch bugs infest the base of grass plants and are other than the insecticide. Chinch bugs can be one of the most difficult pests to control in grasses and the higher application rates may be required to control populations that contain both nymphs and adults during the summer.

NOTE: These settings should be used as starting points when Broadcast Application.

CHINCH BUGS: Chinch bugs infest the base of grass plants and are other than the insecticide. Chinch bugs can be one of the most difficult pests to control in grasses and the higher application rates may be required to control populations that contain both nymphs and adults during the summer.

NOTE: These settings should be used as starting points when Broadcast Application.

CHINCH BUGS: Chinch bugs infest the base of grass plants and are other than the insecticide. Chinch bugs can be one of the most difficult pests to control in grasses and the higher application rates may be required to control populations that contain both nymphs and adults during the summer.

NOTE: These settings should be used as starting points when Broadcast Application.

CHINCH BUGS: Chinch bugs infest the base of grass plants and are other than the insecticide. Chinch bugs can be one of the most difficult pests to control in grasses and the higher application rates may be required to control populations that contain both nymphs and adults during the summer.

NOTE: These settings should be used as starting points when Broadcast Application.

CHINCH BUGS: Chinch bugs infest the base of grass plants and are other than the insecticide. Chinch bugs can be one of the most difficult pests to control in grasses and the higher application rates may be required to control populations that contain both nymphs and adults during the summer.

NOTE: These settings should be used as starting points when Broadcast Application.

CHINCH BUGS: Chinch bugs infest the base of grass plants and are other than the insecticide. Chinch bugs can be one of the most difficult pests to control in grasses and the higher application rates may be required to control populations that contain both nymphs and adults during the summer.

NOTE: These settings should be used as starting points when Broadcast Application.

CHINCH BUGS: Chinch bugs infest the base of grass plants and are other than the insecticide. Chinch bugs can be one of the most difficult pests to control in grasses and the higher application rates may be required to control populations that contain both nymphs and adults during the summer.

NOTE: These settings should be used as starting points when Broadcast Application.

CHINCH BUGS: Chinch bugs infest the base of grass plants and are other than the insecticide. Chinch bugs can be one of the most difficult pests to control in grasses and the higher application rates may be required to control populations that contain both nymphs and adults during the summer.

NOTE: These settings should be used as starting points when Broadcast Application.

CHINCH BUGS: Chinch bugs infest the base of grass plants and are other than the insecticide. Chinch bugs can be one of the most difficult pests to control in grasses and the higher application rates may be required to control populations that contain both nymphs and adults during the summer.

NOTE: These settings should be used as starting points when Broadcast Application.

CHINCH BUGS: Chinch bugs infest the base of grass plants and are other than the insecticide. Chinch bugs can be one of the most difficult pests to control in grasses and the higher application rates may be required to control populations that contain both nymphs and adults during the summer.

NOTE: These settings should be used as starting points when Broadcast Application.

CHINCH BUGS: Chinch bugs infest the base of grass plants and are other than the insecticide. Chinch bugs can be one of the most difficult pests to control in grasses and the higher application rates may be required to control populations that contain both nymphs and adults during the summer.

NOTE: These settings should be used as starting points when Broadcast Application.

CHINCH BUGS: Chinch bugs infest the base of grass plants and are other than the insecticide. Chinch bugs can be one of the most difficult pests to control in grasses and the higher application rates may be required to control populations that contain both nymphs and adults during the summer.

NOTE: These settings should be used as starting points when Broadcast Application.

CHINCH BUGS: Chinch bugs infest the base of grass plants and are other than the insecticide. Chinch bugs can be one of the most difficult pests to control in grasses and the higher application rates may be required to control populations that contain both nymphs and adults during the summer.

NOTE: These settings should be used as starting points when Broadcast Application.

CHINCH BUGS: Chinch bugs infest the base of grass plants and are other than the insecticide. Chinch bugs can be one of the most difficult pests to control in grasses and the higher application rates may be required to control populations that contain both nymphs and adults during the summer.

NOTE: These settings should be used as starting points when Broadcast Application.

CHINCH BUGS: Chinch bugs infest the base of grass plants and are other than the insecticide. Chinch bugs can be one of the most difficult pests to control in grasses and the higher application rates may be required to control populations that contain both nymphs and adults during the summer.

NOTE: These settings should be used as starting points when Broadcast Application.

CHINCH BUGS: Chinch bugs infest the base of grass plants and are other than the insecticide. Chinch bugs can be one of the most difficult pests to control in grasses and the higher application rates may be required to control populations that contain both nymphs and adults during the summer.

NOTE: These settings should be used as starting points when Broadcast Application.

CHINCH BUGS: Chinch bugs infest the base of grass plants and are other than the insecticide. Chinch bugs can be one of the most difficult pests to control in grasses and the higher application rates may be required to control populations that contain both nymphs and adults during the summer.

NOTE: These settings should be used as starting points when Broadcast Application.

CHINCH BUGS: Chinch bugs infest the base of grass plants and are other than the insecticide. Chinch bugs can be one of the most difficult pests to control in grasses and the higher application rates may be required to control populations that contain both nymphs and adults during the summer.

NOTE: These settings should be used as starting points when Broadcast Application.

CHINCH BUGS: Chinch bugs infest the base of grass plants and are other than the insecticide. Chinch bugs can be one of the most difficult pests to control in grasses and the higher application rates may be required to control populations that contain both nymphs and adults during the summer.

NOTE: These settings should be used as starting points when Broadcast Application.

CHINCH BUGS: Chinch bugs infest the base of grass plants and are other than the insecticide. Chinch bugs can be one of the most difficult pests to control in grasses and the higher application rates may be required to control populations that contain both nymphs and adults during the summer.

NOTE: These settings should be used as starting points when Broadcast Application.

CHINCH BUGS: Chinch bugs infest the base of grass plants and are other than the insecticide. Chinch bugs can be one of the most difficult pests to control in grasses and the higher application rates may be required to control populations that contain both nymphs and adults during the summer.

NOTE: These settings should be used as starting points when Broadcast Application.

CHINCH BUGS: Chinch bugs infest the base of grass plants and are other than the insecticide. Chinch bugs can be one of the most difficult pests to control in grasses and the higher application rates may be required to control populations that contain both nymphs and adults during the summer.

NOTE: These settings should be used as starting points when Broadcast Application.

CHINCH BUGS: Chinch bugs infest the base of grass plants and are other than the insecticide. Chinch bugs can be one of the most difficult pests to control in grasses and the higher application rates may be required to control populations that contain both nymphs and adults during the summer.

NOTE: These settings should be used as starting points when Broadcast Application.

CHINCH BUGS: Chinch bugs infest the base of grass plants and are other than the insecticide. Chinch bugs can be one of the most difficult pests to control in grasses and the higher application rates may be required to control populations that contain both nymphs and adults during the summer.

NOTE: These settings should be used as starting points when Broadcast Application.

CHINCH BUGS: Chinch bugs infest the base of grass plants and are other than the insecticide. Chinch bugs can be one of the most difficult pests to control in grasses and the higher application rates may be required to control populations that contain both nymphs and adults during the summer.

NOTE: These settings should be used as starting points when Broadcast Application.

NET CONTENTS: 25 LB



7 49394 00384 5

USA/12/S/15

54/4/12/S/15

USA/AA5832

FP0

280

JPP

7 49394 00384 5

INSECTICIDE

LAWN GRANULAR

WISDOM

Back Reads This Way

DATE / LOT CODE

FRECQ VALVE INFO

377.93

62.07

62.07

190

130

510

530

1060

1230

150

170

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

NET CONTENTS: 25 LB



7 49394 00384 5

USA/12/S/15

54/4/12/S/15

USA/AA5832

FP0

280

JPP

7 49394 00384 5

INSECTICIDE

LAWN GRANULAR

WISDOM

Back Reads This Way

DATE / LOT CODE

FRECQ VALVE INFO

377.93

62.07

62.07

190

130

510

530

1060

1230

150

170