



HORSE GARD™

LICENSED

9526.25

Disc.

LICENSED

PERIOD 2015-2017 LIC. NO.

PERIOD 2012-2014 LIC. NO.

PRECAUTIONARY STATEMENTS:

HAZARDS TO HUMANS AND DOMESTIC ANIMALS:

CAUTION:

Harmful if swallowed or absorbed through skin. Avoid breathing of vapors or spray mist. Avoid contact with skin and eyes. In case of contact, immediately flush eyes with plenty of water. Obtain medical attention if irritation persists. Do not apply directly to food. Before spraying in Medical Care Areas, remove all patients. After spraying, ventilate area for 2 hours before returning patients. Remove pets, birds and cover fish aquariums before spraying.

PHYSICAL OR CHEMICAL HAZARDS:

Extremely flammable. Contents under pressure. Keep away from heat, sparks and heated surfaces. Do not puncture or incinerate container. Exposure to temperatures above 130°F may cause bursting. Do not spray on plastic, painted or varnished surfaces or directly into electronic equipment such as radios, TVs, computers, etc.

FIRST AID:

IF SWALLOWED: Immediately call a poison control center or doctor. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give any liquid to the person. Do not give anything by mouth to an unconscious person. **IF INHALED:** Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. Call a poison control center or doctor for further treatment advice. **IF ON SKIN OR CLOTHING:** Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice. **IF IN EYES:** Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice. **HOT LINE NUMBER** - Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For emergency information, call 1-800-222-1222, 24 hours, 7 days a week. **NOTE TO PHYSICIAN:** Contains petroleum distillates – vomiting may cause aspiration pneumonia.

STORAGE & DISPOSAL

Do not contaminate water, food or feed by storage or disposal. **STORAGE:** Do not store near heat or open flame. Store in cool dry area away from children. **DISPOSAL:** Do Not Puncture or Incinerate! IF EMPTY: Place in trash or offer for recycling, if available. IF PARTLY FILLED: Call your local solid waste agency or -1-800-CLEANUP for disposal instructions.

EPA Reg. No. 10807-450 EPA Est. No. 10807-GA-1

For Additional Products and Information, visit us at
www.Country-Vet.com



CONTAINS NO CFCs OR OTHER OZONE DEPLETING SUBSTANCES.
FEDERAL REGULATIONS PROHIBIT CFC PROPELLANTS IN AEROSOLS.

Manufactured by
AMREP, INC.
990 Industrial Park Drive
Marietta, GA 30062
Made in U.S.A.

Part #34-9314CVA



Swatter™

INSECTICIDE & REPELLENT FOR HORSES



PYRETHRINS ARE MADE FROM CHRYSANTHEMUM FLOWERS

Kills & Repels: Mosquitoes, Flies, Fleas, Barn Flies, Deer Flies, Horn Flies, Stable Flies & Gnats

- Provides Rapid Control of Barn Pests
- 0.50% Pyrethrins for Quick Kill

ACTIVE INGREDIENTS

Pyrethrins	0.50%
*Piperonyl Butoxide, TECHNICAL	4.00%
OTHER INGREDIENTS.....	95.50%
Total	100.00%
*Equivalent to 3.2% (butylcarbityl) (6-propylpiperonyl) ether and 0.8% related compounds. Contains Petroleum Distillates	

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

See Back Panel for Additional Precautionary Statements.

NET WT 1 LB. (16 oz.) (453.6 g)

Barns, Stables, Dairy Farms, Animal Quarters, Hog & Swine Operations, Beef Cattle Operations, Milkrooms and Poultry Operations.

DIRECTIONS FOR USE:

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

FLYING INSECTS: Flies, Mosquitoes, Small Flying Moths, Wasps, Gnats – Calculate cubic feet of space to be treated. Close all doors and windows. Spray in sweeping motion to all parts of the room. Apply for 2-3 seconds per 1,000 cubic feet of space. Sweep up and destroy fallen insects. Repeat treatment as necessary.

CRAWLING INSECTS: Cockroaches, Ants, Spiders, Silverfish – General Volumetric Treatment: Open cabinets and doors in area to be treated. Shut off air conditioners, fans, and close doors and windows before spraying. Spray for 20 seconds per 1,000 cubic feet of space. Direct spray toward areas suspected of harboring insects. Spray into hiding places such as cracks and crevices, behind sinks, cabinets, along baseboards and floors, around drains and plumbing, hitting insects with spray whenever possible. Keep area closed for 15 minutes. Open and ventilate before reoccupying. Repeat treatment as necessary.

FOR INDIANMEAL MOTHS AND ANGOUMOIS GRAIN MOTHS: Close doors and windows. Direct spray into all parts of room especially around stored food product containers, pallets and darkened areas. Spray 7 to 10 seconds for each 1,000 cubic feet of space. Keep area closed for 15 minutes following treatment. Repeat application weekly or as needed.

FOR FLEAS: Spray sleeping quarters, bedding, floor and floor covering where pets are kept. Repeat as necessary. For best results, use with a recommended pet spray on pets. **NOTE,** when using this product in an enclosed area, such as milkroom, bar or stable, remove animals before spraying.

DAIRY FARM USE: Milking Parlor and Milkroom: Close all windows and doors. Direct spray upwards, spraying for 2-3 seconds per 1,000 cubic feet of space. Do not remain in treated areas. Ventilate after treatment. Cover milking utensils and milk to prevent contamination from spray or dead and falling insects. **ANIMAL USE:** Thoroughly spray entire animal from approximately 2 feet distance. Do not spray directly toward animal's eyes. Spot treat withers, shoulders, and back where saliva accumulates from head tossing.

BEEF CATTLE OPERATIONS: In barns, close all windows and doors. Spray at the rate of 2-3 seconds per 1,000 cubic feet of space. Do not remain in treated area. After 15 minutes, ventilate the area. Repeat application as necessary. **ANIMAL USE:** For animal application, thoroughly spray entire animal from approximately 2 feet distance. Do not spray directly toward animal's face or eyes. Repeat application daily or as necessary.

STANCHION BARN USE: Walk behind animals and direct spray over backs, allowing 5 seconds per animal. For most effective results, apply each morning.

HORSE BARN OR STABLES: In horse and pony barns, close all windows and doors. Spray at the rate of 2-3 seconds per 1,000 cubic feet of space. Do not remain in the treated area. After 15 minutes, ventilate the area. Repeat the application as necessary. **ANIMAL USE:** For animal application, lightly mist over back of horses and ponies from a distance of 2 feet. Do not spray directly toward animal's face or eyes. Repeat application daily or as necessary.

HOG AND SWINE OPERATIONS: In hog houses, close all doors and windows. Spray at the rate of 2-3 seconds per 1,000 cubic feet of space. Do not remain in treated area. After 15 minutes, ventilate the area. Repeat the application as necessary. **SWINE USE:** For animal application, direct spray over back of hogs and spray for 5 seconds per hog. Do not spray directly towards the animal's face or eyes. Repeat application daily or as necessary.

POULTRY OPERATIONS: In poultry houses, close all windows and doors. Spray roosts, walls, nests, and cages thoroughly. Spray at the rate of 2-3 seconds per 1,000 cubic feet of space. Do not remain in treated areas. After 15 minutes, ventilate the area. Treat daily or as necessary.