

STATE OF HAWAII  
Department of Agriculture

**ACCEPTED**

LICENSE NO. **9526.36**  
Disc.



**MISTY**®

# Glypho Kill 18

**ACTIVE INGREDIENT:**

Isopropylamine salt of glyphosate: ..... 18.0%

**INERT INGREDIENTS:** ..... 82.0%

**TOTAL:** ..... 100.0%

**THIS PRODUCT CONTAINS:**

1.61 lbs. per U.S. gallon of the active ingredient glyphosate in the form of its isopropylamine salt.

Equivalent to 1.15 lbs. per U.S. gallon of the acid, glyphosate.

EPA REG. NO. 2217-850-10807

EPA EST. NO. 1270-TX-1

KEEP OUT OF REACH OF CHILDREN.


## **CAUTION:**

See packet insert for additional Precautionary Statements and First Aid and Use Directions.

- **For nonselective post emergent control of unwanted vegetation**
- **Systemic weed and grass killer for spot treatment of unwanted weeds and grasses**
- **Kills roots and all in 2-4 weeks**
- **No residual activity in soil**
- **Controls cattails and many other aquatic weeds**

# **B00572**

**NET CONTENTS 1 GALLON (3.8 LITERS)**



# Glypho Kill 18

- **For nonselective post emergent control of unwanted vegetation**
- **Systemic weed and grass killer for spot treatment of unwanted weeds and grasses**
- **Kills roots and all in 2-4 weeks**
- **No residual activity in soil**
- **Controls cattails and many other aquatic weeds.**

**ACTIVE INGREDIENT:**

Isopropylamine salt of glyphosate: ..... 18.0%

**INERT INGREDIENTS:** ..... 82.0%

**TOTAL:** ..... 100.0%

**THIS PRODUCT CONTAINS:**

1.61 lbs. per U.S. gallon of the active ingredient glyphosate in the form of its isopropylamine salt.

Equivalent to 1.15 lbs. per U.S. gallon of the acid, glyphosate.

KEEP OUT OF REACH OF CHILDREN.

**CAUTION:**

EPA REG. NO. 2217-850-10807

EPA EST. NO. 1270-TX-1

See below for additional Precautionary Statements, First Aid, and Use Directions.

**NET CONTENTS: 1 GALLON (3.8 LITERS)**

**BO0572**

MADE IN U.S.A. FOR:



**READ THE ENTIRE LABEL FIRST. OBSERVE ALL PRECAUTIONS AND FOLLOW DIRECTIONS CAREFULLY.**

**PRECAUTIONARY STATEMENTS: HAZARDS TO HUMANS AND DOMESTIC ANIMALS:**

**CAUTION:** Causes moderate eye irritation. Harmful if absorbed through the skin. Harmful if swallowed. Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water after handling.

**FIRST AID: IF SWALLOWED:** Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor. Do not give anything by mouth to an unconscious person. **IF ON SKIN OR CLOTHING:** Take off contaminated clothing. Rinse skin immediately with plenty of water for 15 - 20 minutes. Call a poison control center or doctor for treatment advice. **IF INHALED:** Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. Call a poison control center or doctor for treatment advice. **IF IN EYES:** Hold eye open and rinse slowly and gently with water for 15 - 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice. Have the product container or label with you when calling a poison control center or doctor or going for treatment. In case of medical emergency involving this product, call toll-free 1-877-800-5556.

**DOMESTIC ANIMALS:** This product is considered to be relatively nontoxic to dogs and other domestic animals; however, ingestion of this product or large amounts of freshly sprayed vegetation may result in temporary gastrointestinal irritation (vomiting, diarrhea, colic, etc.) If such symptoms are observed, provide the animal with plenty of fluids to prevent dehydration. Call a veterinarian if symptoms persist for more than 24 hours.

**ENVIRONMENTAL HAZARDS:** Do not contaminate water when disposing of equipment washwater. Treatment of aquatic weeds can result in oxygen depletion or loss due to decomposition of dead plants. This oxygen loss can cause fish suffocation. In case of spill or leak, soak up with an absorbent and remove to a landfill.

**PHYSICAL OR CHEMICAL HAZARDS:** Combustible. Do not use or store near heat or open flame. Spray solutions of this product should be mixed, stored and applied using only stainless steel, aluminum, fiberglass, plastic or plastic-lined steel containers. DO NOT MIX, STORE OR APPLY THIS PRODUCT OR SPRAY SOLUTIONS OF THIS PRODUCT IN GALVANIZED STEEL

OR UNLINED STEEL (EXCEPT STAINLESS STEEL) CONTAINERS OR SPRAY TANKS. This product or spray solutions of this product react with such containers and tanks to produce hydrogen gas which may form a highly combustible gas mixture. This gas mixture could flash or explode causing serious personal injury, if ignited by open flame, spark, welder's torch, lighted cigarette or other ignition source.

**DIRECTIONS FOR USE:** It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

**GENERAL INFORMATION:** This product mixes readily with water and is intended to be applied as a foliar spray for the control or destruction of many herbaceous or woody plants. It may be applied through most standard industrial or field-type sprayers after dilution and thorough mixing with water in accordance with label instructions. This product moves through the plant from the point of foliage contact to and into the root system. Visible effects on most annual weeds occur within 2 to 4 days, but on most perennial weeds may not occur for 7 days or more. Extremely cool or cloudy weather following treatment may slow activity of this product and delay visual effects of control. Visible effects are gradual wilting and yellowing of the plant which advances to complete browning of above-ground growth and deterioration of underground plant parts. Unless otherwise specified on this label, delay application until vegetation has emerged and reached the growth stages described for control. Unemerged plants arising from unattached underground rhizomes or root stocks of perennials will not be affected by the herbicide and will continue to grow. For this reason, best control of most perennial weeds is obtained when treatment is made at late growth stages approaching maturity. Always use the higher rate of this product per acre within the recommended range when weed growth is heavy or dense. Do not treat weeds under poor growing conditions such as drought stress, disease or insect damage, as reduced weed control may result. Reduced results may also occur when treating weeds heavily covered with dust. Reduced control may result when applications are made to annual or perennial weeds that have been mowed, grazed, or cut, and have not been allowed to regrow to the recommended stage for treatment. Rainfall or irrigation occurring within 6 hours after application may reduce effectiveness. Heavy rainfall or irrigation within 2 hours after application may wash the chemical off the foliage and a repeat treatment may be required. This product does not provide residual weed control. For subsequent residual weed control, follow a label-approved herbicide program. Read and carefully observe the cautionary statements and all other information appearing on the labels of all herbicides used. Buyer and all users are responsible for all loss or damage in connection with the use or handling of mixtures of this product with herbicides or other materials that are not expressly recommended in this labeling. Mixing this product with herbicides or other materials not recommended on this label may result in reduced performance. For best results, spray coverage should be uniform and complete.

**USE PRECAUTIONS:** AVOID DRIFT. EXTREME CARE MUST BE USED WHEN APPLYING THIS PRODUCT TO PREVENT INJURY TO DESIRABLE PLANTS AND CROPS. Do not allow the herbicide solution to mist, drip, drift or splash onto desirable vegetation since minute quantities of this product can cause severe damage or destruction to the crop, plants, or other areas on which treatment was not intended. The likelihood of injury occurring from the use of this product is greatest when winds are gusty or in excess of 5 miles per hour or when other conditions, including lesser wind velocities, will allow spray drift to occur. When spraying, avoid combinations of pressure and nozzle types that will result in splatter or fine particles (mist) which are likely to drift. AVOID APPLYING AT EXCESSIVE SPEED OR PRESSURE. **NOTE:** Use of this product in any manner not consistent with this label may result in injury to persons, animals or crops, or other unintended consequences. Keep container closed to prevent spills and contamination.

**SPRAY PREPARATION AND TANK MIXTURES:** This product is a soluble concentrate that dissolves readily in clean water (free of mud or clay). Reduced results may occur if water containing soil is used, such as water from ponds or unlined ditches. Additional surfactants are not required for this formulation.

**This Product Alone With Water As Diluent:** (1) Determine spray volume needed for the treated area. Add one-half the required amount of clean water to the spray tank. (2) Measure the recommended amount of product and add this product slowly with agitation, and complete filling the tank with water. (3) Mix thoroughly and continue agitation while spraying.

**Sprayer Cleanup:** CLEAN THE ENTIRE SPRAYER AFTER APPLICATION OF THIS PRODUCT. Failure to clean the sprayer thoroughly may result in injury to desirable crops which are subsequently sprayed.

**APPLICATION EQUIPMENT AND TECHNIQUES:** Do not apply this product through any type of irrigation system. This product may be applied with the following application equipment: **Hand-held and high-volume spray equipment** - pump-up pressure sprayers, knapsack and backpack sprayers, sprayguns, handwands, lances and other hand-held and motorized spray equipment used to direct the spray onto weed foliage. Applications should be made on a spray-to-wet basis, spray coverage should be uniform and complete. Use only coarse sprays. **Ground equipment** - Power sprayers fitted with a boom or spray wand/gun may be used for broadcast applications and spot treatments. For best spray distribution and coverage, select a spray volume and delivery system that will ensure accurate and uniform coverage. Boom sprayers equipped with appropriate flat fan nozzles, tips, and screens are suitable for broadcast applications. Spray volumes of 3 - 30 gallons per acre with spray pressures adjusted to 20 - 40 psi are appropriate. Use higher spray volumes within the recommended range for dense weed populations. **Controlled droplet applicator (CDA)** - The rate of this product applied per acre by vehicle-mounted CDA equipment must not be less than the amount recommended in this label when applied by conventional broadcast equipment. For vehicle-mounted CDA equipment, apply 3 to 15 gallons of water per acre. **Do not apply as a fine mist. Use coarse sprays only.** Controlled droplet application equipment produces a spray pattern which is not easily visible. Extreme care must be exercised to avoid spray or drift contacting the foliage or any other green tissue of desirable vegetation, as damage or destruction may result. **Selective equipment** - Recirculating sprayers, shielded and hooded sprayers, wiper applicators and sponge bars. **Wiper applicators and sponge bars** - Wiper applicators are devices that physically wipe appropriate amounts of this product directly onto the weed. **Rope or sponge wick applicators** - Apply this product undiluted (100% solution) to weeds listed on this label. **Porous-Plastic Applicators** - Apply this product undiluted (100% solution) in porous-plastic wiper applicators.

**SPRAY PREPARATION FOR APPLICATIONS WITH HAND-HELD HIGH VOLUME EQUIPMENT:** Specific use recommendations for annuals, perennials, woody brush, and trees are presented in the section named WEEDS CONTROLLED. Identify the target weed species, note the weed growth stages and select the appropriate spray concentration presented in WEEDS CONTROLLED. Spray concentrations for application with high volume equipment made on a spray-to-wet basis range from 2.75-7.00% vol/vol. Use the lower spray concentrations in the range for annual weeds and the higher spray concentration for perennial and hard-to-control species. The spray preparation chart for applications on a spray-to-wet basis is shown below in Table 1.

Volume (gallons)	2.75%	3.50%	4.40%	5.25%	7.00%
1 gallon	3.50 fl.oz.	4.50 fl.oz	5.60 fl.oz	6.70 fl.oz	9.00 fl.oz
3 gallons	10.50 fl.oz.	13.50 fl.oz	16.80 fl.oz	20.10 fl.oz	27.00 fl.oz
25 gallons	0.69 gal.	0.88 gal.	1.10 gal.	1.31 gal.	1.75 gal.
50 gallons	1.38 gal.	1.75 gal.	2.20 gal.	2.63 gal.	3.50 gal.
100 gallons	2.75 gal.	3.50 gal.	4.40 gal.	5.25 gal.	7.00 gal.

When using application methods that result in less than complete coverage, use a 17.5% spray concentration for annual and perennial weeds and a 17.5 - 28.0% spray concentration for woody brush and trees. If a straight-stream nozzle is used, start the application at the top of the target vegetation and spray from top to bottom in a "zig-zag" motion. Ensure that at least 50% of the leaf surface is contacted by the spray. Small open-branched trees need only be treated from one side. If foliage is thick or there are multiple root sprouts, application must be made from several sides to ensure adequate coverage. The spray preparation chart for these applications is shown below in Table 2.

Volume (gallons)	17.50%	28.00%
1 gallon	22.40 fl.oz	35.80 fl.oz
25 gallons	4.38 gal.	7.00 gal.
50 gallons	8.75 gal.	14.00 gal.
100 gallons	17.50 gal.	28.00 gal.

**WEEDS CONTROLLED: ANNUAL WEEDS:** Apply to actively growing grass and broadleaf weeds. Allow at least 3 days after treatment before disturbing vegetation. After this period, weeds may be mowed, tilled or burned. To prevent seed production, applications should be made prior to seedhead formation. This product may be used as a broadcast treatment or spot treatment for annual weeds with any of the application equipment listed on this label in noncropland sites.

**Recommendations for Annual Weeds Only**

Application Spray	Concentration, vol/vol%	Dosage rate, gallons/A
Spot treatments, hand held high volume equipment	2.75%	_____
Broadcast treatments	_____	0.7 – 1.1

**Hand-Held High-Volume Application:** Use a 2.75% vol/vol solution of this product in water and apply to foliage of vegetation to be controlled. This

product will provide full or partial control of the following annual weeds when applied as directed. Annual weeds will continue to germinate from seed throughout the growing season. Repeat treatments will be necessary to control later germinating seeds. **Broadcast Application:** Generally, the lower application rates within the specified range will provide satisfactory control of sensitive weed species. The higher application rates within the specified range will be required for dense infestations, for adverse/extreme environmental conditions, or for weeds beyond the appropriate growth stages. Use 0.7 gallons of this product per acre when weeds are less than 6 inches tall. If weeds are greater than 6 inches tall, use 1.1 gallons of this product per acre per 100 gallons of spray solution. Spray volumes of 3 to 10 gallons per acre for ground applications are recommended. **WEED SPECIES:** Anoda, spurred; Barnyardgrass *Echinochloa crus-galli*; Bassia, fivehook *Bassia hirsutioides*; Bittercress; Black nightshade; Bluegrass, annual *Poa annua*; Bluegrass, bulbous *Poa bulbosa*; Brome, downy; Brome, Japanese; Buttercup *Ranunculus spp.*; Carolina geranium; Carpetweed; Cheatgrass; Cheeseweed *Malva parviflora*; Chervil; Chickweed; Chickweed, mouseear *Cerastium vulgatum*; Cocklebur *Xanthium strumarium*; Copperleaf hophornbeam; Corn speedwell; Crabgrass *Digitaria spp.*; Dwarf dandelion *Krigia cespitosa*; Eclipta; Falsedandelion; Falseflax, smallseed *Camelina microcarpa*; Fiddleneck *Amsinckia spp.*; Filaree; Flaxleaf fleabane *Conyza bonariensis*; Fleabane *Erigeron spp.*; Florida pusley; Foxtail *Setaria spp.*; Foxtail, Carolina *Alopecurus carolinianus*; Goosegrass; Groundsel, common *Senecio vulgaris*; Henbit; Horseweed/marestail *Conyza Canadensis*; Itchgrass; Johnsongrass, seedling; Knotweed; Kochia, *Kochia scoparia*; Lambsquarters, common *Chenopodium album*; Lettuce, prickly *Lactuca serriola*; Mayweed; Medusahead; Morningglory, *Ipomoea spp.*; Mustard, blue *Chorispora tenella*; Mustard, tansy *Descurainia pinnata*; Mustard, tumble *Sisymbrium altissimum*; Mustard, wild *Brassica kaber*; Oats, wild *Avena fatua*; Panicum, *Panicum spp.*; Pennycress, field *Thiapsi arvensis*; Pigweed, redroot *Amaranthus retroflexus*; Pigweed, smooth *Amaranthus hybridus*; Plains/Tickseed coreopsis; Puncturevine; Purslane, common; Ragweed, common *Ambrosia artemisiifolia*; Ragweed, giant *Ambrosia trifida*; Rocket, London *Sisymbrium irio*; Sandbur, field *Cenchrus spp.*; Shattercane *Sorghum bicolor*; Shepherdspurse *Capsella bursa-pastoris*; Sicklepod; Signalgrass, broadleaf *Brachiaria platyphylla*; Smartweed, ladythumb; Smartweed, Pennsylvania *Polygonum pensylvanicum*; Sowthistle, annual *Sonchus oleraceus*; Spanishneedles; Speedweed, purslane; Spurge, annual; Spurge, prostrate; Spurge, spotted; Spurry, umbrella *Holosteum umbellatum*; Starthistle, yellow; Stinkgrass *Eragrostis cilianensis Canadensis*; Sunflower *Helianthus annuus*; Teaweed/Prickly sida; Texas panicum; Thistle, Russian *Salsola kali*; Velvetleaf *Abutilon theophrasti*; Virginia copperleaf; Virginia pepperweed; Witchgrass *Panicum capillare*; Woolly cupgrass; Yellow rocket.

**PERENNIAL WEEDS:** This product may be used as a broadcast treatment or spot treatment for perennial weeds with any of the application equipment listed on this label in noncropland sites and aquatic sites.

**Recommendations for Perennial Weeds Only**

Application Spray	Concentration, vol/vol%	Dosage rate, gallons/A
Spot treatments, hand held high volume equipment	2.75 – 5.25%	_____
Broadcast treatments	_____	2.0 – 3.3

Generally, the lower application rates within the specified range will provide satisfactory control of sensitive weed species. The higher application rates within the specified range will be required for dense infestations, for adverse/extreme environmental conditions, or for weeds beyond the appropriate growth stages. Specific use recommendations for certain perennial weeds are presented in the following section and **Table 3**. Apply to actively growing perennial weeds. Best results are obtained when soil moisture is adequate for active weed growth. Unless otherwise directed, allow at least 7 days after application before disturbing vegetation. **NOTE:** If weeds have been mowed or tilled, do not treat until plants have resumed active growth and have reached the recommended stages. Fall treatments must be applied before a killing frost. Repeat treatments may be necessary to control weeds regenerating from underground parts or seed. When applied as recommended under the conditions described, **this product will provide full or partial control for the following perennial weeds:** Alligatorweed *Alternanthera philoxeroides*\*; Anisel/Fennel *Foeniculum vulgare*; Artichoke, Jerusalem *Helianthus tuberosus*; Bahiagrass, *Paspalum notatum*; Beachgrass, European; Bermudagrass, *Cynodon dactylon*; Bermudagrass, water (knotgrass) *Paspalum distichum*; Bindweed, field *Convolvulus arvensis*; Bluegrass, Kentucky *Poa pratensis*; Blueweed, Texas *Helianthus ciliaris*; Brackenfern, *Pteridium aquilinum*; Bromegrass, smooth *Bromus inermis*; Bursage, woolly-leaf; Canarygrass, reed *Phalaris arundinacea*; Cattail, *Typha spp.*; Clover, reed *Trifolium pratense*; Clover, white *Trifolium repens*; Cogongrass, *Imperata cylindrical*; Cordgrass, *Spartina app* †; Cutgrass, giant *Zizaniopsis milliciae*\*; Dalligrass, *Paspalum dilatatum*; Dandelion *Taraxacum officinale*; Dock, curly *Rumex crispus*; Dogbane, hemp *Apocynum cannabinum*; Fescue, *Festuca spp.*; Fescue, tall *Festuca arundinacea*; German ivy; Guineagrass *Panicum maximum*; Hemlock, poison *Conium maculatum*; Horsenettle *Solanum carolinense*; Horseradish *Armoracia rusticana*; Ice plant *Mesembryanthemum crystallinum*; Johnsongrass, *Sorghum halepense*; Kikuyugrass *Pennisetum clandestinum*; Knapweed *Centauria repens*; Lantana *Lantana camara*; Lespedeza, common *Lespedeza striata*; Lespedeza, serices *Lespedeza cuneata*; Lossesstrife, purple *Lythrum salicaria* †; Lotus, American *Nelumbo lutea*; Maidencane *Panicum hematiomum*; Milkweed *Asclepias spp.*; Muhly, wirestem *Muhlenbergia frondosa*; Mullein, common *Verbascum thapsus*; Napiergrass *Pennisetum purpureum*; Nightshade, *silverleaf Solanum elaeagnifolium*; Nutsedge, purple, *Cyperus rotundus*, yellow, *Cyperus esculentus*; Orchardgrass, *Dactylis glomerata*; Pampasgrass, *Cortaderia jubata*; Paragrass, *Brachiada mutica*; Pepperweed, perennial; Phragmites *Phragmites spp.* †; Quackgrass *Agropyron repens*; Redvine †; Reed, giant *Arundo donax*; Ryegrass, perennial *Lolium perenne*; Smartweed, swamp *Polygonum coccineum*; Spatterdock, *Nuphar luteum* †; Spurge, leafy †; Starthistle, yellow *Centauria soisitalis*; Sweet potato, wild *Ipomoea pandurata*\*; Thistle, artichoke *Cynara cardunculus*; Thistle, Canada; Timothy *Phleum pratense*; Torpedograss *Panicum repens*\*; Trumpetcreeper †; Tules, common *Scirpus acutus* †; Vaseygrass *Paspalum urvillei*; Velvetgrass *Holcus spp.*; Waterhyacinth *Eichhornia crassipes* †; Waterlettuce *Pistia stratiotes* †; Waterprimrose *Ludwigia spp.* †; Wheatgrass, western. (\*Partial control). (†Not for use in California).

**Table 3. Recommendations for Perennial Weeds, Dosage Rates, Restrictions, and Comments.**

**Alligatorweed:** Apply 2.6 gallons of this product per acre or apply a 4.4% solution with hand-held and high volume equipment. Apply when most of the target plants are in bloom. Repeat applications will be required to maintain such control. **Bermudagrass:** Apply 3.3 gallons of this product per acre as a broadcast spray, or as a 5.25% solution with hand-held equipment. Apply when target plants are actively growing and when seedheads appear. **Bindweed, field; silverleaf nightshade; Texas blueweed:** Apply 2.6 – 3.3 gallons of this product per acre as a broadcast spray, west of the Mississippi River and 2.0 – 2.6 gallons per acre, east of the Mississippi River. With hand-held and high volume equipment, use a 5.25% solution. Apply when target plants are actively growing and are at or beyond full bloom. **For silverleaf nightshade,** best results can be obtained when application is made after berries are formed. Do not treat when weeds are under drought stress. New leaf development indicates active growth. For best results apply in late summer or fall. **Brackenfern:** Apply 2.0 – 2.6 gallons of this product per acre as a broadcast spray or as a 2.75 – 3.5% solution with hand-held and high volume equipment. Apply to fully expanded fronds which are at least 18 inches long. **Cattail:** Apply 2.0 – 2.6 gallons of this product per acre as a broadcast spray or as a 2.75% solution with hand-held and high volume equipment. Apply when target plants are actively growing and are at or beyond the early to full bloom stage of growth. Best results are achieved when application is made during the summer or fall months. **Cogongrass:** Apply 2.0 – 3.3 gallons of this product per acre as a broadcast spray. Apply when cogongrass is at least 18 inches tall and actively growing in late summer or fall. Allow 7 or more days after application before tillage or mowing. Due to uneven stages of growth and the dense nature of vegetation preventing good spray coverage, repeat treatments may be necessary to maintain control. **Cordgrass (Not for use in California):** Apply 2.0 – 3.3 gallons of this product per acre as a broadcast spray or as a 3.5 - 7.0% solution with hand-held and high volume equipment. Schedule applications in order to allow 6 hours before treated plants are covered by tidewater. The presence of silt or debris on the cordgrass will reduce performance. It may be necessary to wash target plants before application to improve uptake of this product into the plant. **Cutgrass, giant:** Apply 2.6 gallons of this product per acre as a broadcast spray or as a 3.5% solution with hand-held and high volume equipment. Repeat applications will be required to maintain control, especially where vegetation is partially submerged in water. Allow regrowth to the 7 to 10 leaf stage before retreatment. **Dogbane, hemp; knapweed; horseradish:** Apply 2.6 gallons of this product per acre as a broadcast spray or as a 5.25% solution with hand-held and high volume equipment. Apply when target plants are actively growing, and most have reached the late bud to flower stage of growth. For best results apply in late summer or fall. **Fescue, tall:** Apply 2.0 gallons of this product per acre as a broadcast spray or as a 3.5% solution with hand-held and high volume equipment. Apply when target plants are actively growing, and most have reached the boot to head stage of growth. When applied prior to boot stage, control may be reduced. **Guineagrass:** Apply 2.0 gallons of this product per acre as a broadcast spray or as 2.75% solution with hand-held and high volume equipment. Apply when target plants are actively growing and when most have reached at least the 7-leaf stage of growth. **Johnsongrass; Kentucky bluegrass; smooth bromegrass; reed canarygrass; orchardgrass; perennial ryegrass; timothy; western wheatgrass:** Apply 1.3 – 2.0 gallons of this product per acre as a broadcast spray or as a 2.75% solution with hand-held and high volume equipment. Apply when target plants are actively growing, and most have reached the boot to head stage of growth. When applied prior to boot stage, control may be reduced. In fall, apply before plants have turned brown. **Lantana:** Apply this product as a 2.75 - 3.5% solution with hand-held and high volume equipment. Apply to actively growing lantana at or beyond the bloom stage of growth. Use the higher rate for plants that have reached the woody stage of growth. **Purple loosestrife (Not for use in California):** Apply 1.7 gallons of this product per acre as a broadcast spray or as a 3.5 – 5.25% solution with hand-held and high volume equipment. Treat when plants are actively growing at or beyond the bloom stage of growth. Best results are achieved when application is made during summer or fall months. Fall treatments must be applied before a killing frost. **Lotus, American:** Apply 1.7 gallons

of this product as a broadcast spray or as 2.75% solution with hand-held and high volume equipment. Treat when plants are actively growing at or beyond the bloom stage of growth. Best results are achieved when applications are made during summer or fall months. Fall treatments must be applied before a killing frost. Repeat treatments may be necessary to control regrowth from underground parts and seeds. **Maidencane; paragrass:** Apply 2.6 gallons of this product as a broadcast spray or as 2.75% solution with hand-held and high volume equipment. Sequential treatments will be required, especially to vegetation partially submerged in water. Under these conditions, allow for regrowth to the 7 to 10 leaf stage prior to retreatment. **Milkweed, common:** Apply 2.0 gallons of this product per acre as a broadcast spray or as a 5.25% solution with hand-held and high volume equipment. Apply when target plants are actively growing and have reached the late bud to flower stage of growth. **Nutsedge, purple, yellow:** Apply 2.0 gallons of this product per acre as a broadcast spray or as a 2.75% solution with hand-held and high volume equipment. Apply when target plants are in flower or when new nutlets can be found at rhizome tips. Nutlets which have not germinated will not be controlled and may germinate following treatment. Repeat treatments will be required for long-term control. **Pampasgrass:** Apply a 5.25% solution of this product with hand-held and high volume equipment when plants are actively growing. **Phragmites, common reed:** In Florida, apply 3.3 gallons per acre of this product as a broadcast spray, or as a 5.25% solution with hand-held and high volume equipment. In other areas of the US, apply 1.7 – 2.6 gallons per acre as a broadcast spray or as a 2.75% solution with hand-held and high volume equipment. For partial control of phragmites in Florida and the counties of other states bordering the Gulf of Mexico. For best results treat during late summer or fall months when plants are actively growing and in full bloom. Due to the dense nature of the vegetation which may prevent good spray coverage and uneven stages of growth, repeat treatments may be necessary to maintain control. Visual control symptoms will be slow to develop. **Quackgrass; Kikuyugrass; wirestem muhly:** Apply 1.3 – 2.0 gallons of this product per acre as a broadcast spray or as a 2.75% solution with hand-held and high volume equipment. Apply when most quackgrass or wirestem muhly is at least 8 inches high (3 to 4 leaf stage of growth) and actively growing. Allow 3 or more days after application before tillage. **Giant reed; iceplant:** For control of giant reed and iceplant, apply a 5.25% solution of this product with hand-held and high volume equipment when plants are actively growing. For giant reeds best results are obtained when application is made in late summer to fall. **Spatterdock (Not for use in California):** Apply 2.6 gallons of this product per acre as a broadcast spray or as a 2.75% solution with hand-held and high volume equipment. Apply when most of the plants are in full bloom. For best results apply in late summer or fall. **Wild sweet potato:** Apply as a 5.25% solution with hand-held and high volume equipment. Apply to actively growing weeds at or beyond the bloom stage of growth. Repeat applications will be required. Allow the plant to reach the recommended stage of growth before retreatment. **Thistle, Canada:** Apply 1.3 – 2.0 gallons of this product per acre as a broadcast spray or as a 5.25% solution with hand-held and high volume equipment. Apply when target plants are actively growing and are at or beyond the bud stage of growth. **Torpedograss:** Apply 2.6 – 3.3 gallons per acre of this product as a broadcast spray or as a 2.75 – 5.25% solution with hand-held and high volume equipment. Use the lower rates under terrestrial conditions, and the higher rates under partially submerged or floating-mat conditions. Repeat treatment will be required to maintain control. **Tules, common (Not for use in California):** Apply as a 5.25% solution with hand-held and high volume equipment. Apply this product with hand-held equipment. Apply to actively growing plants at or beyond the seedhead stage of growth. After application visual symptoms will be slow to appear and may not occur for 3 or more weeks. **Waterhyacinth (Not for use in California):** Apply 2.2 - 2.6 gallons of this product per acre as a broadcast spray or as a 2.75 – 3.5% solution with hand-held and high volume equipment. Apply when target plants are actively growing and at or beyond the early bloom stage of growth. After application, visual symptoms may require 3 or more weeks to appear with complete necrosis and decomposition usually occurring within 60 to 90 days. Use the higher rates when more rapid visual effects are required. **Waterlettuce (Not for use in California):** Apply as a 2.75 – 3.5% solution with hand-held and high volume equipment. Use higher rates where infestations are heavy. Best results are obtained from mid-summer through winter applications. Spring application may require retreatment. **Waterprimrose (Not for use in California):** Apply as a 2.75% solution with hand-held and high volume equipment. Apply this product using hand-held equipment to plants that are actively growing at or beyond the bloom stage of growth, but before fall color changes occur. Thorough coverage is necessary for best control. **Other perennials listed on this label:** Apply 2.0 – 3.3 gallons of this product per acre as a broadcast spray or as a 2.75 – 5.25% solution with hand-held and high volume equipment. Apply when target plants are actively growing and most have reached early head or early bud stage of growth.

**WOODY BRUSH AND TREES:** When applied as recommended under the conditions described, this product CONTROLS or PARTIALLY CONTROLS the following woody brush, herbaceous vines and trees: Alder, *Alnus spp.*; Ash *Fraxinus spp.*\*; Aspen, quaking *Populus tremuloides*; Beech; Birch, *Betula spp.*; Blackberry, *Rubus spp.*; Blackgum; Bracken; Broom: French *Cytisus monspessulanus*, Scotch *Cytisus scopadus*; Buckwheat, California *Edogonum fasciculatum*\*; Cascara pagrada *Rhamnus purshiana*\*; Catsclaw *Acacia greggii*\*; Ceanothus, *Ceanothus spp.*; Chamise, *Adenostoma fasciculatum*; Cherry: Bitter *Prunus emarginata*, Black *Prunus serotina*, Pin *Prunus pensylvanica*; Coyotebrush, *Baccharis consanguinea*; Creeper, Virginia *Parthenocissus quinquefolia*\*; Deerweed; Dewberry, *Rubus trivialis*; Dogwood, *Cornus spp.*; Elderberry, *Sambucus spp.*; Elm, *Ulmus spp.* ; Eucalyptus, bluegum *Eucalyptus spp.*; Hasardia *Haplopappus squamosus*\*; Hawthorn, *Crataegus spp.*; Hazel, *Corylus spp.*; Hickory, *Carya spp.*; Holly; Honeysuckle, *Lonicera spp.*; Hornbeam, American *Carpinus caroliniana*; Kudzu, *Pueraria lobata*; Locust, black *Robinia pseudoacacia*\*; Madrone; Manzanita, *Arctostaphylos spp.*; Maple: Red *Acer rubrum*\*, Sugar *Acer saccharum*, Vine *Acer circinatum*\*; Monkey Flower *Mimulus guttatus*\*; Oak: Black *Quercus yelutina*\*, Northern Pin *Quercus palustris*, Post *Quercus stellata*, Red *Quercus rubra*, Southern Red *Quercus falcate*, White *Quercus alba*\*; Peppertree, Brazilian *Schinus terebinthifolius*; Persimmon, *Diospyros spp.* ; Pine; Poison ivy, *Rhus radicans*; Poison oak, *Rhus toxicodendron*; Poplar, yellow (Tulip tree) *Liriodendron tulipifera*\*; *Prunus spp.*; Raspberry, *Rubus spp.*; Redbud, eastern *Cercis Canadensis*; Rose, multiflora *Rosa multiflora*; Russian olive *Elaeagnus angustifolia*; Sage, black and white *Salvia spp.*; Sagebrush, California *Artemisia californica*; Salmonberry, *Rubus spectabilis*; Saltcedar *Tamarix spp.*; Saltbush; Sea myrtle *Baccharis halimifolia*; *Sassafras spp.*; Sourwood *Oxydendrum arboreum*\*; Sumac: Poison *Rhus vernix*\*, Smooth *Rhus glabra*\*, Winged *Rhus copallina*\*; Sweetgum *Liquidambar styraciflua*; Swordfern *Polystichum munium*; Tallow tree, Chinese *Sapium sebiferum*; Thimbleberry, *Rubus parviflorus*; Tobacco, tree *Nicotiana glauca*\*; Toyon; Trumpetcreeper *Campsis radicans*; Waxmyrtle, southern *Myrica cerifera*\*; Willow *Salix spp.*; Yerba santa. (\*Partial control). **NOTE:** If brush has been mowed or tilled or trees have been cut, do not treat until regrowth has reached the recommended stages of growth. Apply this product when

plants are actively growing and, unless otherwise directed, after full leaf expansion. Use the higher rate for larger plants and/or dense areas of growth. On vines, use the higher rate for plants that have reached the woody stage of growth. Best results are obtained when application is made in late summer or fall after fruit formation. In arid areas, best results are obtained when application is made in the spring to early summer when brush species are at high moisture content and are flowering. This product may be used as a broadcast treatment or spot treatment for woody brush and trees with any of the application equipment listed on this label in noncropland and aquatic sites.

Recommendations for Woody Brush and Trees Only		
ApplicationSpray	Concentration, vol%/vol%	Dosage rate, gallons/A
Spot treatments, hand held high volume equipment as full cover spray.	2.75 – 7.0%	—
Spot treatments, hand held high volume equipment without full coverage.	17.5 – 28.0%	—
Broadcast treatments	—	1.3 – 3.3

Generally, the lower application rates within the specified range will provide satisfactory control of sensitive weed species. The higher application rates within the specified range will be required for dense infestations, for adverse/extreme environmental conditions, or for weeds beyond the appropriate growth stages. Ensure thorough coverage when using hand-held equipment. Symptoms may not appear prior to frost or senescence with fall treatments. Allow 7 or more days after application before tillage, mowing or removal. Repeat treatments may be necessary to control plants regenerating from underground parts or seed. Some autumn colors on undesirable deciduous species are acceptable provided no major leaf drop has occurred. Reduced performance may result if fall treatments are made following a frost. Applied as a 17.5 - 28.0% solution as a directed application as described in the HAND-HELD AND HIGH-VOLUME EQUIPMENT section, this product will control or partially control all the species listed in this section of this label. Use the higher rate for dense stands and larger woody brush and trees. Specific use recommendations for certain woody brush and trees are presented in Table 4.

**Table 4. Recommendations for Woody Brush and Trees, Dosage Rates, Precautions and Comments.**

**Alder; blackberry; dewberry; honeysuckle; post oak; raspberry:** Apply 2.0 – 2.6 gallons per acre of this product as a broadcast spray or as a 2.75 –4.4% solution with hand-held and high volume equipment. **Aspen, quaking; hawthorn; trumpetcreeper:** Apply 1.3 – 2.0 gallons of this product per acre as a broadcast spray or as a 2.75 –4.4% solution with hand-held and high volume equipment. **Birch; elderberry; hazel; salmonberry; thimbleberry:** Apply as a 2.75% solution with hand-held and high volume equipment. **Broom, French, Scotch:** Apply a 4.4 – 5.25% solution with hand-held and high volume equipment. **Buckwheat, California; hasardia; monkey flower, tobacco tree:** Apply as a 2.75 –5.25% solution with hand-held and high volume equipment. Thorough coverage of foliage is necessary for best results. **Castorbean:** Apply a 5.25% solution with hand-held and high volume equipment. **Catsclaw:** Apply as a 4.4 – 5.25% solution with hand-held and high volume equipment. For partial control, apply with hand-held equipment when at least 50% of the new leaves are fully developed. **Cherry, bitter, black, pin; oak, southern red; sweetgum; Prunus:** Apply 1.3 – 3.3 gallons of this product per acre as a broadcast spray or as a 3.5 – 5.25% solution with hand-held and high volume equipment. **Coyotebrush:** Apply as a 4.4 – 5.25% solution with hand-held and high volume equipment. Apply with hand-held equipment when at least 50% of the new leaves are fully developed. **Dogwood; hickory; saltcedar:** Apply 2.6 – 3.3 gallons of this product per acre as a broadcast spray or as a 3.5 – 7.0% solution with hand-held and high volume equipment. For partial control. **Eucalyptus, bluegum:** Apply as a 5.25% solution with hand-held and high volume equipment. For control of eucalyptus resprouts, apply this product with hand-held equipment when resprouts are 6 - 12 feet tall. Ensure complete coverage. Apply when plants are growing actively. Avoid application to drought-stressed plants. **Kudzu:** Apply 2.6 gallons of this product per acre as a broadcast spray or as a 5.25% solution with hand-held and high volume equipment. Repeat applications will be required to maintain control. **Maple, red:** Apply as a 2.75 – 4.4% solution with hand-held and high volume equipment. Apply with hand-held equipment when leaves are fully developed. **Maple, sugar; oak, northern pin, red:** Apply as a 2.75 – 4.4% solution with hand-held and high volume equipment. Apply with hand-held equipment when at least 50% of the new leaves are fully developed. **Peppertree, Brazilian (Holly, Florida); waxmyrtle, southern:** Apply as a 5.25% solution with hand-held and high volume equipment. For partial control. **Poison ivy; poison oak:** Apply 2.6 – 3.3 gallons of this product per acre as a broadcast spray or as a 5.25% solution with hand-held and high volume equipment. Repeat applications may be required to maintain control. Fall treatments must be applied before leaves lose green color. **Rose, multiflora:** Apply 1.3 gallons of this product per acre as a broadcast spray or as a 2.75% solution with hand-held and high volume equipment. Treatments should be made prior to leaf deterioration by leaf-feeding insects. **Sage, black; sagebrush, California; chamise; tallow tree, Chinese:** Apply as a 2.75% solution with hand-held and high volume equipment. Thorough coverage of foliage is necessary for best results. **Saltbush, sea myrtle:** Apply as a 3.5% solution with hand-held and high volume equipment. **Willow:** Apply 2.0 gallons of this product per acre as a broadcast spray or as a 2.75% solution with hand-held and high volume equipment. **Other woody brush and trees listed:** Apply 1.3 – 3.0 gallons of this product per acre as a broadcast spray or as a 2.75 – 5.25% solution with hand-held and high volume equipment. For partial control.

**AQUATIC AND NONCROP SITES: Aquatic sites and wetlands:** This product may be applied to emerged weeds, woody brush and trees in all bodies of fresh and brackish water which may be flowing, non-flowing, or transient. This includes lakes, rivers, streams, ponds, estuaries, rice levees, seeps, irrigation and drainage ditches, canals, reservoirs, wastewater treatment facilities, wildlife habitat restoration and management areas, and similar sites including the following: **Aquatic food crop:** Agricultural drainage systems, irrigation systems, lakes/ponds/reservoirs (with human or wildlife use) stream/rivers/channeled water, drainage ditches, canals, irrigation ditches, estuaries. **Aquatic non-food industrial:** aquatic areas/water, drainage systems, waste water systems. **Aquatic non-food outdoor:** aquatic areas/water, ditches, ditch banks. This product may be used in and around water and wetlands found near power, telephone and pipeline rights-of-way sites, including where these sites are adjacent to or surrounding domestic water supply reservoirs, supply streams, lakes and ponds. If aquatic sites are present in the non-crop

area and are part of the intended treatment, read and observe the following directions: **This product does not control plants which are completely submerged or have a majority of foliage under water.** There is no restriction on the use of treated water for irrigation, recreation, or domestic purposes. Consult local State fish and game agency and local public water control authorities before applying this product in and around public waters. Permits may be required to treat in such areas. **NOTE:** Do not apply this product within ½ mile upstream of an active potable water intake in flowing water (i.e. river, stream, etc.) or within ½ mile of an active potable water intake in a standing body of water such as lake, pond, or reservoir. To make aquatic applications around and within ½ mile of active potable water intakes the water intake must be turned off for a minimum period of 48 hours after the application. The water intake may be turned on prior to 48 hours if the glyphosate level in the intake water is below 0.7 parts per million as determined by laboratory analysis. These aquatic applications may be made ONLY in those cases where there are alternative water sources or holding ponds which would permit the turning off of an active potable water intake for a minimum period of 48 hours after the applications. This restriction does not apply to intermittent inadvertent overspray of water in terrestrial use sites. For treatments after drawdown of water or in dry ditches, allow 7 or more days after treatment before reintroduction of water to achieve maximum weed control. Apply this product within 1 day after drawdown to ensure application to actively growing weeds. Floating mats of vegetation may require retreatment. Avoid wash-off of sprayed foliage by spray boat or recreational boat backwash or by rainfall within 6 hours of application. Do not retreat within 24 hours following the initial treatment. Applications made to moving bodies of water must be made while traveling upstream to prevent concentration of this herbicide in water. When making any bankside applications, do not overlap more than 1 foot into open water. Do not spray in bodies of water where weeds do not exist. The maximum application rate of 3.3 gallons of product per acre must not be exceeded in any single broadcast application that is being made over water. When emerged infestations require treatment of the total surface area of impounded water, treating the area in strips may avoid oxygen depletion due to decaying vegetation. Oxygen depletion may result in fish kill.

**NONCROPLAND SITES AND AQUATIC SITES WITHIN NONCROPLAND SITES:** This product may also be used to control the listed weeds, woody brush, and trees in terrestrial noncrop sites and/or in aquatic sites associated with these areas: **Terrestrial Non-Food Crop:** ornamental and/or shade trees, ornamental woody shrubs and vines, ornamental plants, agricultural rights-of-way/fencerows/hedgerows, agricultural uncultivated areas, airports/landing fields, golf course turf, industrial areas, outdoor buildings/structures, rights-of-way/fence rows/hedgerows, nonagricultural uncultivated areas/soils, ornamental and/or shade trees, cemeteries, vacant lots, paved areas (private roads/sidewalks), recreational areas, urban areas, golf courses, parks, schools, rights-of-way (highway, power lines, pipeline utility, railroads), roadsides, industrial sites, storage yards, lumber yards, petroleum tank farms, petroleum pumping station, parking areas, shelter belts, farm yards, wildlife habitat restoration and management areas. **Outdoor Residential:** ornamental and/or shade trees, ornamental herbaceous plants, ornamental woody shrubs and vines, household/domestic dwellings outdoor premises. This product may be used as a broadcast treatment or spot treatment with any of the application equipment listed on this label in noncropland sites. **For spot treatments:** Specific use recommendations for annuals, perennials, woody brush, and trees are presented in the section named WEEDS CONTROLLED. See Tables 3 and 4. **For broadcast treatments:** Specific use recommendations for noncropland sites including dosage rates, spray volumes, and restrictions are presented in Table 5.

**Table 5. Recommendations for Noncropland Use Patterns, Dosage Rates, Spray Volumes, Restrictions, and Comments for Broadcast Applications Appropriate for Each Site.**

Amount of Product Gals./Acre	Spray Volume Gals./Acre	Amount of Product fl.oz./1000²	Spray Volume Gals./1000²
Annual weeds:			
0.7 – 1.1	3 - 40	2.0 – 3.2	≤ 1.0
Perennial weeds:	3 - 40	6.0 – 10.0	≤ 1.0
2.0 – 3.3			
Woody brush & trees:	3 - 40	4.0 – 10.0	≤ 1.0
1.3 – 3.3			

**Site:** Farmsteads, farm roads, roadsides, fence rows, rights-of-way, farm buildings, golf courses, industrial sites, parks, cemeteries & other noncropland sites. **Major Weed Species:** Annual and perennial weeds including crabgrass, foxtail, johnsongrass, kochia, morningglory, multiflora rose, pigweed, trumpetcreeper, thistle, woody brush.

**Precautions, Restrictions & Comments:** SITE PREPARATION(preplant, preemergence, renovation): TRIM AND EDGE: This product may be used to trim and edge around trees, buildings, sidewalks and roads in noncropland sites. BARE GROUND: Repeated applications of this product may be used on emerged weeds to maintain bare ground. NONSELECTIVE POSTEMERGENCE: Annuals and perennials: Apply to foliage of emerged weeds. Apply at early growth stages of annual weeds and when perennial weeds are approaching maturity. Woody brush and trees: Apply this product after full leaf expansion, unless otherwise directed. Use the higher rate for larger plants and/or dense areas of growth. On vines, use the higher rate for plants that have reached the woody stage of growth. Best results are obtained when application is made in late summer or fall after fruit formation. In arid areas, best results are obtained when applications are made in the spring to early summer when brush species are at high moisture content and are flowering. Symptoms may not appear prior to frost or senescence with fall treatments. Allow 7 or more days after application before tillage, mowing or removal. Repeat treatments may be necessary to control plants regenerating from underground parts or seed. Reduced performance may result if fall treatments are made following a frost.

**WILDLIFE HABITAT RESTORATION AND MANAGEMENT AREAS:** This product is recommended for the restoration and/or maintenance of native habitat and in wildlife management areas. **Habitat restoration and maintenance:** When applied as directed, exotic and other undesirable vegetation may be controlled in habitat management areas. Applications may be made to allow recovery of native plant species, to open up water to attract waterfowl, and for similar broad-spectrum vegetation control requirements in habitat management areas. Spot treatments may be made to selectively remove unwanted plants for habitat enhancement. For spot treatments, care should be exercised to keep spray off of desirable plants. **Wildlife food plots:** This product may be used as a site preparation treatment prior to planting wildlife food plots. Apply as directed to control vegetation in the plot area. Any wildlife food species may be planted after applying this product, or native species may be allowed to reinfest the area. If tillage is needed to prepare a seedbed, wait 7 days after applying this product before tilling to allow for maximum effectiveness.

**CUT STUMP APPLICATION:** Cut stump treatments are effective on stumps with diameters larger than 3 - 4 inches in diameter. Woody vegetation may be controlled by treating freshly cut stumps of trees and resprouts with this product. Apply this product using suitable equipment to ensure coverage of the entire cambium. Cut vegetation close to the soil surface. Apply a 100% solution (undiluted) of this product to freshly cut surface immediately after cutting. Treat the entire circumference of the stump by drenching the cambium (the outermost wood between the pith and the bark). Delay in applying this product may result in reduced performance. For best results, trees should be cut during periods of active growth and full leaf expansion. The most favorable period for cut stump treatment occurs after full leaf stage in the spring and continues into early summer. Brush and trees treated in the late fall may not be fully controlled. When used according to directions for cut stump application, this product will control, partially control or suppress most woody brush and tree species, some of which are listed below: Alder *Alnus spp.*; Coyote brush *Baccharis consanfuinea*\*; Dogwood *Cornus spp.*; Eucalyptus *Eucalyptus spp.*; Hickory *Carya spp.*; Madrone *Arbutus menziesii*; Maple *Acer spp.*\*; Oak *Quercus spp.*; Poplar *Populus spp.*\*; Reed, giant *Arundo donax*; Saltcedar *Tamarix spp.*; Sweetgum *Liquidambar styraciflua*\*; Sycamore *Platanus occidentalis*\*; Tanoak *Lithocarpus densiflorus*; Willow *Salix spp.* (Not for use in California.) **INJECT AND FRILL APPLICATION:** Woody vegetation may be controlled by injection or frill application of this product. Apply this product using suitable equipment which must penetrate into living tissue. Apply the equivalent of 3.5 - 4.0 ml of this product per 2 - 3 inches of trunk diameter. This is best achieved by applying a 100% concentration (undiluted) of this product either to a continuous frill around the tree or as cuts evenly spaced around the tree below all branches. As tree diameter increases in size, better results are achieved with a continuous frill or more closely spaced cuttings. Avoid application techniques that allow runoff to occur from frill or cut areas in species that exude sap freely after frills or cutting. In species such as these, make frill or cut at an oblique angle so as to produce a cupping effect and use undiluted material. For best results, applications should be made during periods of active growth and full leaf expansion. **This treatment will CONTROL the following woody species:** Oak *Quercus spp*; Poplar *Populus spp.*; Sweet gum *Liquidambar styraciflua*; Sycamore *Platanus occidentalis*. **This treatment will SUPPRESS the following woody species:** Black gum *Nyssa sylvatica*\*; Dogwood *Cornus spp.*; Hickory *Carya spp.*; Maple, red *Acer rubrum*. (\*This product is not approved for this use on this species in the State of California.)

**STORAGE & DISPOSAL:** Do not contaminate water, food, or feed by storage or disposal. **STORAGE:** STORE ABOVE 10° F (12°C) TO KEEP PRODUCT FROM CRYSTALLIZING. Crystals will settle to the bottom. If allowed to crystallize, place in a warm room (68°F or 20°C) for several days to allow crystals to redissolve, then shake well before using. **PESTICIDE DISPOSAL:** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility. **CONTAINER HANDLING:** Nonrefillable container. Do not reuse or refill this container. Offer for recycling, if available, or puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke. **(Nonrefillable with capacities equal to or less than 5 gallons:)** Triple rinse container promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. **(Nonrefillable with capacities greater than 5 gallons:)** Triple rinse container promptly after emptying. Triple rinse as follows: Empty remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times.

**LIMITED WARRANTY AND DISCLAIMER** FOR USE ONLY AS DIRECTED. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE MANUFACTURER NEITHER MAKES NOR INTENDS ANY EXPRESS OR IMPLIED WARRANTIES, INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, WHICH ARE HEREBY EXPRESSLY DISCLAIMED. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO CASE SHALL THE MANUFACTURER BE LIABLE FOR INCIDENTAL, CONSEQUENTIAL, OR SPECIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. If these terms are not acceptable, return this product unopened immediately to the point of purchase, and the purchase price will be refunded in full. The terms of this LIMITED WARRANTY STATEMENT cannot be varied by any written or verbal statements or agreements at the point of sale or elsewhere.

EPA REG. NO. 2217-850-10807      EPA EST. NO. 1270-TX-1

1045115 05/13