

STATE OF HAWAII  
Department of Agriculture

**ACCEPTED**

LICENSE NO.

**9529.260**



# REDIGO™

## 480

Net  
Contents:

**2.5 Gallons**

**GROUP 3 FUNGICIDE**

***For: control of specified diseases on listed crops.***

**ACTIVE INGREDIENT:**

Prothioconazole, 2-[2-(1-Chlorocyclopropyl)-3-(2-chlorophenyl)-2-hydroxypropyl]-1,2-dihydro-3H-1,2,4-triazole-3-thione: . . . . .

**41.00%**

**OTHER INGREDIENTS:** . . . . . **59.00%**

**TOTAL: 100.00%**

Contains 4.00 pounds Prothioconazole per U.S. gallon

**EPA Reg. No. 264-825**

**KEEP OUT OF REACH OF CHILDREN  
CAUTION**

For **MEDICAL** And **TRANSPORTATION** Emergencies **ONLY**

Call 24 Hours A Day 1-800-334-7577

For **PRODUCT USE** Information Call 1-866-99BAYER (1-866-992-2937)

Produced for:  
Bayer CropScience LP  
P.O. Box 12014, 2 T.W. Alexander Drive  
Research Triangle Park, North Carolina 27709  
REDIGO is a trademark of Bayer.  
©2016 Bayer CropScience

160324A 04/16

US00000000A

## FIRST AID

<b>If Swallowed:</b>	<ul style="list-style-type: none"><li>• Immediately call a poison control center or doctor for treatment advice.</li><li>• Have person sip a glass of water if able to swallow.</li><li>• Do not induce vomiting unless told to do so by a poison control center or doctor.</li><li>• Do not give anything by mouth to an unconscious person.</li></ul>
<b>If Inhaled:</b>	<ul style="list-style-type: none"><li>• Move person to fresh air.</li><li>• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible.</li><li>• Call a poison control center or doctor for further treatment advice.</li></ul>
<b>If on Skin or Clothing:</b>	<ul style="list-style-type: none"><li>• Take off contaminated clothing.</li><li>• Rinse skin immediately with plenty of water for 15-20 minutes.</li><li>• Call a poison control center or doctor for treatment advice.</li></ul>
<b>If in eyes:</b>	<ul style="list-style-type: none"><li>• Hold eye open and rinse slowly and gently with water for 15-20 minutes.</li><li>• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li><li>• Call a poison control center or doctor for treatment advice.</li></ul>
<p><b>In case of emergency call toll free the Bayer CropScience Emergency Response Telephone No. 1-800-334-7577.</b></p> <p><b>Have a product container or label with you when calling a poison control center or doctor, or going for treatment.</b></p>	
Note to Physician: No specific antidote. Treat symptomatically.	

---

## PRECAUTIONARY STATEMENTS

---

### HAZARDS TO HUMANS AND DOMESTIC ANIMALS

#### CAUTION

- Harmful if swallowed or inhaled.
- Causes moderate eye irritation.
- Avoid contact with eyes and clothing. Avoid breathing vapor or spray mist.
- Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco.
- Remove and wash contaminated clothing before reuse.

#### PERSONAL PROTECTIVE EQUIPMENT (PPE)

##### Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Chemical resistant gloves made of any waterproof material
- Shoes plus socks

#### ENGINEERING CONTROLS

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

## USER SAFETY RECOMMENDATIONS

- Users should wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Users should remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Users should remove PPE immediately after handling this product.

## USER SAFETY REQUIREMENTS

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

## ENVIRONMENTAL HAZARDS

This product is toxic to estuarine/marine invertebrates, and freshwater/estuaries/marine aquatic plants. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwater or rinseate.

Prothioconazole-desthio (a degradate of prothioconazole) is known to leach through soil into ground water under certain conditions as a result of label use. Use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in ground-water contamination.

Drift and runoff are hazardous to aquatic organisms in water adjacent to treated areas. This product has a high potential for runoff for several months or more after application. Poorly draining soils and soils with shallow water tables are more prone to produce runoff that contains this product. A level, well maintained vegetative buffer strip between areas to which this product is applied and surface water features such as ponds, streams, and springs will reduce the potential for contamination of water from rainfall-runoff. Runoff of this product will be reduced by avoiding applications when rainfall is forecasted to occur within 48 hours.

## CONDITIONS OF SALE AND LIMITATIONS OF WARRANTY AND LIABILITY

Read the entire Directions for Use, Conditions, Disclaimer of Warranties and Limitations of Liability before using this product. If terms are not acceptable, return the unopened product container at once.

By using this product, user or buyer accepts the following Conditions, Disclaimer of Warranties and Limitations of Liability.

Treatment of highly mechanically damaged seed, or seed of known low vigor and poor quality, may result in reduced germination and/or reduction of seed and seedling vigor. Treat and conduct germination tests on a small portion of seed before committing the total seed lot to a selected chemical treatment. Due to seed quality conditions beyond the control of Bayer CropScience, no claims are made to guarantee germination of carry-over seed.

**CONDITIONS:** The directions for use of this product are believed to be adequate and must be followed carefully. However, it is impossible to eliminate all risks associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of Bayer CropScience. To the extent consistent with applicable law, all such risks shall be assumed by the user or buyer.

*(continued)*

**DISCLAIMER OF WARRANTIES:** TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BAYER CROPSCIENCE MAKES NO OTHER WARRANTIES, EXPRESS OR IMPLIED, OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE OR OTHERWISE, THAT EXTEND BEYOND THE STATEMENTS MADE ON THIS LABEL. No agent of Bayer CropScience is authorized to make any warranties beyond those contained herein or to modify the warranties contained herein. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BAYER CROPSCIENCE DISCLAIMS ANY LIABILITY WHATSOEVER FOR SPECIAL, INCIDENTAL OR CONSEQUENTIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT.

**LIMITATIONS OF LIABILITY:** TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER FOR ANY AND ALL LOSSES, INJURIES OR DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER IN CONTRACT, WARRANTY, TORT, NEGLIGENCE, STRICT LIABILITY OR OTHERWISE, SHALL NOT EXCEED THE PURCHASE PRICE PAID, OR AT BAYER CROPSCIENCE'S ELECTION, THE REPLACEMENT OF PRODUCT.

---

## DIRECTIONS FOR USE

---

**It is a violation of Federal law to use this product in a manner inconsistent with its labeling.**

**Read the entire label before using this product.**

For use only in commercial seed treatment equipment. Not for use in hopper box, planter box, or slurry box.

### PRODUCT INFORMATION

Redigo™ 480:

- is a broad-spectrum systemic seed treatment fungicide for the control of Ascomycetes, Basidiomycetes and Deuteromycetes diseases in a variety of crops including alfalfa, barley, buckwheat, cotton, dried shelled pea and bean (except for soybean) subgroup, millet (Pearl & Proso), oats, rice, rye, soybeans, triticale, and wheat.

### APPLICATION INSTRUCTIONS

- Redigo™ 480 provides control or suppression of many important diseases of alfalfa, barley, buckwheat, cotton, dried shelled pea and bean (except for soybean) subgroup, millet (Pearl & Proso), oats, rice, rye, soybeans, triticale, and wheat.
- Apply using standard slurry or mist-type seed treatment equipment. Uniform application of seed is necessary to ensure seed safety and best disease protection. Product should be diluted with sufficient water to ensure complete seed coverage. Consult a seed treatment specialist regarding slurry rates recommended for the crop to be treated with Redigo 480. For best results, apply to high quality, properly cleaned seed.
- When reference is made to disease suppression, suppression can mean either erratic control from good to fair or consistent control at a level below that obtained with the best commercial disease control products.
- Under conditions conducive to extended infection periods or high disease pressure, additional fungicide applications beyond the number allowed by this label may be needed. Under these conditions use another fungicide registered for the crop/disease.
- Equipment must be properly calibrated before use.

## USE RESTRICTIONS

- Not for direct application to seeds in a hopper-box, planter-box, or slurry-box.
- This product does not contain colorant. The purchaser of this product is responsible for ensuring that all seed treated with this product are adequately dyed with a suitable color to prevent its accidental use as food for man or feed for animals. Refer to 21CFR, part 2.25. Any colorant added to treated seed must be cleared for use under 40CFR, Part 153.155.
- In California – Registered only for use in commercial seed treating establishments. Not allowed for “on-farm” use.

Refer to the specific use directions and restrictions in each Crop, Crop Group or Crop Subgroup table.

## RESISTANCE MANAGEMENT RECOMMENDATIONS

- Redigo 480 is a Group 3 fungicide, which exhibits no known cross-resistance to other fungicide groups. However, fungal pathogens are known to develop resistance to products with the same mode of action when used repeatedly.
- Any fungal population may contain or develop individuals that are resistant to Redigo 480 and other Group 3 fungicides. If Group 3 fungicides are used repeatedly in the same field or in successive years as the primary method of control for targeted diseases, the resistant isolates may eventually dominate the fungal population.
- Because resistance development cannot be predicted, the use of this product should conform to resistance management strategies established for the crop and use area. Such strategies may include rotation and /or tank mixing with products having different modes of action or limiting the total number of applications per season.
- Contact your local extension specialist, certified crop advisor, and/or manufacturer for fungicide resistance management and/or integrated disease management recommendations for specific crops and pathogen populations. Bayer CropScience encourages responsible resistance management to ensure effective long-term control of the fungal diseases on this label.

---

## SPECIFIC CROP DIRECTIONS

---

### CROP USE DIRECTIONS

### APPLICATION DIRECTIONS FOR SEED TREATMENT

ALFALFA			
Disease(s) Controlled			
Seed rot and damping-off caused by <i>Rhizoctonia</i>			
Rate of Redigo 480			
seeds/lb <sup>1</sup>	fl oz/100 lbs seed	fl oz/100,000 seeds	mg ai/seed
227,000	0.48	0.002	0.0003
Crop Specific Directions and Restrictions:			
<ul style="list-style-type: none"><li>• <sup>1</sup>Seeds/lb values based on the average for rice (227,000 seeds/lb).</li><li>• Application rate specified for planting rate of 3,405,000 alfalfa seed per acre.</li></ul>			

**BEANS AND PEAS (DRIED) INCLUDING SOYBEAN**

**Crops of Crop Group 6C Including:** Soybean, Adzuki Bean, Blackeyed Pea, Broad Bean, Catjang, Chickpea, Cowpea, Crowder Pea, Field Bean, Field Pea, Guar, Kidney Bean, Lablab Bean, Lentil, Lima Bean, Moth Bean, Mung Bean, Navy Bean, Pigeon Pea, Pinto Bean, Rice Bean, Southern Pea, Tepary Bean, Urd Bean.

**Disease(s) Controlled**

Seed rot and damping-off caused by *Rhizoctonia*, *Fusarium*  
Seed decay

**Rate of Redigo 480**

Soybean	seeds/lb <sup>1</sup>	fl oz/100 lbs seed	fl oz/100,000 seeds	mg ai/seed
	1500 - 3600	0.16 - 0.32	0.0622 - 0.2987	0.0063 - 0.0303

**Rate of Redigo 480**

Crop group 6-C <sup>1</sup>	seeds/lb <sup>1</sup>	fl oz/100 lbs seed	fl oz/100,000 seeds	mg ai/seed
<b>Common Dry Beans</b>	800 - 1814	0.16 - 0.32	0.0882 - 0.4000	0.0125 - 0.0568
<b>Lentils</b>	19000	0.16 - 0.32	0.0084 - 0.0168	0.0012 - 0.0024
<b>Field Peas</b>	1600 - 5000	0.16 - 0.32	0.0320 - 0.2000	0.0045 - 0.0284
<b>Southern Peas</b>	3200 - 4000	0.16 - 0.32	0.0400 - 0.1000	0.0057 - 0.0142

**Crop Specific Directions and Restrictions:**

- Application rate specified for planting range of 125,000 to 200,000 soybean seed per acre.
- <sup>1</sup>For seed/lb information relative to other specific beans or peas from crop group 6-C, consult with your Agricultural Extension Agent or Seed Company.
- Application rate specified for planting range of 125,000 to 200,000 soybean seed per acre
- Application rate specified for planting range of 25,000 to 130,000 of Dry Beans per acre, 275,000 to 325,000 of Field Peas per acre, 24,000 to 105,000 of Southern Peas per acre.

**CEREAL GRAINS**

Barley, Buckwheat, Millet (Pearl and Proso), Oats, Rye, Triticale, Wheat, Corn and Rice (see corn and rice).

**Disease(s) Controlled**

Common bunt	Loose smut	Seed decay
Covered smut	Stinking smut	Common root rot, foot rot, and crown rot (early season suppression)
False loose smut	Stem smut	Rust, <i>Septoria</i> and powdery mildew (early season suppression)
Flag smut	True loose smut	
Leaf stripe	Seed rot, pre-emergence damping-off and seedling blight caused by soilborne <i>Rhizoctonia solani</i> , <i>Fusarium</i> , <i>Cochliobolus</i>	

**Rate of Redigo 480**

	seeds/lb <sup>1</sup>	fl oz/100 lbs seed	fl oz/100,000 seeds	mg ai/seed
<b>Barley</b>	11,700	0.16 - 0.8	0.014 - 0.068	0.0019 - 0.0097
<b>Buckwheat</b>	20,412	0.16 - 0.8	0.008 - 0.039	0.0011 - 0.0056
<b>Millet (Pearl and Proso)</b>	85,000	0.16 - 0.8	0.002 - 0.009	0.0003 - 0.0013
<b>Oats</b>	15,000	0.16 - 0.8	0.011 - 0.053	0.015 - 0.0076
<b>Rye</b>	18,000	0.16 - 0.8	0.009 - 0.044	0.013 - 0.0063
<b>Triticale</b>	16,000	0.16 - 0.8	0.010 - 0.050	0.0014 - 0.0071
<b>Wheat</b>	15,000	0.16 - 0.8	0.011 - 0.053	0.0015 - 0.0076

(continued)

## CEREAL GRAINS (continued)

### Crop Specific Directions and Restrictions:

- <sup>1</sup>Seeds/lb values based on the averages for barley (11,700 seeds/lb), buckwheat (20,412 seeds/lb), pearl millet (85,000 seeds/lb), spring oat (15,000 seeds/lb), rye (18,000 seeds/lb), triticale (16,000 seeds/lb) and winter wheat (15,000 seeds/lb).
- Corn and Rice (see corn and rice).
- For environments with Pythium pressure, combine Redigo 480 with Allegiance Flowable Seed Treatment Fungicide at rates recommended on label.
- Application rate specified for planting rate range of 351,000 to 1,300,000 barley seed per acre.
- Application rate specified for planting rate range of 734,832 to 1,469,664 buckwheat seed per acre.
- Application rate specified for planting rate range of 425,000 to 2,550,000 millet seed per acre.
- Application rate specified for planting rate range of 900,000 to 1,350,000 oat seed per acre.
- Application rate specified for planting rate range of 1,080,000 to 1,620,000 rye seed per acre.
- Application rate specified for planting rate of 1,744,000 triticale seed per acre.
- Application rate specified for planting rate of 1,740,000 wheat seed per acre.

## COTTON

### Disease(s) Controlled

Seedborne decay caused by *Fusarium* spp., *Aspergillus* spp.,  
*Cladosporium* spp., *Penicillium* spp., *Alternaria* spp.

Soilborne seedling disease caused by *Fusarium* spp., *Rhizoctonia solani*

### Rate of Redigo 480

seeds/lb <sup>1</sup>	fl oz/100 lbs seed	fl oz/100,000 seeds	mg ai/seed
4500	0.15 - 0.40	0.035 - 0.090	0.005 - 0.013

### Disease(s) Controlled

Seedling disease caused heavy infestations of *Fusarium* spp. and *Rhizoctonia solani*

### Rate of Redigo 480

seeds/lb <sup>1</sup>	fl oz/100 lbs seed	fl oz/100,000 seeds	mg ai/seed
4500	0.40 - 0.81	0.090 - 0.18	0.013 - 0.026

### Crop Specific Directions and Restrictions:

- Application rate specified for planting rate range of 29,000 - 85,000 cotton seed per acre

## CORN

Field Corn, Field Corn Grown For Seed, Popcorn, and Sweet Corn

### Disease(s) Controlled

To aid in the control or suppression of seedborn disease and early season seedling disease caused by *Fusarium* spp. and *Rhizoctonia solani*

### Rate of Redigo 480

	seeds/lb <sup>1</sup>	fl oz/100 lbs seed	fl oz/100,000 seeds	mg ai/seed
Field Corn, Field Corn Grown For Seed,	1681	0.08 - 0.8	0.038 - 0.381	0.0068 - 0.0676
Popcorn	3061	0.08 - 0.8	0.021 - 0.210	0.0037 - 0.0371
Sweet Corn	3150	0.08 - 0.8	0.020 - 0.203	0.0036 - 0.0361

### Crop Specific Directions and Restrictions:

- <sup>1</sup>Seeds/lb values based on averages for Field Corn (1681 seeds/lb), Popcorn (3061 seeds/lb), and Sweet Corn (3150 seeds/lb).
- Application rate specified for planting rate range of 8,800 to 40,250 field corn seed per acre.
- Application rate specified for planting rate range of 20,000 to 30,000 popcorn seed per acre.
- Application rate specified for planting rate range of 13,068 to 59,739 sweet corn seed per acre.

<b>RICE</b>			
<b>Disease(s) Controlled</b>			
Seed rot and damping-off caused by <i>Rhizoctonia</i> and <i>Fusarium</i> Seed decay			
<b>Rate of Redigo 480</b>			
<b>seeds/lb<sup>1</sup></b>	<b>fl oz/100 lbs seed</b>	<b>fl oz/100,000 seeds</b>	<b>mg ai/seed</b>
21850	0.16 - 0.32	0.0073 - 0.015	0.0011 - 0.0021
<b>Crop Specific Directions and Restrictions:</b>			
<ul style="list-style-type: none"> <li>• <sup>1</sup>Seeds/lb values based on the average for rice (21850 seeds/lb).</li> <li>• Do not plant treated rice seed directly into a flooded field. Do not soak treated rice seed.</li> <li>• Application rate specified for planting rate range of 1,742,400 to 2,439,360 rice seed per acre.</li> </ul>			

## SEED TAG LABELING

The Federal Seed Act requires that the container of seed treated with Redigo 480 must be labeled with the following statements:

- This seed has been treated with:
  - **Redigo 480**, which contains **41% Prothioconazole**.
- Do not use treated seed for food, feed, or oil production. User is responsible for ensuring that the seed bag meets all the requirements under the Federal Seed Act.

In addition, the US Environmental Protection Agency requires the following statements on the container of seed treated with **Redigo 480**:

- Treated areas may be replanted with any crop specified on this label as soon as practical after last application. For crops not listed on this label, do not plant back within 30 days of last application.
- **Store treated seeds away from food and feedstuffs.**
- **Do not allow children, pets, or livestock to have access to treated seed.**
- **Wear long-sleeved shirt, long pants and chemical-resistant gloves when handling treated seed.**
- Excess treated seed may be used for ethanol production only if (1) by-products are not used for livestock feed and (2) no measurable residues of pesticide remain in ethanol by-products that are used in agronomic practice.
- Treated seeds exposed on soil surface may be hazardous to wildlife. Cover or collect treated seeds spilled during loading.
- Dispose of all excess treated seed. Leftover treated seed may be buried away from water sources in accordance with local requirements.
- Do not contaminate water bodies when disposing of planting equipment wash waters.
- Treated at manufacturer's recommended rate
- Treated seed must be adequately covered with soil at planting. Plant at a minimum depth of 1/2 inch.
- Dispose of seed packaging or containers in accordance with local requirements. Do not use empty seed bags for any other purpose.
- To reduce seed dust which can drift onto blooming crops or weeds, ensure that planting equipment is functioning properly in accordance with manufacturer's recommendations. Surplus seed or empty seed containers should be stored or disposed according to local federal regulations.

*(continued)*

- After the seeds have been planted, do not enter or allow worker entry into the treated areas during the restricted-entry interval (REI) of 12 hours. Exception: Once the seeds are planted in soil or other planting media, the Worker Protection Standard allows workers to enter the treated area without restriction if there will be no worker contact with the treated seeds in the soil or with the planting media.

**For PRODUCT USE Information Call 1-866-99BAYER (1-866-992-2937)  
For MEDICAL, TRANSPORTATION or OTHER Emergencies ONLY Call 24  
Hours A Day 1-800-334-7577**

## STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage, disposal or cleaning of equipment.

### **Pesticide storage**

Store in a cool, dry place and in such a manner as to prevent cross contamination with other pesticides, fertilizers, food, and feed. Store in original container and out of the reach of children, preferably in a locked storage area.

Handle and open container in a manner as to prevent spillage. If container is leaking, invert to prevent leakage. If the container is leaking or material is spilled for any reason or cause, carefully dam up spilled material to prevent runoff. Refer to Precautionary Statements on label for hazards associated with the handling of this material. Do not walk through spilled material. Absorb spilled material with absorbing type compounds and dispose of as directed for pesticides below. In spill or leak incidents, keep unauthorized people away. You may contact the Bayer CropScience Emergency Response Team for decontamination procedures or any other assistance that may be necessary. The Bayer CropScience Emergency Response Telephone No. is 1-800-334-7577.

### **Pesticide disposal**

Wastes resulting from the use of this product must be disposed of on-site or at an approved waste disposal facility.

### **Container handling**

#### **Dilutable Seed Treatment Products in Non-Refillable Plastic Containers Rigid, Non-refillable containers (equal to or less than 5 gallons)**

Non-refillable container. Do not reuse or refill this container. Offer for recycling, if available. Triple rinse or pressure rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times.

Pressure rinse as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 PSI for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

Once container is rinsed, offer for recycling if available or puncture and dispose of in a sanitary landfill.

#### **Rigid Non-refillable containers that are too large to shake (i.e., with capacities greater than 5 gallons or 50 lbs)**

Non-refillable container. Do not reuse or refill this container. After emptying product from container, either return container to Bayer CropScience per instructions from Bayer CropScience Customer Service Center (1-800-527-4781), or rinse and either recycle or dispose of the container as follows:

**Bottom Discharge IBC (e.g. – Schuetz Caged IBC or Snyder Square Stackable)**

Pressure rinsing the container before final disposal is the responsibility of the person disposing of the container. To pressure rinse the container before final disposal, empty the remaining contents from the IBC into application equipment or mix tank. Raise the bottom of the IBC by 1.5 inches on the side which is opposite of the bottom discharge valve to promote more complete product removal. Completely remove the top lid of the IBC. Use water pressurized to at least 40 PSI to rinse all interior portions. Continuously pump or drain rinsate into application equipment or rinsate collection system while pressure rinsing. Continue pressure rinsing for 2 minutes or until rinsate becomes clear. Replace the lid and close bottom valve.

**Top Discharge IBC, Drums, Kegs (e.g.– Snyder 120 Next Gen, Bonar B120, Drums, and Kegs)**

Triple rinsing the container before final disposal is the responsibility of the person disposing of the container. To triple rinse the container before final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill the container at least 10 percent full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Rinse all interior surfaces. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this procedure two more times.

Once container is rinsed, offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration.





# Redigo™ 480

GROUP 3 FUNGICIDE

For: control of specified diseases on listed crops.

## ACTIVE INGREDIENT:

Prothioconazole, 2-[2-(1-Chlorocyclopropyl)-3-(2-chlorophenyl)-2-hydroxypropyl]-1,2-dihydro-3H-1,2,4-triazole-3-thione: . . . . . 41.00%

OTHER INGREDIENTS: . . . . . 59.00%

TOTAL: 100.00%

Contains 4.00 pounds Prothioconazole per U.S. gallon

EPA Reg. No. 264-825

## KEEP OUT OF REACH OF CHILDREN CAUTION

For **MEDICAL** And **TRANSPORTATION** Emergencies **ONLY**

Call 24 Hours A Day 1-800-334-7577

For **PRODUCT USE** Information Call 1-866-99BAYER (1-866-992-2937)

### PRECAUTIONARY STATEMENTS

#### HAZARDS TO HUMANS AND DOMESTIC ANIMALS CAUTION

- Harmful if swallowed or inhaled.
- Causes moderate eye irritation.
- Avoid contact with eyes and clothing. Avoid breathing vapor or spray mist.
- Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco.
- Remove and wash contaminated clothing before reuse.

### STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage, disposal or cleaning of equipment.

#### Pesticide storage

Store in a cool, dry place and in such a manner as to prevent cross contamination with other pesticides, fertilizers, food, and feed. Store in original container and out of the reach of children, preferably in a locked storage area.

Handle and open container in a manner as to prevent spillage. If container is leaking, invert to prevent leakage. If the container is leaking or material is spilled for any reason or cause, carefully dam up spilled material to prevent runoff. Refer to Precautionary Statements on label for hazards associated with the handling of this material. Do not walk through spilled material. Absorb spilled material with absorbing type compounds and dispose of as directed for pesticides below. In spill or leak incidents, keep unauthorized people away. You may contact the Bayer CropScience Emergency Response Team for decontamination procedures or any other assistance that may be necessary. The Bayer CropScience Emergency Response Telephone No. is 1-800-334-7577.

#### Pesticide disposal

Wastes resulting from the use of this product must be disposed of on-site or at an approved waste disposal facility.

#### Container handling

##### Dilutable Seed Treatment Products in Non-Refillable Plastic Containers

#### Rigid, Non-refillable containers (equal to or less than 5 gallons)

Non-refillable container. Do not reuse or refill this container. Offer for recycling, if available. Triple rinse or pressure rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times.

Pressure rinse as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 PSI for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

Once container is rinsed, offer for recycling if available or puncture and dispose of in a sanitary landfill.

Bayer CropScience LP

P.O. Box 12014, 2 T.W. Alexander Drive  
Research Triangle Park, North Carolina 27709  
©2016 Bayer CropScience

REDIGO is a trademark of Bayer.

US00000000A 160324A 04/16

NET CONTENTS: 2 1/2 GALLONS



(01) 0 0785740 0000 \*