



ClickSan™ Disinfectant/Sanitizer



ACTIVE INGREDIENTS:	
Octyl decyl dimethyl ammonium chloride.....	6.9120%
Dioctyl dimethyl ammonium chloride.....	2.7648%
Didecyl dimethyl ammonium chloride.....	4.1472%
Alkyl (C ₁₂ , 50%; C ₁₂ , 40%; C ₁₆ , 10%) dimethyl benzyl ammonium chloride.....	9.2160%
Other Ingredients.....	76.9600%
Total.....	100.0000%



KEEP OUT OF REACH OF CHILDREN

DANGER

FOR INDUSTRIAL USE

9553.45

LICENSED

PERIOD 2015-2017 LIC. NO.

Medical Emergency: (877) 231-2615 or call collect 0 (952) 853-1713

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

DANGER. Corrosive. Causes irreversible eye damage and skin burns. Do not get in eyes, on skin, or on clothing. Harmful if swallowed, inhaled or absorbed through the skin. Wear protective eyewear (goggles, face shield or safety glasses), protective clothing and protective gloves (rubber, chemical resistant) when handling. Avoid breathing spray mist. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco. Remove and wash contaminated clothing before reuse.



An **ECOLAB** Company
Distributed by:
KAY CHEMICAL COMPANY
8300 Capital Drive
Greensboro, NC 27409-9790, USA
Manufactured in USA
Customer Service : (800) 529-5458



LICENSED

PERIOD 2012-2014 LIC. NO.

EPA Reg. No. 6836-305-5389
EPA Est. No. 58046-TX-2(X), 5389-NC-1(G)
Superscript refers to the first letter of the date code.



LICENSED

PERIOD 2009-2011 LIC. NO.

PULL TAB AT LABEL CORNER TO SEE
FIRST AID AND DIRECTIONS FOR USE

Net Contents: 1 US Gallon (3.78 L)

FIRST AID

IF IN EYES: Hold eye open and rinse slowly and gently with water for 15 - 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.

IF ON SKIN OR CLOTHING: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15 - 20 minutes.

IF SWALLOWED: Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor. Do not give anything by mouth to an unconscious person.

IF INHALED: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible.

Call a poison control center or doctor for treatment advice. Have the product container or label with you when calling a Poison Control Center or doctor or going for treatment.

NOTE TO PHYSICIAN: Probable mucosal damage may contraindicate the use of gastric lavage.

ClickSan™ Disinfectant/Sanitizer kills:

Bacteria: *Staphylococcus aureus* (*Staph*), *Salmonella enterica* (*Salmonella*), *Brevibacterium ammoniagenes* (*Brevibacterium*), *Campylobacter jejuni* (*Campylobacter*), *Escherichia coli* [*E. coli*], *Escherichia coli* pathogenic strain 0157:H7 (*pathogenic E. coli*), *Klebsiella pneumoniae* [*Klebsiella*], *Listeria monocytogenes* [*Listeria*], (*Community Associated*) *Methicillin-Resistant Staphylococcus aureus* - [*CA-MRSA*] [*MRSA*] (*NR5123*) (*USA400*), *Pseudomonas cepacia* [*Pseudomonas*], *Salmonella typhi* [*Salmonella*], *Yersinia enterocolitica*
Viruses: *Adenovirus Type 5, *Herpes Simplex Virus Type 1 (Herpes), *Herpes Simplex Virus Type 2 (Herpes), *Influenza A2 / Hong Kong [Influenza] [Flu] Virus, *Norwalk Virus - Feline Calicivirus, *SARS Associated Human Coronavirus, *Vaccinia Virus (Pox Virus)

Fungi: *Aspergillus niger*

ClickSan™ Disinfectant/Sanitizer is an effective sanitizer at 150 ppm active quat for use on food contact surfaces in 400 ppm hard water against: *Escherichia coli*, *Staphylococcus aureus*

ClickSan™ Disinfectant/Sanitizer is an effective sanitizer at 200 ppm active quat for use on food contact surfaces in 500 ppm hard water against: *Campylobacter jejuni*, (*Community Associated*) *Methicillin-Resistant Staphylococcus aureus* - [*CA-MRSA*] [*MRSA*] (*NR5384*) (*USA300*), (*Community Associated*) *Methicillin-Resistant Staphylococcus aureus* - [*CA-MRSA*] [*MRSA*] (*NR5123*) (*USA400*), *Escherichia coli*, *Escherichia coli* 0157:H7, *Klebsiella pneumoniae*, *Listeria monocytogenes*, *Salmonella enterica*, *Shigella sonnei*, *Staphylococcus aureus*, *Vibrio cholerae*, *Yersinia enterocolitica*

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Dilution:

Disinfection (1:512)..... 1 oz. per 4 gallons of water.....(450 ppm active quat)
Sanitizer (1:544) 4 oz. per 18 gallons of water.....(400 ppm active quat)
Sanitizer (1:1088) 2 oz. per 18 gallons of water.....(200 ppm active quat)
Sanitizer (1:1472) 1 1/2 oz. per 18 gallons of water(150 ppm active quat)

DISINFECTION / VIRUCIDAL* / FUNGICIDAL DIRECTIONS:

Add 1 oz. of ClickSan™ Disinfectant/Sanitizer per 4 gallons of water (or equivalent dilution) to disinfect hard, nonporous surfaces. Before use in federally inspected meat and poultry food processing plants and dairies, food products and packaging materials must be removed from the room or carefully protected. Apply use-solution with a cloth, mop, sponge, sprayer or by immersion, through wetting surfaces.

For sprayer applications, use a coarse spray device. Spray 6 - 8 inches from surface; rub with brush, sponge or cloth. Do not breathe spray.

NOTE: For spray applications, cover or remove all food products.

Treated surfaces must remain wet for 10 minutes. Wipe dry with a clean cloth, sponge or mop or allow to air dry.

Rinse food contact surfaces such as counter tops, tables, picnic tables, exteriors of appliances and / or stove tops with potable water prior to reuse. Do not use on glasses, dishes, or utensils as a disinfectant.

For heavily soiled areas, preclean first. Prepare a fresh solution daily or more often if the use solution becomes visibly diluted or soiled.

TO SANITIZE FOOD PROCESSING EQUIPMENT, UTENSILS, AND OTHER FOOD CONTACT ARTICLES REGULATED BY 40 CFR § 180.940(a) (IN A THREE COMPARTMENT SINK):

1. Scrape, flush, or presoak articles (whether mobile or stationary) to remove gross food particles and soil.
2. Thoroughly wash articles with an appropriate detergent or cleaner.
3. Rinse articles thoroughly with potable water.
4. Sanitize by immersing articles with a use-solution of 1 1/2 - 4 oz. of ClickSan™ Disinfectant/Sanitizer per 18 gallons of water (150-400 ppm active quat) (or equivalent dilution) for at least 60 seconds. Articles too large for immersing must be thoroughly wetted by rinsing, spraying, or swabbing.
5. Remove immersed items from solution to drain and then air dry. Non-immersed items must also be allowed to air dry. Do not rinse.

U.S. PUBLIC HEALTH SERVICE

CLEANING AND SANITIZING: Equipment and utensils must be thoroughly pre-flushed or pre-scraped and when necessary, pre-soaked to remove gross food particles and soil.

1. Thoroughly wash equipment and utensils in hot detergent solution.
2. Rinse utensils and equipment thoroughly with potable water.
3. Sanitize equipment and utensils by immersion in a use solution of 1 1/2 - 4 oz. of ClickSan™ Disinfectant/Sanitizer per 18 gallons of water (150-400 ppm active quat) (or equivalent dilution) for at least 60 seconds at a temperature of 75°F.
4. For equipment and utensils too large to sanitize by immersion, apply a use-solution of 1 1/2 - 4 oz. of ClickSan™ Disinfectant/Sanitizer per 18 gallons of water (150-400 ppm active quat) (or equivalent dilution) by rinsing, spraying or swabbing until thoroughly wet.
5. Allow sanitized surface to drain and air dry. Do not rinse.

WISCONSIN STATE BOARD OF HEALTH DIRECTIONS FOR EATING ESTABLISHMENTS

1. Scrape and pre-wash utensils and glasses whenever possible.
2. Wash with a good detergent or compatible cleaner.
3. Rinse with potable water.
4. Sanitize in a solution of 2-4 oz. of ClickSan™ Disinfectant/Sanitizer per 18 gallons of water (200-400 ppm active quat). Immerse all utensils for at least two minutes or for contact time specified by governing sanitary codes.
5. Place sanitized utensils on a rack or drain board to air dry.
6. Prepare a fresh sanitizing solution at least daily or when visibly soiled or diluted.

NOTE: A clean potable water rinse following sanitization is not permitted under HFS 196, Appendix 7-204.1I of the Wisconsin Administrative Code (reference 40 CFR 180.940 (a)).

STORAGE AND DISPOSAL

Store in original container in areas inaccessible to children.

Nonrefillable container. Do not reuse or refill this container. Clean container promptly after emptying. Triple rinse as follows: Fill container 1/4 full with water and reclose the container. Agitate vigorously and dispose of rinsate. Repeat two more times. Wrap container and put in trash or offer for recycling if available.