

25% MORE FREE!

GORDON'S®

**Contains
Glyphosate**

PRONTO®
FAST ACTING

**Weed & Grass
KILLER CONCENTRATE**

**Makes Up To
30 Gallons**

9554.110

- **Lawn Renovation**
 - **Garden Preparation**
 - **Fenceline Cleanup**
- and more!*



Rain Fast In Hours

EVEN KILLS THE ROOTS!

ACTIVE INGREDIENT:
Isopropylamine salt of
glyphosate* 25.0%

OTHER INGREDIENTS: 75.0%

TOTAL 100.0%

KEEP OUT OF REACH OF CHILDREN

CAUTION

See back panel for additional
Precautionary Statements
and First Aid.

*N-(phosphonomethyl)glycine

954/2-2005

NET CONTENTS: 40 FL. OZ.



LICENSED

PERIOD **2012-2014** LIC. NO.



Department of Agriculture
STATE OF HAWAII

LICENSED

PERIOD **2015-2017** LIC. NO.



Department of Agriculture
STATE OF HAWAII

LICENSED

PERIOD **2009-2011** LIC. NO.

GORDON'S®

PRONTO®
FAST ACTING

Weed & Grass
KILLER CONCENTRATE

Contains Glyphosate
Makes Up To 30 Gallons
Even Kills The Roots!

***Peel open for additional
Precautionary Statements
and First Aid.***

954/6-2006 AP090805
EPA REG. NO. 2217-846
EPA EST. NO. 2217-KS-2

Manufactured By

G **pbi / GORDON**
CORPORATION

An Employee-Owned Company
1217 West 12th Street
Kansas City, Missouri 64101

consumer.pbigordon.com



40 FL. OZ.

GORDON'S®

PRONTO®
FAST ACTING

Weed & Grass
KILLER CONCENTRATE

- **Lawn Renovation**
- **Garden Preparation**
- **Fenceline Cleanup and more!**

ACTIVE INGREDIENT:

Isopropylamine salt of glyphosate* 25.0%

OTHER INGREDIENTS: 75.0%

TOTAL 100.0%

*N-(phosphonomethyl)glycine

KEEP OUT OF REACH OF CHILDREN

CAUTION

See next page for additional
Precautionary Statements and First Aid.



**READ THE ENTIRE LABEL FIRST.
OBSERVE ALL PRECAUTIONS AND
FOLLOW DIRECTIONS CAREFULLY.**

PRECAUTIONARY STATEMENTS

Hazards to Humans & Domestic Animals

CAUTION: Causes moderate eye irritation. Harmful if absorbed through skin. Harmful if swallowed. Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water.

First Aid

If swallowed:	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by a poison control center or doctor. • Do not give anything by mouth to an unconscious person.
If on skin or clothing:	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.
If inhaled:	<ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give

First Aid (Continued)

	<p>artificial respiration, preferably mouth-to-mouth if possible.</p> <ul style="list-style-type: none"> • Call a poison control center or doctor for treatment advice.
If in eyes:	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
<p>Have the product container or label with you when calling a poison control center or doctor or going for treatment.</p>	

ENVIRONMENTAL HAZARDS:

Do not apply directly to water. Do not contaminate water when disposing of equipment washwater and rinsate.

PHYSICAL OR CHEMICAL HAZARDS:

Combustible. Do not use or store near heat or open flame.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Keep children and pets out of treated areas until sprays have dried.

STORAGE & DISPOSAL

PESTICIDE STORAGE: Store in original container in a locked storage area inaccessible to children and domestic animals. Protect container from freezing.

PESTICIDE DISPOSAL: Securely wrap original container in several layers of newspaper and discard in trash.

CONTAINER DISPOSAL: Do not reuse container. Rinse thoroughly before discarding in trash.

HOW IT WORKS:

Facts and Tips

- Pronto® Fast Acting Weed & Grass Killer Concentrate is a multipurpose herbicide designed to control a broad spectrum of emerged annual and perennial weed species. Pronto® Weed & Grass Killer Concentrate will control unwanted species of grass, sedge, broadleaf weeds, and certain woody species around trees, shrubs, fences.
- This product kills plants by inhibiting the production of a substance found only in plants. It enters plants through the sprayed foliage and moves to the roots.
- This product is nonselective and will kill or injure all vegetation contacted by the spray. Do not spray plants or grasses you like — they may die too. Not recommended for spot weed control in lawns since this product kills lawn grasses.
- No soil residual activity — any product not absorbed by plants breaks down into natural materials without moving in or on the soil to untreated plants.

- Plants that have not emerged at the time of application will not be controlled.
- Time to symptoms — Best results will be obtained when the weed species are actively growing. Visible effects are gradual — wilting and yellowing, advancing to complete browning and root destruction. Weeds usually yellow and wilt within 2 to 4 days and plant death occurs in 1 to 2 weeks.

Product Description and Spray Coverage: This product is a nonselective herbicide that will kill all vegetation contacted. This herbicide kills unwanted plants by inhibiting the production of a substance found only in plants. It enters plants through the sprayed foliage and moves to the roots. Avoid direct spray or drift onto desirable plants, lawn grasses, shrubs and trees. Protect desirable plants and vegetation with a piece of plastic or cardboard. If desirable plants are accidentally sprayed, wash plants with water immediately. Any glyphosate not absorbed by the plant is broken down into natural materials. This product has no soil residual activity and thus will not affect nearby untreated plants.

WHERE TO USE:

For General Weed Control and Spot Spraying

- Use to prepare sites for large plantings, for planting flower beds, for planting fruit trees, ornamentals, vegetables, garden sites and to renovate lawns.
- Along and on sidewalks, gravel/brick walkways, paths, curbs, paved areas (private roads and streets), driveways, parking areas, recreational areas.

- Along ornamental lawns and turfgrass. Edging along/around lawns/yards and landscape plantings.
- In areas for lawn replacement or lawn renovation/replacement.
- Around or near ornamental/shade trees, fruit and nut trees, grapevines, shrubs, groundcovers, flower beds, mulched landscape beds.
- For garden plot preparation, vegetable plot preparation, and small fruits and berries plot preparation.
- For flower garden preparation.
- Along fences, hedgerows, under fencelines.
- Around rocks, signposts, light posts, or on fringe areas.
- Around or along buildings/structures, foundations, patios, houses, mobile homes, vacant lots, urban areas, domestic outdoor premises, and other areas associated with household or home life.
- Urban areas where total vegetation control is desired.
- Farmsteads where total vegetation control is desired: around farmstead building foundations, along fences, hedgerows, fencerows, under fence-lines, around shelter belts and other areas where nonselective weed control is desired.

TOUGH WEED CONTROL **(ANNUALS, PERENNIALS, BRUSH)**

**CONTROLS WEEDS AND GRASSES INCLUDING
LAWN AND TURFGRASS SPECIES: Pronto® Weed &**

Grass Killer Concentrate will provide full or partial control (suppression) of these lawn and turfgrass species:

Bahiagrass
 Bentgrass, colonial, creeping and redtop
 Bermudagrass, common and hybrid
 Bluegrass, annual, rough, and Kentucky
 Bromegrass, downy and smooth
 Buffalograss
 Canarygrass, reed
 Carpetgrass
 Centipedegrass
 Fescues, chewings, hard, red, tall, sheep
 Foxtail, green, yellow, creeping, meadow, giant
 Johnsongrass
 Kikuyugrass
 Orchardgrass
 Paspalum *spp.*
 Ryegrass, annual, Italian, perennial, and hardy
 St. Augustinegrass
 Zoysiagrass or Japanese lawngrass

WOODY SPECIES:

Pronto® Weed & Grass Killer Concentrate will provide full or partial control (suppression) of these woody species:

Alder	Dewberry
Blackberry	Elm
Barberry	Elderberry
Caenothus <i>spp.</i>	Honeysuckle
Cherry	Hickory
Coyote brush	Kudzu

Lantana	Oak
Maple	Quaking aspen
Multiflora rose	Raspberry
Mock orange*	Sumac
Manzanita	Trumpet creeper
Pine	Wild blackberry
Poison ivy	Western red cedar*
Poison oak	Willow
Poison hemlock	*Not for use in California

ANNUAL AND PERENNIAL WEED SPECIES:

Pronto® Weed & Grass Killer Concentrate will provide full or partial control (suppression) of these annual and perennial weed species:

Artichoke, Jerusalem	Creeping beggarweed
Barnyardgrass	Creeping Charlie
Beggarweed	Curly dock
Bittercress	Dallisgrass
Black medic	Dandelion
Broadleaf plantain	Evening primrose
Buckhorn plantain	Fall panicum
Bull thistle	False dandelion
Burclover	Fennel
Canada thistle	Fiddleneck
Chickweed	Field bindweed
Clover	Field pennycress
Cocklebur	Field sandbur
Common groundsel	Filaree
Common mullien	Florida pusley
Common ragweed	Garden spurge
Coral-bead	Goosegrass
Crabgrass	Green foxtail

Guineagrass	Quackgrass
Hemp sesbania	Ragweed
Henbit	Red clover
Horsenettle	Red sorrel*
Horseradish	Redroot pigweed
Horseweed/marestail	Shepherdspurse
Iceplant	Smartweed
Ironweed	Smooth catsear
Knapweed	Smooth pigweed
Knawel	Sowthistle
Knotweed	Spotted spurge
Lambsquarters	Spurge
London rocket	Sunflower
Lovegrass	Tansy ragwort
Mallow	Tansy mustard
Mayweed	Texas panicum
Milkweed	Thistle
Mouseear chickweed	Timothy
Nimblewill	Toadflax
Oleander*	Torpedograss
Oxalis	Trailing lantana
Pampas grass	Tumble mustard
Pennsylvania smartweed	Vaseygrass
Pigweed	Velvetleaf
Plantain	Virginia-creeper
Prickly lettuce	White clover
Primrose	Whitetop
Prostrate spurge	Wild barley
Puncturevine	Wild carrot
Purple cudweed	Wild geranium
Purple nutsedge	Wild morningglory
Purslane	Wild mustard

Wild oats
Wild sweet potato
Witchgrass
Yarrow
Yellow nutsedge

Yellow oxalis
Yellow starthistle
and many other weeds

*Not for use in California

WHEN TO USE [WHEN TO APPLY]:

Apply to actively growing weeds when the vegetation is green. Use when the wind is calm to avoid drift. Make applications on warm sunny days when the day-time temperature is above 60°F. Weed control is accelerated by warm sunny weather. Weeds usually begin to wilt within hours with complete kill in 1 to 2 weeks. Rain or watering 2 hours after application will not wash away effectiveness. Hard-to-control weeds and thick lawn grasses (like bermudagrass) may require a second application.

HOW TO USE [HOW TO APPLY]:

This product is formulated for spot treatment of individual weeds and grasses or for treatment of large areas as instructed on this label.

To apply this product use any of the following equipment:

- Plastic, aluminum or stainless steel tank sprayer
- Ortho Dial 'N Spray®
- Ortho Spray-Ette®
- A hose end sprayer recommended for herbicides.
- Hand-trigger pump sprayer.
- **Do not use or store in galvanized or unlined steel sprayer, or through any irrigation system or sprinkling can.**

- Apply to evenly and thoroughly wet weeds and grasses.
- Use a tank type sprayer with a coarse spray nozzle for spot treatments.
- Use the Ortho Dial 'N Spray[®] or other hose end sprayer to treat large areas such as for lawn renovation or to prepare sites for planting flower beds or vegetable gardens.
- Treated weeds begin to wilt in about 2 to 4 days with complete kill of the weed including the root in 2 to 4 weeks.
- Pronto[®] Weed & Grass Killer Concentrate does not move or build up in the soil to effect untreated plants growing nearby.
- Ornamental plantings can be made 3 days after treatment. Vegetable gardens can be seeded or planted 3 days after treatment.
- For lawn replacement, lawns may be replanted 1 week after spraying.
- Measure the correct amount of Pronto[®] Weed & Grass Killer Concentrate and water to add to sprayer from the directions on next page.

TABLE 1. SPRAY PREPARATION WITH HOSE END SPRAYERS AND TANK SPRAYERS

SPRAYER TYPE	MIXING INSTRUCTIONS		TREAT
	Annuals or General Weed Control	Tough Weed Control Perennial Weeds	
Standard Tank Sprayer	<ul style="list-style-type: none">• Use 2.25 fl. oz. of Pronto® Weed & Grass Killer Concentrate in 1 gallon of water.	<ul style="list-style-type: none">• Use 4.5 fl. oz. of Pronto® Weed & Grass Killer Concentrate in 1 gallon of water.	300 sq. ft. (30 ft x 10 ft)
Ortho Dial 'N Spray® Hose End Sprayer	<ul style="list-style-type: none">• Add 2.25 fl. oz. of Pronto® Weed & Grass Killer Concentrate to sprayer jar.• DO NOT ADD WATER.• Set dial to 3 oz. and apply.	<ul style="list-style-type: none">• Add 4.5 fl. oz. of Pronto® Weed & Grass Killer Concentrate to sprayer jar.• DO NOT ADD WATER.• Set dial to 5 oz. and apply.	300 sq. ft.

(Continued on next page)

TABLE 1. SPRAY PREPARATION WITH HOSE END SPRAYERS AND TANK SPRAYERS (Cont.)

SPRAYER TYPE	MIXING INSTRUCTIONS		TREAT
	Annuals or General Weed Control	Tough Weed Control Perennial Weeds	
Ortho Spray-Ette® Hose end Sprayer	<ul style="list-style-type: none">• Add 8 fl. oz. of Pronto® Weed & Grass Killer Concentrate to sprayer jar.• Fill to 4 gallon mark.	<ul style="list-style-type: none">• Add 16 fl. oz. of Pronto® Weed & Grass Killer Concentrate to sprayer jar.• Fill to 4 gallon mark.	1200 sq. ft. (30 ft x 40 ft)
Hand Trigger Sprayers	<ul style="list-style-type: none">• Use 1 tbs. (0.5 fl. oz.) of Pronto® Weed & Grass Killer Concentrate in 24 fl.oz. of water.	<ul style="list-style-type: none">• Use 2 tbs. (1.0 fl. oz.) of Pronto® Weed & Grass Killer Concentrate in 24 fl.oz. of water.	Use for spot treatments or single weeds

EQUIVALENT MEASUREMENTS

- 1 teaspoon (tsp.)
 1 tablespoon (tbs.) = 3 teaspoons (tsp.)
 1 fluid ounce (fl.oz.) = 2 tablespoons (tbs.)
 2 fluid ounces (fl.oz.) = 4 tablespoons (tbs.)

Pronto® Weed & Grass Killer Concentrate in these container sizes will treat these approximate areas:

CONTAINER SIZE	ANNUAL WEEDS (Treated Area, Sq. Ft.)	PERENNIAL WEEDS (Treated Area, Sq. Ft.)
1 pint	2,400 sq. ft.	1,200 sq. ft.
1 quart	4,800 sq. ft.	2,400 sq. ft.
2 quarts (1/2 gallon)	9,600 sq. ft.	4,800 sq. ft.
1 gallon	19,200 sq. ft.	9,600 sq. ft.

CLEANING APPLICATION EQUIPMENT:

To clean sprayer after use, rinse spray equipment thoroughly and flush all sprayer components with water 3 times after use. Spray rinsate on bare soil or gravel surface. Do not apply rinsate to treated weeds. Only after a very thorough cleaning can the sprayer or spray equipment be used for application of other products.

WHERE TO USE

For General Weed Control: Use near fences, in paths, patios, sidewalks, curbs and driveways, around ornamental trees, shrubs, flower beds and plantings, around buildings. Control weeds in brick and gravel walks. Use for trimming and edging landscapes. NOT recommended for spot weed treatment in lawns, since glyphosate kills all green plants, including lawn grasses.

Spot Treatments: Spray directly on actively growing weeds in and around flower beds, ornamental trees, fruit and nut trees, grapevines, shrubs, fences, driveways and walkways. Do not spray fruit and nut trees or grapevines directly.

For Landscaping:

- Prepare areas for planting of ornamentals, trees, shrubs, desert landscapes, rock gardens, flower beds or similar plantings.
- Delay mowing, tilling, or disturbing the treated vegetation for at least seven (7) days after application of this product for the best results. However, if the treated vegetation is not disturbed, planting or transplanting of trees, flowers, shrubs, bedding plants may be done at one (1) day after application.

For Lawn Renovation and Lawn Replacement:

- Spot treatments or broadcast applications can be used to establish or renovate warm season and cool season turfgrass from seed or planting stock. This product may be applied before planting, seeding, sprigging, plugging, or sodding.
- Use Pronto® Weed & Grass Killer Concentrate to remove the existing lawn and kill weeds and grasses.

Uniform and complete spray coverage of all existing vegetation is required.

- Bermudagrass, nutsedge, torpedograss and deep rooted perennials may require a second treatment in 5 to 7 days.
- For best results, apply in spring or fall when daytime temperatures are at least 60°F.
- Best results are obtained when one mowing is skipped before and after the application of this product.
- If the lawn is dry, water lawn before applying Pronto® Weed & Grass Killer Concentrate. Water again 2 to 3 days after application.
- Hose end sprayers and tank sprayers are appropriate for localized spot treatments and broadcast (entire) applications on small turfgrass areas, e.g. ~10,000 sq. ft. Refer to Table 1 for the spray preparation using hose end sprayers and tank sprayers.
- Conventional sprayers fitted with spray booms and spray guns are appropriate for broadcast (entire) applications on large turfgrass areas, e.g. ~20,000 sq. ft. Refer to Table 2 for the spray preparation using power spray equipment suitable for large turfgrass areas.
- Seeding, sprigging, plugging or sodding may be done at seven (7) days after application of this product. Prepare the soil for planting by raking, roto-tilling, vertical cutting, or other tillage method.
- Rake up and remove loosened thatch or debris.
- Mix in starter fertilizer and soil amendments if needed. Establish the lawn according to local cultural practices.

**TABLE 2. SPRAY PREPARATION FOR LAWN RENOVATION AND LAWN REPLACEMENT
USING POWER SPRAY EQUIPMENT WITH BROADCAST APPLICATIONS**

Site	Major Weed Species	Amount of Product, Quarts/Acre	Spray Volume, Gals./Acre	Amount of Product, fl. oz./1000 sq. ft.	Spray Volume, Gals./1000 sq. ft.
Lawn & turf seedbeds (prior to establishment):	Many annual and perennial grasses and broadleaf weeds, including annual bluegrass, crabgrass, lambsquarters and ragweed.	Annual weeds: 1.8 - 3.6	≤ 25	1.3 - 2.6	< 1
		Perennial weeds: 5.5 - 9.5	≤ 25	4.0 - 7.0	< 1
	Bahiagrass	5.5 - 9.5	≤ 25	4.0 - 7.0	< 1
	Bentgrass, colonial creeping	2.7 - 3.6	≤ 25	2.0 - 2.6	< 1

(Continued on next page)

TABLE 2. SPRAY PREPARATION FOR LAWN RENOVATION AND LAWN REPLACEMENT USING POWER SPRAY EQUIPMENT WITH BROADCAST APPLICATIONS (Cont.)

Site	Major Weed Species	Amount of Product, Quarts/Acre	Spray Volume, Gals./Acre	Amount of Product, fl. oz./ 1000 sq. ft.	Spray Volume, Gals./ 1000 sq. ft.
	Bermudagrass, common and hybrid	5.5 - 9.5	≤ 25	4.0 - 7.0	< 1
	Bluegrass, annual rough and Kentucky	1.8 - 3.6	≤ 25	1.3 - 2.6	< 1
	Bromegrass, downy and smooth	1.8 - 3.6	≤ 25	1.3 - 2.6	< 1
	Orchardgrass	1.8 - 3.6	≤ 25	1.3 - 2.6	< 1
	Canarygrass, reed	3.6 - 5.5	≤ 25	2.6 - 4.0	< 1

(Continued on next page)

**TABLE 2. SPRAY PREPARATION FOR LAWN RENOVATION AND LAWN REPLACEMENT
USING POWER SPRAY EQUIPMENT WITH BROADCAST APPLICATIONS (Cont.)**

Site	Major Weed Species	Amount of Product, Quarts/Acre	Spray Volume, Gals./Acre	Amount of Product, fl. oz./ 1000 sq. ft.	Spray Volume, Gals./ 1000 sq. ft.
	Fescue, tall	3.6 - 5.5	≤ 25	2.6 - 4.0	< 1
	Fescues, chewings, hard, red, sheep	5.5 - 9.5	≤ 25	4.0 - 7.0	< 1
	Johnsongrass and ryegrass annual, perennial	1.8 - 5.5	≤ 25	1.3 - 4.0	< 1
	Kikuyugrass	3.6 - 5.5	≤ 25	2.6 - 4.0	< 1
	Quackgrass and yellow nutsedge	3.6 - 5.5	≤ 25	2.6 - 4.0	< 1

(Continued on next page)

TABLE 2. SPRAY PREPARATION FOR LAWN RENOVATION AND LAWN REPLACEMENT USING POWER SPRAY EQUIPMENT WITH BROADCAST APPLICATIONS (Cont.)

Site	Major Weed Species	Amount of Product, Quarts/Acre	Spray Volume, Gals./Acre	Amount of Product, fl. oz./ 1000 sq. ft.	Spray Volume, Gals./ 1000 sq. ft.
	Other perennial grasses listed on this label	5.5 - 9.5	≤ 25	4.0 - 7.0	< 1

PRECAUTIONS, RESTRICTIONS & COMMENTS FOR LAWN RENOVATION AND LAWN REPLACEMENT

SITE PREPARATION (preplant, preemergence, renovation): This product may be applied before planting or renovation of turf. Applications must be made prior to the emergence of the turf to avoid turf injury. For maximum control of existing vegetation, delay planting to determine if any regrowth from escaped underground plant parts occurs. Where repeat treatments are necessary, sufficient regrowth must be attained prior to application. For warm season grasses, such as bermudagrass, summer or fall applications provide best control.

Do not disturb soil or underground plant parts before treatment.

Tillage or renovation techniques such as vertical mowing, coring or slicing should be delayed for 7 days after application to allow proper translocation into underground plant parts.

This product does not provide residual control.

FARMSTEADS AND FENCEROWS: Pronto® Weed & Grass Killer Concentrate will control kochia, multiflora rose, thistle in areas where total vegetation control is desired. Specific instructions for farmsteads are presented in Table 3.

TABLE 3. SPRAY PREPARATION FOR FARMSTEADS USING POWER SPRAY EQUIPMENT

Site	Major Weed Species	Amount of Product, Quarts/Acre	Spray Volume, Gals./Acre	Amount of Product, fl. oz./1000 sq. ft.	Spray Volume, Gals./1000 sq. ft.
Farmsteads where total vegetation control is desired: around farmstead building foundations, along fences, hedgerows, fencerows, under fencelines, around shelter belts and other area where nonselective weed control is desired.	Annual and perennial weeds, including crabgrass, foxtail, Johnsongrass, kochia, morningglory, multiflora rose, pigweed, trumpetcreeper, thistle, woody brush.	Annual weeds: 1.8 - 3.6	3 - 40	1.3 - 2.6	≤ 1.0
		Perennial weeds: 5.5 - 9.5	3 - 40	4.0 - 7.0	≤ 1.0
		Woody brush & trees: 5.5 - 9.5	3 - 40	4.0 - 7.0	≤ 1.0

See next page for Table 3 – Precautions, Restrictions & Comments

TABLE 3. SPRAY PREPARATION FOR FARMSTEADS USING POWER SPRAY EQUIPMENT (Cont.)

PRECAUTIONS, RESTRICTIONS & COMMENTS:

SITE PREPARATION (preplant, preemergence, renovation):

BARE GROUND: Repeated applications of this product may be used on emerged weeds to maintain bare ground.

NONSELECTIVE POSTEMERGENCE:

Annuals and Perennials: Apply to foliage of emerged weeds. Apply at early growth stages of annual weeds and when perennial weeds are approaching maturity.

Woody brush and trees: Apply this product after full leaf expansion. Use the higher rate for larger plants and/or dense areas of growth. On vines, use the higher rate for plants that have reached the woody stage of growth. Best results are obtained when applications are made in late summer or fall after fruit formation.

In arid areas, best results are obtained when applications are made in the spring to early summer when brush species are at high moisture content and are flowering.

Allow 7 or more days after application before tillage, mowing or removal. Repeat treatments may be necessary to control plants regenerating from underground parts or seed. Reduced performance may result if fall treatments are made following a frost.

GARDENS (VEGETABLES, HERBS, SMALL FRUITS AND BERRIES):

This product may be applied before planting or transplanting of vegetables (garden crops), herbs and small fruits and berries. This product may be used as a broad spectrum postemergent herbicide prior to planting to control annual and perennial weeds including field bindweed, Johnsongrass, or dallisgrass.

Spray the undesirable vegetation on a spray-to-wet basis. Uniform and complete spray coverage of all existing vegetation is required.

RESEEDING AND REPLANTING IN SPRAYED AREAS

ORNAMENTALS AND FLOWERS: Reseed or replant one (1) day after application.

VEGETABLES: Wait three (3) days before planting or transplanting the vegetables listed below:

Beans	Endive
Beet greens	Garlic
Broccoli	Gourds
Brussels sprout	Honeydew melon
Cabbage	Honeyball melon
Cantaloupe	Horseradish
Carrot	Jerusalem artichoke
Cauliflower	Kale
Casaba melon	Kohlrabi
Chard (Swiss)	Leek
Chicory	Lentils
Collards	Lettuce
Corn	Mango melon
Crenshaw melon	Melons
Cucumber	Muskmelon
Eggplant	Mustard greens

Okra	Rutabaga
Onion	Spinach
Rape greens	Summer squash
Parsley	Sweet potato
Parsnips	Tomatillo
Peas	Tomato*
Pepper	Turnip
Persian melon	Watercress
Potato	Watermelon
Pumpkin	Winter squash
Radish	Yams
Rhubarb	

*Tomatoes planted from seed only. Wait thirty (30) days for transplants. For vegetables not listed or for tomato transplants, wait thirty (30) days before planting.

SMALL FRUITS AND BERRIES: Blackberry, black raspberry, blueberry, boysenberry, currant, dewberry, elderberry, gooseberry, huckleberry, loganberry, olallieberry, red raspberry and red raspberry may be planted by seed, rooted cutting, or transplant 7 days after spraying Pronto® Weed & Grass Killer Concentrate. Wait thirty (30) days after application before planting of all other unlisted small fruits and berries.

FRUIT TREES, NUT TREES, CANEBERRIES OR GRAPEVINES: If this product is used to control weeds around fruit or nut trees, caneberreries or grapevines, wait twenty-one (21) days before harvesting and eating the fruits or nuts.

Use the following table to determine the amount of Pronto® Weed & Grass Killer Concentrate to mix with water for use in gardens (vegetables, herbs, small fruits and berries). Close cap tightly after use.

TABLE 4. SPRAY PREPARATION WITH HOSE END SPRAYERS AND TANK SPRAYERS FOR USE IN GARDENS, (VEGETABLES, HERBS, SMALL FRUITS, AND BERRIES)

SPRAYER TYPE	MIXING INSTRUCTIONS		TREAT
	Annuals or General Weed Control	Tough Weed Control Perennial Weeds	
Standard Tank Sprayer	<ul style="list-style-type: none"> • Use 2.25 fl. oz. of Pronto® Weed & Grass Killer Concentrate in 1 gallon of water. 	<ul style="list-style-type: none"> • Use 4.5 fl. oz. of Pronto® Weed & Grass Killer Concentrate in 1 gallon of water. 	300 sq. ft. (30 ft x 10 ft)
Ortho Dial 'N Spray® Hose End Sprayer	<ul style="list-style-type: none"> • Add 2.25 fl. oz. of Pronto® Weed & Grass Killer Concentrate to sprayer jar. • DO NOT ADD WATER. • Set dial to 3 oz. and apply. 	<ul style="list-style-type: none"> • Add 4.5 fl. oz. of Pronto® Weed & Grass Killer Concentrate to sprayer jar. • DO NOT ADD WATER. • Set dial to 5 oz. and apply. 	300 sq. ft.

(Continued on next page)

TABLE 4. SPRAY PREPARATION WITH HOSE END SPRAYERS AND TANK SPRAYERS FOR USE IN GARDENS, (VEGETABLES, HERBS, SMALL FRUITS, AND BERRIES) (Cont.)

SPRAYER TYPE	MIXING INSTRUCTIONS		TREAT
	Annuals or General Weed Control	Tough Weed Control Perennial Weeds	
Ortho Spray-Ette® Hose end Sprayer	<ul style="list-style-type: none"> • Add 8 fl. oz. of Pronto® Weed & Grass Killer Concentrate to sprayer jar. • Fill to 4 gallon mark. 	<ul style="list-style-type: none"> • Add 16 fl. oz. of Pronto® Weed & Grass Killer Concentrate to sprayer jar. • Fill to 4 gallon mark. 	1200 sq. ft. (30 ft x 40 ft)
Hand Trigger Sprayers	<ul style="list-style-type: none"> • Use 0.5 fl. oz. (1 tbs.) of Pronto® Weed & Grass Killer Concentrate in 24 fl. oz. of water. 	<ul style="list-style-type: none"> • Use 1 fl. oz. (2 tbs.) of Pronto® Weed & Grass Killer Concentrate in 24 fl. oz. of water. 	Use for spot treatments or single weeds

LIMITED WARRANTY AND DISCLAIMER

The manufacturer warrants only that the chemical composition of this product conforms to the ingredient statement given on the label, and that the product is reasonably suited for the labeled use when applied according to the Directions for Use.

THE MANUFACTURER NEITHER MAKES NOR INTENDS ANY OTHER EXPRESS OR IMPLIED WARRANTIES, INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, WHICH ARE EXPRESSLY DISCLAIMED.

This limited warranty does not extend to the use of the product inconsistent with label instructions, warnings or cautions, or to use of the product under abnormal conditions such as drought, excessive rainfall, tornadoes, hurricanes, etc. These factors are beyond the control of the manufacturer or the seller. Any damages arising from a breach of the manufacturer's warranty shall be limited to direct damages, and shall not include indirect or consequential damages such as loss of profits or values, except as otherwise provided by law.

The terms of this Limited Warranty and Disclaimer cannot be varied by any written or verbal statements or agreements. No employee or agent of the seller is authorized to vary or exceed the terms of this Limited Warranty and Disclaimer in any manner.

Pronto® is a registered trademark of PBI/GORDON Corporation

Ortho®, Ortho Dial 'N Spray®, and Ortho Spray-Ette® are registered trademarks of Monsanto Company.

**STOP! READ THE ENTIRE LABEL
FIRST. OBSERVE ALL
PRECAUTIONS AND FOLLOW
DIRECTIONS CAREFULLY.**

PRECAUTIONARY STATEMENTS

Hazards to Humans & Domestic Animals

CAUTION: Causes moderate eye irritation. Harmful if absorbed through skin. Harmful if swallowed. Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water.

First Aid

If swallowed: Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.

If on skin or clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

If inhaled: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. Call a poison control center or doctor for treatment advice.

If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor or going for treatment.

PEEL HERE TO OPEN ▲

954/6-2006 AP090805
EPA REG. NO. 2217-846
EPA EST. NO. 2217-KS-2

Manufactured By

**G pbi/gordon
CORPORATION**

An Employee-Owned Company
1217 West 12th Street
Kansas City, Missouri 64101

consumer.pbigordon.com