

PHYSICAL AND CHEMICAL HAZARDS
Do not use or store near heat or open flame.

ENVIRONMENTAL HAZARDS

This product is extremely toxic to fish and aquatic organisms. Do not apply directly to any body of water. Do not contaminate water when disposing of equipment washwater. This pesticide is toxic to honey bees and other beneficial pollinators exposed to an application. Do not apply when bees are actively visiting blooming plants (vegetables, flowers, fruit/ornamental trees). Do not allow this product to come in direct contact with beehives at any time.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

PESTICIDE STORAGE: Store in original container. Keep area inaccessible to children and pets.

DISPOSAL: If empty: Non-refillable container. Do not reuse or refill this container. Offer for recycling, if available. Do not rinse unless required for recycling. If recycling is not available, then dispose of container in a sanitary landfill or, if allowed by State and local authorities, by burning, if burned, stay out of smoke.

If partly filled: Call your local solid waste agency for disposal instructions. Never pour unused product down any indoor or outdoor drain or on the ground.

HOME GARDENS: FOR USE ON HOME GARDENS. NOT FOR USE ON PLANTS BEING GROWN FOR SALE OR OTHER COMMERCIAL USE.

Wearing household latex or rubber gloves, use a dust applicator such as a shaker can, dust gun, puffer or other blubious duster, rotary duster or other appropriate duster as available. Start applying at the farthest corner of the treatment area and work backward to avoid contact with dusted surfaces. Dust lightly to cover both upper and lower leaf surfaces with a thin, even film of dust. (Many insects hide on the underside of the leaf). Apply when air is calm to avoid drift and contact with eyes and skin.

Controls a wide range of insect pests. Use at first signs of insect infestation, preferably before damage from feeding occurs. Note the pre-harvest (time between use and harvest) interval with each crop listed.

ORNAMENTALS: DIRECTIONS FOR APPLICATION

CROP	TARGET PEST	REMARKS
Asparagus Fern	Japanese Beetle, Asparagus Beetle, Lygus Bug	Dust lightly. Apply to the fern stage of asparagus plant after spear harvest when adult stage of the asparagus beetle, lygus bugs and the adult stage of the Japanese beetle are present.
Roses	Beet Armyworm, Cabbage Looper, Omnivorous Leafroller	Dust lightly.

TREE CROPS: DIRECTIONS FOR APPLICATION

Dust lightly, using a dust gun or rotary duster.

CROP	TARGET PEST	REMARKS
Apples	Apple Aphid, Redbanded Leafhopper, Obliquebanded Leafroller, Plum Curculio, White Apple Leafhopper, Spotted Tentiform Leafminer, Tarnished Plant Bug	Dust lightly. Do not apply after petal fall.
Peaches	Peach Twig Borer, Oriental Fruit Moth, Green Fruitworm, Lesser Peachtree Borer, Plum Curculio, Tarnished Plant Bug	Dust lightly. Apply up to 5 times per season. May be applied until 14 days before harvest.
Walnuts	Navel Orangeworm, Codling Moth, Walnut Husk Fly	Dust lightly. Apply up to 8 times per season. May be applied until 1 day before harvest.

WARRANTY STATEMENT

Read the entire Directions for Use and Warranty before using this Product. By using this product, user or buyer accepts the following warranty.

The directions for use of this product are believed to adequate and must be followed carefully. It is impossible to eliminate all risks associated with the use of this product. Unintended consequences may result because of unknown factors. All such risks shall be assumed by the user or buyer.

PBI/Gordon Corporation is committed to providing high quality products. To the extent consistent with applicable law, PBI/Gordon Corporation makes no warranties, express or implied, of merchantability or of fitness for a particular purpose or otherwise, that extend beyond statements on this label. To the extent consistent with applicable law, PBI/Gordon Corporation, the manufacturer, or the Seller shall not be liable for indirect, special, incidental or consequential damages. To the extent consistent with applicable law, the exclusive remedy of the user or buyer shall not exceed the purchase price paid.

GARDEN GUARD® is a registered trademark of PBI/Gordon Corporation.

535/12-2009 AP080808
EPA REG. NO. 53888-33-2217
EPA EST. NO. 2217-KS-2

pbi/gordon corporation
An Employee-Owned Company
DISTRIBUTED BY
PBI/GORDON CORPORATION
1217 WEST 12TH STREET
KANSAS CITY, MISSOURI 64101
www.pbigordon.com



GORDON'S GARDEN GUARD INSECTICIDE

For Outdoor Use on Home Gardens Controls A Wide Range Of Insect Pests On Garden Vegetables As Listed



ACTIVE INGREDIENT:
Permethrin (3-phenoxyphenyl) methyl (+/-) cis, trans-3-(2,2-dichloroethenyl)-2,2-dimethylcyclopropanecarboxylate* 0.25%
INERT INGREDIENTS: 99.75%
*cis:trans ratio. Min. 35% (+/-) cis. Max. 65% (+/-) trans. TOTAL 100.00%

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

See side panel for additional Precautionary Statements.

NET WEIGHT: THREE POUNDS



READ THE ENTIRE LABEL FIRST BEFORE EACH USE. OBSERVE ALL PRECAUTIONS AND FOLLOW DIRECTIONS CAREFULLY.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Humans: Harmful if swallowed or if inhaled. Remove contaminated clothing and wash clothing before reuse. Avoid breathing dust. Avoid contact with eyes or clothing. Wash thoroughly with soap and water after handling and before smoking or eating. Avoid contamination of feed and foodstuffs. **Animals:** Harmful if swallowed, inhaled. Consult a veterinarian before using this product on debilitated, aged, pregnant or nursing animals or animals on medication. Sensitivities may occur after using ANY pesticide product for pets. If signs of sensitivity occur bathe your pet with mild soap and rinse with large amounts of water. If signs continue, consult a veterinarian immediately.

FIRST AID

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-877-800-5556 for emergency medical treatment information.

If in eyes:	<ul style="list-style-type: none"> Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.
If on skin or clothing:	<ul style="list-style-type: none"> Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.
If inhaled:	<ul style="list-style-type: none"> Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control center or doctor for further treatment advice.

VEGETABLES: DIRECTIONS FOR APPLICATION

CROP	TARGET PEST	REMARKS
Asparagus	Cutworm Complex, Asparagus Beetle	Dust lightly. Apply up to 4 times per season. May be applied until 1 day before harvest.
Broccoli, Brussels Sprouts, Cauliflower	Cabbage Looper, Imported Cabbageworm, Diamondback Moth, Cabbage Aphid	Dust lightly. Apply up to 8 times per season. May be applied until 1 day before harvest.
Cabbage	Cabbage Looper, Imported Cabbageworm, Diamondback Moth, Cabbage Aphid	Dust lightly. Apply up to 5 times per season. May be applied until 1 day before harvest.
Cucumbers, Melons, Pumpkins, Squash, Watermelon	Leafminer, Squash Bug, Cabbage Looper, Cucumber Beetle (adult), Cutworm, Melon Worm, Pickle Worm, Plant Bug (including Lygus and Stink Bug), Rindworm, Squash Vine Borer	Dust lightly. Apply up to 8 times per season. May be applied until harvest.
Celery, Lettuce (head and leaf), Parsley, Rhubarb, Spinach	Vegetable Leafminer, Cabbage Looper, Cutworm Complex, Fall Armyworm, Beet Armyworm, Corn Earworm, Leafhopper, European Corn Borer	Dust lightly. Apply up to 5 times per season. May be applied until 1 day before harvest.
Eggplant	Colorado Potato Beetle, Vegetable Leafminer, Cabbage Looper, Flea Beetle	Dust lightly. Apply up to 10 times per season. May be applied until 3 days before harvest.
Onions (Dry bulb and Garlic)	Onion Thrips, Armyworm, Cutworm, Leafminer, Onion Maggot (adult), Stink Bug	Dust lightly. Apply up to 6 times per season. May be applied until 1 day before harvest.
Peppers (Bell)	Pepper Weevil, Vegetable Leafminer, Cabbage Looper, Flea Beetle, European Corn Borer	Dust lightly. Apply up to 8 times per season. May be applied until 3 days before harvest.
Potatoes	Colorado Potato Beetle, Potato Leafhopper, Aster Leafhopper, Potato Flea Beetle, Potato Aphid, Potato Tuberworm, Potato Psyllid, Cabbage Looper	Dust lightly. Apply up to 8 times per season. May be applied until 14 days before harvest.
Sweet Corn	Corn Earworm, European Corn Borer, Fall Armyworm, Southern Armyworm, Corn Rootworm (adult), Cutworm, Aster Leafhopper, Flea Beetle	Dust lightly. Apply up to 6 times per season. May be applied until 1 day before harvest.
Tomatoes	Tomato Fruitworm, Cabbage Looper, Tomato Pinworm, Vegetable Leafminer, Beet Armyworm, Southern Armyworm, Colorado Potato Beetle, Granulate Cutworm, Hornworm	Dust lightly. Apply up to 6 times per season. May be applied until day of harvest.

RECEIVED
JAN 20 2010



STATE OF HAWAII
Department of Agriculture

ACCEPTED
LICENSE NO. 9554.117