



Department of Agriculture
STATE OF HAWAII



LICENSED

PERIOD **2015-2017** LIC. NO.

9554.156

Trimec NUTSEGE

Plus

LAWN WEED KILLER

CONCENTRATE

▶ **KILLS YELLOW NUTSEGE & 200+ LISTED BROADLEAF WEEDS***



Treats Northern & Southern Grasses

ACTIVE INGREDIENTS:

2,4-D, dimethylamine salt	6.10%
Mecoprop-p, dimethylamine salt	2.20%
Dicamba, dimethylamine salt	0.57%
Sulfentrazone	0.15%

OTHER INGREDIENTS: 90.98%

TOTAL 100.00%

See back panel for substatement
of ingredient statement.

*See inside booklet for a complete list of weeds controlled.

See attached booklet for complete Precautionary
Statements and Directions for Use (including
First Aid and Storage and Disposal).

**KEEP OUT OF
REACH OF CHILDREN
CAUTION**

**NET CONTENTS:
ONE PINT (16 Fl. Oz.)**

855/6-2013



Trimec NUTSEGE

Plus

LAWN WEED KILLER

C O N C E N T R A T E



For questions or comments call toll-free **800.884.3179**
Mon.-Fri. 8:00 a.m. - 4:30 p.m. CST



or visit: **GordonsUSA.com**



**pbi / Gordon
CORPORATION**

An Employee-Owned Company

THIS PRODUCT CONTAINS:

- 0.44 lb 2,4-D acid equivalent per gallon or 5.08%
- 0.16 lb Mecoprop-p acid equivalent per gallon or 1.82%
- 0.04 lb Dicamba acid equivalent per gallon or 0.47%
- 0.01 lb Sulfentrazone per gallon or 0.15%

Isomer Specific by AOAC Method.

See inside pages for complete
Precautionary Statements and
Directions for Use (including
First Aid and Storage and Disposal).

855/6-2013 AP082611
EPA REG. NO. 2217-880
EPA EST. NO. 2217-KS-1



(ONE PINT)



Trimec[®] NUTSEGE[®]

Plus[™] LAWN WEED KILLER

CONCENTRATE

Kills Yellow Nutsedge & 200+ Broadleaf Weeds*

Treats Northern & Southern Grasses

ACTIVE INGREDIENTS:

2,4-D, dimethylamine salt	6.10%
Mecoprop-p, dimethylamine salt	2.20%
Dicamba, dimethylamine salt	0.57%
Sulfentrazone	0.15%
OTHER INGREDIENTS:	90.98%
	TOTAL 100.00%

THIS PRODUCT CONTAINS:

- 0.44 lb 2,4-D acid equivalent per gallon or 5.08%
- 0.16 lb Mecoprop-p acid equivalent per gallon or 1.82%
- 0.04 lb Dicamba acid equivalent per gallon or 0.47%
- 0.01 lb Sulfentrazone per gallon or 0.15%

Isomer Specific by AOAC Method.

*As listed in attached booklet.

KEEP OUT OF REACH OF CHILDREN

CAUTION



READ THE ENTIRE LABEL FIRST. OBSERVE ALL PRECAUTIONS AND FOLLOW DIRECTIONS CAREFULLY.

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION: Harmful if swallowed. Causes moderate eye irritation. Avoid contact with eyes or clothing. Wear long-sleeved shirt and long pants, shoes plus socks and gloves. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco.

First Aid	
If swallowed:	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to by a poison control center or doctor. • Do not give anything to an unconscious person.
If in eyes:	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact 1-877-800-5556 for emergency medical treatment advice.	

Environmental Hazards

This pesticide is toxic to fish and aquatic invertebrates and may adversely affect non-target plants. Do not apply directly to water. Do not contaminate water when disposing of equipment washwaters or rinsate.

This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistern or well may result in contamination of drinking water or groundwater.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

SHAKE WELL BEFORE USING.

PRECAUTIONS AND RESTRICTIONS:

- Do not apply this product in a way that will contact any person or pet, either directly or through drift. Keep people and pets out of the area during application.
- Do not allow people or pets to enter the treated area until sprays have dried.

1. Weeds Controlled

GRASSY WEEDS:

Yellow Nutsedge

BROADLEAF WEEDS:

Trimec[®] Nutsedge Plus Lawn Weed Killer Concentrate will control or suppress the following broadleaf weeds.

Broadleaf Weeds		
Amaranth, purple	Fleabane, rough	Pusley, florida
Amaranth, slender	Galinsoga, hairy	Ragweed, large flower
Aster, heath	Galinsoga, smallflower	Ragweed, bur
Aster, white heath	Goldenrod	Ragweed, common
Aster, purple	Ground ivy	Ragweed, lancleaf
Aster, white prairie	Groundsel	Ragweed, western
Artichoke	Gumweed	Redstem filaree
Austrian fieldcress	Hairy fleabane	Rough cinquefoil
Bedstraw, catchweed	Hawkweed, meadow	Scarlet pimpernel
Bedstraw, smooth	Hawkweed, mouseear	Shepherd's purse
Bedstraw, Northern	Hawkweed, orange	Smartweed, ladysthumb
Bedstraw, pineywoods	Hawkweed, yellow	Smartweed, pale
Beggarticks, hairy	Healall	Smartweed, pennsylvania
Betony, Florida	Heartleaf drymary	Smooth chaff-flower
Bindweed, Field	Hemp	Smoothdock
Bindweed, Hedge	Henbit	Sorrel, red (sheep)
Bitter Wintercress	Hoary cress	Sowthistle, annual
Bittercress, hairy	Horsenettle	Sowthistle, spiny
Bitterweed	Horseweed	Spanish needles
Black-eyed Susan	Jimsonweed	Speedwell, bi-lobed
Black Medic	Johnny Jump-up violet	Speedwell, common
Blood flower milkweed	Knawel	Speedwell, corn
Blue lettuce	Knotweed, prostrate	Speedwell, ivyleaf
Brassbuttons	Kochia	Speedwell, persian
Bristly oxtongue	Lambsquarters	Speedwell, purslane
Broomweed	Lespedeza	Speedwell, slender
Buckhorn	Mallow, alkali	Speedwell, snow
Bullnettle	Mallow, bristly	Speedwell, thymeleaf
Burclover	Mallow, common	Spiny amaranth
Burdock	Mallow, dwarf	Spurge, prostrate
Burweed, lawn (spurweed)	Mallow, venice	Spurge, spotted
Buttercup, bulbous	Marcela	Spurweed
Buttercup, creeping	Matchweed	Star of Bethlehem
Buttonweed, common (poorjoe)	Mexicanweed	Strawberry, India mock
Buttonweed, virginia	Moneywort	Tansy ragwort
Carolina geranium (wild)	Morningglory, bigroot	Tanweed
Carpetweed	Morningglory, ivyleaf	Thistle, blessed
Catnip	Morningglory, mexican	Thistle, bull
Catsear, spotted (false dandelion)	Morningglory, tall	Thistle, canada
Chamber bitter	Morningglory, wolly	Thistle, flodman (prairie)
Chickweed, common	Mugwort	Thistle, musk
Chickweed, mouseear	Mustard, ball	Thistle, platte
Chickweed, sticky	Mustard, black	Thistle, plumeless
Chicory	Mustard, blue	Thistle, russian
Cinquefoil	Mustard, hedge	Thistle, scotch
Clover, crimson	Mustard, indian,	Thistle, wavyleaf
Clover, alyce	Mustard, tansy	Thistle, yellow (horrible)
Clover, hop	Mustard, tumble	Trailing crownvetch
Clover, rabbit foot	Mustard, white	Velvetleaf
Clover, red	Mustard, wormseed	Vervain, blue
Clover, strawberry	Nettle, stinging (burning)	Vervain, hoary
Clover, yellow sweet	Nettle, tall	Vervain, prostrate
Clover, white sweet	Nettle, wood	Vervain, tall
Clover, white	Nettle, wood	Vetch, bird
Cockle	Parsley-piert	Vetch, hairy
Cocklebur	Parsnip	Vetch, milk
Cocklebur, oriental	Pearlwort	Vetch, narrowleaf
Cocklebur, spiny	Pennycress (frenchweed)	Vetch, wild
Common mullein	Pepperweed, perinnia	Virginia creeper
Corn Spurry	Pepperweed, virginia (Peppergrass)	Western salsify
Creeping Beggarweed	Pigweed, prostate	Wild aster
Creeping Jenny (Creeping Charlie)	Pigweed, redroot	Wild buckwheat
Cupid shaving Brush	Pigweed, russian	Wild carrot
Cudweed	Pigweed, smooth	Wild four-o'clock
Daisy, English	Pigweed, tumble	Wild garlic
Daisy, oxeye	Pineapple weed	Wild lettuce
Dandelion, common	Plains coreopsis (tickseed)	Wild mustard
Dandelion, false	Plantain, blackseed	Wild mustard (yellow rocket)
Day flower	Plantain, bracted	Wild onion
Deadnettle	Plantain, broadleaf	Wild parsnip
Dichondra	Plantain, buckhorn	Wild radish
Dock, broadleaf	Plantain, hoary	Wild rape
Dock, curly	Plantain, narrowleaf	Wild strawberry
Dogbane	Plantain, slender	Wild sweet potato
Dogfennel	Plantain, woolly	Wild vetch
Dollarweed (Pennywort)	Poison ivy	Wild violet
Dove weed	Poison oak	Woodsorrel, creeping
Elderberry	Pokeweed	Woodsorrel, violet
Falseflax	Prairie Sunflower	Woodsorrel, yellow (Oxalis)
Falseflower	Prickly lettuce (compass plant)	Woolly croton
Fiddleneck	Prickly sida	Wormseed
Filaree, redstem	Prostrate knotweed	Yarrow, common
Filaree, whitestem	Puncturevine	Yarrow, western
Fleabane, daisy (annual)	Purshlane, common	Yarrow, yellow
	Pusley, brazil	

2. Where To Use

This product is intended for use by homeowners on lawns including:

Cool season turfgrass:

- Kentucky bluegrass
- Perennial ryegrass
- Fescue spp., including tall, red and fine leaf fescues
- Bentgrass

Warm season turfgrass:

- Bermudagrass
- Bahiagrass
- Zoysiagrass
- St. Augustinegrass*
- Centipedegrass

*NOTE: Do not use on 'Floritam' or other improved varieties of St. Augustinegrass in Florida. Do not use this product on carpetgrass, dichondra, legumes, or lawns containing desirable clovers.

3. When To Use

Apply when daytime temperatures are between 50° and 85°F

SPRING: Spray when weeds are young and actively growing with adequate soil moisture for best results.

FALL: Spray when weeds are young and actively growing with adequate soil moisture for best results. Fall applications will control weeds that may otherwise go dormant through the winter and resprout the following spring.

Limitations on broadcast applications to residential turfgrass

The maximum application rate is 5.0 fl.oz. of product per 1,000 sq.ft. per application. The maximum number of broadcast applications is limited to 2 per year with a minimum of 30 days between applications. The maximum seasonal rate is 10 fl.oz. of product per 1,000 sq.ft.

3.1 Spot Treatment

To treat small areas or individual weeds use a pressure sprayer for better control. Do not use a hose-end sprayer for Spot Treatments. MIX 1.25 FL.OZ. (2.5 TABLESPOONS) OF PRODUCT IN 1 GALLON OF WATER. Shake well before spraying. Spray the target weeds thoroughly. Nozzle adjustments may be necessary to deliver small spray droplets of the target weeds for uniform coverage.

Limitations on spot treatments:

Spot treatment is defined as a treatment area no greater than 1,000 sq.ft. per acre. The maximum application rate is 5.0 fl.oz. per 1,000 sq.ft. per application. The maximum number of spot treatments is limited to 2 per year with a minimum of 30 days between applications.

3.2 Entire Lawn Treatments

Three easy steps to mix and apply with hand-operated sprayers:

- 1) Measure the total lawn area to be treated. This is done by multiplying the length of the area by the width to obtain total square feet. Prepare only enough spray solution to treat the area calculated.
- 2) Select the application rate from Table 1 below. Carefully measure the proper amount of product to treat 500 sq.ft. If more or less than 500 sq.ft. is to be treated, adjust the amount of product accordingly.
- 3) Shake the spray solution before and during application. Use a spray pattern that gives small droplets without any fine mist.

FOR HAND-OPERATED SPRAYERS (BACKPACK, KNAPSACK, TANK-TYPE, OR PUMP-UP)

Table 1. Amount of product to be diluted in 1 gallon water required to treat 500 sq. ft. of lawn with pressure sprayers.	
Turfgrass	Amount of Product
St. Augustinegrass*, Bermudagrass, Bentgrass, and Centipedegrass	1.25 fl.oz. (2.5 Tbsp)
Kentucky bluegrass, Fescue spp., Perennial ryegrass, Zoysiagrass and Bahiagrass	2.5 fl.oz. (5 Tbsp)

*Do not use on 'Floritam' or other improved varieties of St. Augustinegrass in Florida.
For pressure sprayers, mix the indicated amount of product listed above with 1 gallon of water for each 500 sq.ft. to be treated.

FOR DIAL-IN HOSE END SPRAYERS WITH NO PRE-MIXING

- 1) From the Table below, determine amount of Trimec® Nutsedge Plus Lawn Weed Killer Concentrate to be applied for each 1,000 sq.ft. of lawn.
- 2) Fill sprayer jar with enough concentrate to cover area to be treated. Do not add additional water to the sprayer jar.
- 3) Set dial on sprayer to the "Rate per Gallon" setting from the Table below.
- 4) Attach sprayer to hose. Extend hose to the farthest area of lawn to be treated and work back toward faucet so that you do not walk over treated areas. Apply evenly to lawn.
- 5) Unused product can be returned to the original container when completed.

Table 2. Application rates for turfgrass with hose-end sprayers		
Turfgrass	Dial settings or Rate per gallon	Amount of herbicide applied per 1000 sq. ft.
St. Augustinegrass*, Bentgrass, Bermudagrass, and Centipedegrass	2 Tablespoons (6 teaspoons) (1 ounce)	2.5 fl.oz.
Kentucky bluegrass, Fescue spp., Perennial Ryegrass, Zoysiagrass and Bahiagrass	4 Tablespoons (12 teaspoons) (2 ounces)	5.0 fl.oz.

*Do not use on 'Floritam' or other improved varieties of St. Augustinegrass in Florida.

An example for a hose-end sprayer is shown below:

- For Kentucky bluegrass: To treat 1,000 sq.ft. of lawn, measure 5.0 fl.oz. of product into sprayer jar. **Do not add water.** Set dial on 4 Tbsp and spray uniformly.
- **To BEGIN spraying:** Attach hose to sprayer. Extend the garden hose to the farthest section of lawn to be treated. Rotate sprayer nozzle tip to direct spray up or down. Turn on water and squeeze trigger or rotate on/off knob to begin.
- **AFTER spraying:** Unused product or concentrate can be poured back into the original container. Keep the dial setting and clean the sprayer by rinsing the sprayer jar with water for one minute.

Check the directions you received when you purchased your sprayer for further details on the proper use of your dial sprayer.

FOR CONCENTRATE-PLUS-WATER SPRAYERS OR HOSE-END SPRAYERS WITH PRE-MIXING

For sprayers that require the addition of water as well as concentrate prior to application:

- 1) From the following Table 3, determine amount of concentrate required to treat each 1,000 sq.ft. of lawn.
- 2) Pour concentrate into sprayer jar to the required "fl.oz." level.
- 3) Then add the appropriate amount of water to the "gals." level. Replace sprayer top on jar and shake well before spraying.
- 4) Attach sprayer to hose. Extend hose to the farthest area of lawn to be treated and work back toward faucet so that you do not walk over treated areas. Apply evenly to lawn.

Table 3. Application rates for turfgrass with hose-end sprayers that require pre-mixing.			
Size of area to be treated	St. Augustinegrass, Bermudagrass, Bentgrass, Centipedegrass	Kentucky bluegrass, Fescue spp., Perennial ryegrass, Zoysiagrass and Bahiagrass	Gallons of water
1,000 sq.ft	2.5 fl.oz.	5.0 fl.oz.	2 gal
5,000 sq.ft.	12.5 fl.oz.	25.0 fl.oz.	10 gal
6,000 sq.ft.	15.0 fl.oz.	30.0 fl.oz.	12 gal

*Do not use on 'Floritam' or other improved varieties of St. Augustinegrass in Florida.

EXAMPLE: To treat 1,000 sq.ft. of fescue pour in concentrate to the 5 fl.oz. mark and add water to the 2 gallon mark. Check the directions you received when you purchased your sprayer for further details on the proper use of your sprayer.

4. Use Tips

- 1) Spray during growing season when weeds are actively growing.
- 2) For best results, mow lawn 2 to 3 days before treating weeds.
- 3) To avoid risk of lawn injury, do not spray drought-stressed lawns.
- 4) Do not water within 6 hours after treatment. Poor weed control may result if rainfall occurs within 3 hours of application.
- 5) Do not apply this product to newly seeded grasses until well-established or after the third mowing.
- 6) Treated areas may be reseeded 3 to 4 weeks after application.
- 7) Adjust nozzle to produce a coarse spray. Coarse sprays are less likely to drift. Avoid fine mists.
- 8) If certain tough weeds are not dead within 3 weeks, repeat application using spot treatment directions.

PROHIBITIONS:

- Do not use on 'Floritam' or other improved varieties of St. Augustinegrass in Florida.
- Do not use this product on carpetgrass, dichondra, legumes, or lawns containing desirable clovers.
- Do not treat when air temperatures exceed 85°F as damage to grass may occur.
- Application at rates above specified on this label may cause injury to turfgrass.
- Do not apply this product to food crops (gardens), forage crops, or ornamental plants including flowers, trees, shrubs, hedges, woody ornamentals, or groundcovers established in landscape plantings.
- Do not spray exposed roots of ornamentals and trees.

Other Use Precautions:

- Do not apply as a fine mist because of potential for injury to desirable plants. Do not use this product on or near desirable plants, including within the drip line of the roots of desirable trees and shrubs, since injury may result.
- Do not apply directly to or near water, storm drains, gutters, sewers, or drainage ditches. Do not apply within 25 feet of rivers, fish ponds, lakes, streams, reservoirs, marshes, estuaries, bays, and oceans. Do not apply when windy. To prevent product run-off, do not over water the treated area(s) to the point of runoff or apply when raining or when rain is expected that day. Rinse applicator over lawn area only.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage and disposal.

PESTICIDE STORAGE: Store in original container in a locked storage area inaccessible to children or pets.

PESTICIDE DISPOSAL AND CONTAINER HANDLING: Nonrefillable container. Do not reuse or refill this container. **If empty:** Place in trash or offer for recycling if available. **If partly filled:** Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

LIMITED WARRANTY AND DISCLAIMER

FOR USE ONLY AS DIRECTED. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE MANUFACTURER NEITHER MAKES NOR INTENDS ANY EXPRESS OR IMPLIED WARRANTIES, INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, WHICH ARE HEREBY EXPRESSLY DISCLAIMED. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO CASE SHALL THE MANUFACTURER BE LIABLE FOR INCIDENTAL, CONSEQUENTIAL, OR SPECIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. If these terms are not acceptable, return this product unopened immediately to the point of purchase, and the purchase price will be refunded in full. The terms of this LIMITED WARRANTY AND DISCLAIMER cannot be varied by any written or verbal statements or agreements at the point of sale or elsewhere.

Trimec® is a registered trademark of PBI-Gordon Corporation.

855/6-2013 AP082611
EPA REG. NO. 2217-880

EPA EST. NO. 2217-KS-1

MANUFACTURED BY
PBI/GORDON CORPORATION
1217 WEST 12TH STREET
KANSAS CITY, MISSOURI 64101
www.GordonsUSA.com

STOP! READ THE ENTIRE LABEL FIRST. OBSERVE ALL PRECAUTIONS AND FOLLOW DIRECTIONS CAREFULLY.

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION: Harmful if swallowed. Causes moderate eye irritation. Avoid contact with eyes or clothing. Wear long-sleeved shirt and long pants, shoes plus socks and gloves. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco.

First Aid	
If swallowed:	<ul style="list-style-type: none">• Call a poison control center or doctor immediately for treatment advice.• Have person sip a glass of water if able to swallow.• Do not induce vomiting unless told to by a poison control center or doctor.• Do not give anything to an unconscious person.
If in eyes:	<ul style="list-style-type: none">• Hold eye open and rinse slowly and gently with water for 15-20 minutes.• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.• Call a poison control center or doctor for treatment advice.
Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact 1-877-800-5556 for emergency medical treatment advice.	

855/6-2013 AP082611
EPA REG. NO. 2217-880
EPA EST. NO. 2217-KS-1

MANUFACTURED BY
PBI/GORDON CORPORATION
1217 WEST 12TH STREET
KANSAS CITY, MISSOURI 64101
www.GordonsUSA.com

PEEL HERE TO OPEN ▲