




Department of Agriculture
STATE OF HAWAII

LICENSED

PERIOD 2015-2017 LIC. NO. 9554.165



LIVESTOCK Backrubber & Pour-On



For Use on Beef Cattle, Dairy Cattle, Sheep and Swine.
For Reduction of Flies, Mosquitoes, Ticks and Lice.

ACTIVE INGREDIENT:	
Permethrin*	0.25%
INERT INGREDIENTS**:	99.75%
TOTAL 100.00%	

* *cis/trans* ratio: Min. 35% (±) *cis* and max. 65% (±) *trans*.

** Contains petroleum distillates.

See back panel for additional
Precautionary Statements
and First Aid.

**NET CONTENTS:
2.5 U.S. GALLONS**

**KEEP OUT OF
REACH OF CHILDREN
CAUTION**

939/9-2014



**READ ENTIRE LABEL BEFORE EACH USE.
OBSERVE ALL PRECAUTIONS AND
FOLLOW DIRECTIONS CAREFULLY.**

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION: Harmful if absorbed through skin. Avoid contact with skin, eyes, and clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet.

FIRST AID

If swallowed:	<ul style="list-style-type: none"> • Immediately call a poison control center or doctor. • Do not induce vomiting unless told to do so by a poison control center or doctor. • Do not give any liquid to the person. • Do not give anything by mouth to an unconscious person.
If on skin or clothing:	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.
If in eyes:	<ul style="list-style-type: none"> • Hold eyelid open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment.

Note to Physician: Contains petroleum distillates – vomiting may cause aspiration pneumonia.

Environmental Hazards

This pesticide is extremely toxic to aquatic organisms, including fish and invertebrates. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area.

This pesticide is highly toxic to bees exposed to direct treatment on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds while bees are actively visiting the treatment area.

Physical or Chemical Hazards

Do not use or store near heat or open flame.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

FOR APPLICATION TO BEEF CATTLE, DAIRY CATTLE, CALVES, SHEEP AND SWINE: To reduce annoyance from flies, mosquitoes, lice, keds (sheep-tick and louse fly), and ticks by the following application methods.

BACKRUBBER METHOD: Fill reservoir or saturate applicator. Recharge as needed to maintain effective results. Locate mechanism to give animals free access.

POUR ON TECHNIQUE: Apply undiluted to beef cattle, dairy cattle, calves and sheep along back and down face at ½ to 2 fl. oz. per 100 lbs. body weight of animal up to a maximum of 20 fl. oz. per animal. For Keds (sheep-tick and louse fly) on sheep, pour along backline over neck, shoulders and rump, applying 1 fl. oz. per 50 lbs. body weight of animal, up to a maximum of 12 fl. oz. for any one animal. Use as needed but not more than once every 2 weeks. Treat lactating dairy cows after milking is completed.

NOTE: This product is not effective for grub control on cattle.

FOR APPLICATION TO SWINE: Use ½ fl. oz. per 100 lbs. of body weight of animal up to a maximum of 4.0 fl. oz. per animal. Use as needed but not more than once every 2 weeks. Do not slaughter swine within 5 days of last treatment.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

PESTICIDE STORAGE: Store in a cool, dry, and secure area. Store in original sealed container away from food or feed.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER HANDLING: Nonrefillable container: Do not reuse or refill this container. If empty: Place in trash or offer for recycling if available. If partly filled: Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

LIMITED WARRANTY AND DISCLAIMER

The manufacturer warrants only that the chemical composition of this product conforms to the ingredient statement given on the label, and that the product is reasonably suited for the labeled use when applied according to the Directions for Use.

THE MANUFACTURER NEITHER MAKES NOR INTENDS ANY EXPRESS OR IMPLIED WARRANTIES, INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, WHICH ARE EXPRESSLY DISCLAIMED. This limited warranty does not extend to the use of the product inconsistent with label instructions, warnings or cautions. Any damages arising from a breach of the manufacturer's warranty shall be limited to direct damages, and shall not include indirect or consequential damages such as loss of profits or values, except as otherwise provided by law.

The terms of this Limited Warranty and Disclaimer cannot be varied by any written or verbal statements or agreements. No employee or agent of the seller is authorized to vary or exceed the terms of this Limited Warranty and Disclaimer in any manner.

939/9-2014 AP111813
EPA REG. NO. 2217-861
EPA EST. NO. 2217-KS-2



An Employee-Owned Company

MANUFACTURED BY
PBI/GORDON CORPORATION
1217 WEST 12TH STREET
KANSAS CITY, MISSOURI 64101
www.pbigordon.com

