



STATE OF HAWAII
Department of Agriculture

ACCEPTED

LICENSE NO. **9554.82**

13361 Amine 400 1-G. (Floater) FAF 10/10/06 11:09 AM Page 1



AMINE 400 2,4-D WEED KILLER

Controls most common broadleaf weeds in Pastures & Rangelands,
Corn, Sorghum, Wheat, and Barley.

Also, for use in

- Lawns • Roadsides • Drainage Ditchbanks • Rights-of-Way • Railroads
- Fencerows • Industrial Sites

ACTIVE INGREDIENT:

*Dimethylamine salt of
2,4-dichlorophenoxyacetic
acid 46.4%

INERT INGREDIENTS: 53.6%
TOTAL 100.0%

This Product Contains:

*3.8 lbs. 2,4-dichlorophenoxyacetic acid
equivalent per gallon or 38.6%.
Isomer Specific by AOAC Methods.

814/10-2006 AP052005
EPA REG. NO. 2217-2
EPA EST. NO. 2217-KS-1



NET CONTENTS: ONE U.S. GALLON

**KEEP OUT OF REACH OF CHILDREN
DANGER – PELIGRO**

Si Usted no entiende la etiqueta, busque a
alguien para que se la explique a Usted en
detalle. (If you do not understand the label,
find someone to explain it to you in detail.)

**See inside pages for additional
Precautionary Statements and First Aid.**



0 70251 81472 5



**READ THE ENTIRE LABEL FIRST.
OBSERVE ALL PRECAUTIONS AND
FOLLOW DIRECTIONS CAREFULLY.**

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

DANGER: Corrosive. Causes irreversible eye damage. Causes skin irritation. Do not get in eyes, on skin or clothing. This product is harmful or fatal if swallowed.

Personal Protective Equipment (PPE):

Applicators and other handlers must wear long-sleeved shirt and long pants, waterproof gloves, shoes plus socks, and protective eyewear.

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

Engineering Control Statements:

Containers over 1 gallon and less than 5 gallons: Mixers and loaders who do not use a mechanical system (probe and pump) to transfer the contents of this container must wear coveralls or a chemical-resistant apron in addition to the other required PPE.

Containers of 5 gallons or more: Do not open-pour product from this container. A mechanical system (such as a probe and pump or spigot) must be used for transferring the contents of this container. If the contents of a non-refillable pesticide container are emptied, the probe must be rinsed before removal. If the mechanical system is used in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d) (4)], the handler PPE requirements may be reduced or modified as specified in the WPS.

When handlers use enclosed cabs or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d) (4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

User Safety Recommendations:

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

FIRST AID

<p>If in eyes:</p>	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
<p>If on skin or clothing:</p>	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.
<p>If swallowed:</p>	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by a poison control center or doctor. • Do not give anything by mouth to an unconscious person. <p><i>(cont. on next page)</i></p>

FIRST AID (cont.)**If inhaled:**

- Move person to fresh air.
- If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible.
- Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact 1-877-800-5556 for emergency medical treatment advice.

NOTE TO PHYSICIAN: Probable mucosal damage may contraindicate the use of gastric lavage.

ENVIRONMENTAL HAZARDS: Do not apply when weather conditions favor drift from target area. Do not contaminate domestic or irrigation waters.

This product is toxic to aquatic invertebrates. Drift or runoff may adversely affect aquatic invertebrates and non-target plants. For terrestrial uses, do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwater.

Most cases of groundwater contamination involving phenoxy herbicides such as 2,4-D have been associated with mixing/loading and disposal sites. Caution should be exercised when handling 2,4-D pesticides at such sites to prevent contamination of groundwater supplies. Use of closed systems for mixing or transferring this pesticide will reduce the probability of spills. Placement of the mixing/loading equipment on an impervious pad to contain spills will help prevent groundwater contamination.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application.

For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard 40 CFR part 170.

This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 48 hours.

For early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, wear coveralls, waterproof gloves, socks plus shoes, and protective eyewear.

STORAGE & DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

PESTICIDE STORAGE: Store in original container in a locked storage area inaccessible to children or pets. Keep from freezing.

PESTICIDE DISPOSAL: Pesticides wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law and may contaminate groundwater. If these wastes cannot be

disposed of by use according to label instructions, contact your state Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL: For plastic containers: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke. For metal containers: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

USE PRECAUTIONS

Do not apply this product through any type of irrigation system.

Do not apply in tank mixtures with other 2,4-D products.

Do not overdose! Avoid spray drift to cotton, soybeans, tomatoes, tobacco, grapes, fruit trees, flowers, garden crops, ornamental plants, shrubs, trees and other sensitive plants. Do not apply near these plants since small quantities of wind drifted herbicide may cause severe injury.

Do not apply when wind speed is sufficient to cause drift. Do not apply when an air temperature inversion exists. An inversion may be detected by creating a smoke column and observing a layering effect. Do not apply when temperatures exceed 90°F and humidity is high.

To aid in avoiding spray drift, use coarse sprays and low pressure. Do not use nozzles which produce fine spray droplets under high pressure. The use of thickening agents or antidrift additives and drift reducing equipment is of value in preventing spray drift. Care should be taken not to make applications where runoff could carry the chemical to nontarget areas. Local spray conditions will vary. Check local recommendations first.

WEEDS CONTROLLED

Annual and Biennial Weeds:

Beggarticks	Morningglory (annual)
Bitterweed	Musk thistle
Broomweed	Mustard
Bull thistle	Parsnip
Burdock	Pennycress
Carpetweed	Peppergrass
Cinquefoil	Pigweed
Cockle	Prickly lettuce
Cocklebur	Primrose
Coffeeweed	Puncturevine
Croton	Radish (wild)
Devil's claw	Ragweed (common)
Fleabane (daisy)	Russian thistle
Flixweed	Shepherdspurse
Frenchweed	Smartweed
Galinsoga	Sneezeweed
Goatsbeard	Sowthistle (common)
Goosefoot	Spanishneedles
Gumweed	Sunflower
Jewelweed	Tumbleweed
Jimsonweed	Velvetleaf
Kochia	Vervain
Knotweed	Vetch
Lambsquarters	Wild carrot
Lettuce (wild)	Wild parsnip
Mallow	Witchweed
Marshelder	Wormwood
Marijuana	Yellow starthistle

Perennial Weeds:

Artichoke	Dandelion
Aster	Dock
Austrian fieldcress	Dogbane
Bindweed	Goldenrod
Blackeyed Susan	Ground ivy
Blue lettuce	Heal-all
Canada thistle	Hoary cress
Catnip	Horsetail
Chicory	Ironweed
Clover (many types)	<i>(cont. on next page)</i>

WEEDS CONTROLLED *(cont.)*

Perennial Weeds *(cont.)*:

Locoweed	Ragweed	Strawberry (wild)	Yellow rocket
Nettle	Rushes	Tall buttercup	Wild garlic
Orange hawkweed	Southern wild rose	Tanweed	Wild onion
Plantain	Southistle	Toadflax	Wild sweet potato
Povertyweed	Stinging nettle	Vervain	

FOR USE IN SMALL GRAINS

CROPS	DOSAGE	SPRAY	INSTRUCTIONS
	RATE PTS./ACRE	VOLUME GALS./ACRE	
Wheat, Barley, Rye, Oats, Winter Grains: Annual and biennial weeds	½ to 2*	8 or more	Apply after crop is fully tillered (about 4" to 8" high) and before jointing. Do not spray crop in boot to dough stage. Oats are more sensitive to 2,4-D than other crops and should be sprayed in spring when well established and tillered and before jointing; (use ½-1 pint per acre). Do not spray crop in boot to dough stage.
Perennial broadleaf weeds	1 to 2*	8 or more	Apply when weeds are near bud stage. Do not spray crop in boot to dough stage.
Spring Grains: Annual broadleaf weeds	½ to 2*	8 or more	Apply after crop is fully tillered (about 4" to 8" high) and before jointing. Do not spray crop in boot to dough stage.
Perennial broadleaf weeds	1 to 2*	8 or more	Apply when weeds are near bud stage. Do not spray crop in boot to dough stage.

***NOTES ABOUT THE ABOVE:** Use the lower rate if small annual and biennial weeds are the major problems. Use the higher rate if the weeds present are in the hard-to-control categories as determined by local experience. The higher rates increase the risk of crop injury and should be used only where the weed control problem justifies the crop damage risk.

Do not forage or graze treated fields within 2 weeks after treatment.

For Aerial Applications on Cereal Grains — Spray volumes of 3 to 10 gallons of water per acre are appropriate.

FOR EMERGENCY WEED CONTROL IN WHEAT

Perennial broadleaf weeds: Apply 3 pints per acre when weeds are approaching bud stage, but do not spray grain in the boot to dough stage. The application rate of 3 pints of product (1.5 lb. acid equivalent) per acre can produce injury to wheat. Balance the severity of your weed problem against the possibility of crop damage. Where perennial weeds are scattered, spot treatment is suggested to minimize the extent of crop injury. Do not feed treated straw to livestock.

FOR USE IN ROW CROPS, PASTURES, AND RANGELANDS

CROPS	DOSAGE RATE PTS./ACRE	SPRAY VOLUME GALS./ACRE	INSTRUCTIONS
Corn (Field & Sweet): Preemergence Postemergence annual broadleaf weeds Perennial broadleaf weeds	2 to 3	15 to 30	Apply just before corn emerges.
	½ to 1	8 to 15	Apply when most weeds have germinated. Corn is susceptible to injury at time of emergence and shortly after unfolding of leaves. Do not spray during this period. Do not spray corn during the tassel to hard dough stages. Use drop nozzles when corn is 10" tall to place spray below its tops. Do not cultivate soon after spraying while corn is brittle. Do not forage or feed corn fodder for 7 days following application.
	1 to 1½	8 to 15	Apply when weeds are in bud to bloom stage. Use drop nozzles after corn is 10" tall. Do not spray corn during the tassel to hard dough stages. 2,4-D may cause brittleness to corn. Winds or cultivation may cause stalk breakage while brittle. Certain single cross corn hybrids may be more susceptible to 2,4-D injury than others.
Grain Sorghum Postemergence	1	6 to 10	Apply when sorghum is 4" to 12" tall. Use drop nozzles to keep spray off sorghum plants, when sorghum is over 10" high. Do not forage or feed fodder for 7 days following application.
Rice	1½ to 2½	5 to 10	To control curly indigo and other broadleaf weeds, apply 7 to 10 weeks after planting when rice is fully tillered. Do not spray rice in boot stage.
Sugarcane Preemergence Postemergence	4	15 to 20	Apply before canes appear for control of emerged broadleaf weeds.
	1½ to 2	10 to 30	Apply after cane emerges and through lay-by.
Rangelands, Pastures & Turf Areas: Golf Courses, Parks, Cemeteries (Do not use on golf greens nor on dichondra or other broadleaf herbaceous groundcovers)	2 to 4*	3 to 60	Use the lower rate if annual and biennial weeds are the major problem, the higher rate for perennial weeds. The maximum application rate to pasture and rangeland is 2 pounds 2,4-D acid equivalent per acre per application per site. Do not apply to newly seeded turf until grass has been cut several times. Where bentgrass predominates, make 2 applications of 1 pint per acre at 3 week intervals. Do not use on susceptible southern grasses such as St. Augustinegrass, bentgrass, dichondra, and clover may be injured by this treatment. Observe the following intervals: a. A 7-day grazing interval after treatment for dairy cattle; b. A 30-day preharvest interval for grass cut for hay; and c. A preslaughter interval for meat animals of 3 days.

*Use 2 or more gallons of water per acre for aerial application.

**To convert local recommendations into terms of
Amine 400 2,4-D Weed Killer use the following table.**

2,4-D Acid Equivalent	1 lb.	¾ lb.	½ lb.	¾ lb.	¼ lb.	½ lb.	¼ lb.
Amine 400 2,4-D Weed Killer	2 pt.	1½ pt.	1 pt	¾ pt.	½ pt.	¾ pt.	¼ pt.

FOR USE IN REDUCED OR NO-TILLAGE SOYBEANS (Preplant Only)

GENERAL INFORMATION

AMINE 400 2,4-D WEED KILLER is a phenoxy type herbicide that provides postemergence control of many susceptible annual and perennial broadleaf weeds. AMINE 400 2,4-D WEED KILLER may be applied prior to planting soybeans to provide foliar burndown control of susceptible annual and perennial broadleaf weeds and certain broadleaf cover crops such as those listed on this label. AMINE 400 2,4-D WEED KILLER should only be applied preplant to soybeans in situations such as reduced tillage production systems, where emergent weeds are present. Apply only according to the application instructions.

MIXING INSTRUCTIONS

Mix AMINE 400 2,4-D WEED KILLER only with water, unless otherwise directed on this label. Compatible crop oil concentrates, agricultural surfactants and fluid fertilizers approved for use on growing crops may increase the herbicidal effectiveness of 2,4-D on certain weeds and may be added to the spray tank. Read and follow all directions and precautions on this label and on all labels of adjuvants or fertilizers mixed with this product.

APPLICATION PROCEDURES

Apply using air or ground equipment in sufficient gallonage to obtain adequate coverage of weeds. Use 2 or more gallons of water per acre in aerial equipment and 10 or more gallons of water per acre in ground equipment.

APPLICATION TIMING AND USE RATES FOR AMINE SALTS

Maximum Amount Of Amine 400 to Apply/acre	Maximum Rate (Pounds 2,4-D a.e./acre)	When to Apply (Days Prior To Planting Soybeans)
1 Pint	0.5	NOT LESS THAN 15 DAYS
1 Quart	1.0	NOT LESS THAN 30 DAYS

WEEDS CONTROLLED

Alfalfa*	Dandelion	Mousetail	Smartweed,
Bindweed*	Dock, curly*	Mustard, wild	Pennsylvania
Bullnettle	Eveningprimrose, cutleaf	Onion, wild*	Sowthistle, annual
Bittercress, smallflowered	Garlic, wild*	Pennycress, field	Speedwell
Buttercup, smallflowered	Horseweed or Maretail	Plantain	Thistle, Canada*
Carolina geranium	Ironweed	Purslane, common	Thistle, bull
Cinquefoil, common and rough	Lambsquarters, common	Ragweed, common	Velvetleaf
Clover, red*	Lettuce, prickly	Ragweed, giant	Vetch, hairy*
Cocklebur, common	Morningglory, annual	Shepherdspurse	Virginia copperleaf

*These species are only partially controlled.

In general, weeds should be small, actively growing and free of stress caused by extremes in climatic conditions, diseases, or insect damage at the time of treatment. The response of individual weed species to AMINE 400 2,4-D WEED KILLER is variable. Consult your local county or State Agricultural Extension Service or crop consultant for advice.

APPLICATION RESTRICTIONS AND PRECAUTIONS FOR SOYBEANS (Preplant Only)

Important Notice: Unacceptable injury to soybeans planted in fields previously treated with AMINE 400 2,4-D WEED KILLER may occur. Whether or not soybean injury occurs and the extent of the injury will depend on weather (temperature and rainfall) from herbicide application until soybean emergence and agronomic factors such as the amount of weed vegetation and previous crop residue present. Injury is more likely under cool rainy conditions and where there is less weed vegetation and crop residue present.

Apply a maximum of one application per growing season regardless of the treatment rate.

Do not apply AMINE 400 2,4-D WEED KILLER when weather conditions such as air temperature inversions or wind favor drift from treated areas to susceptible plants.

Livestock Grazing Restrictions: Do not feed hay, forage, or fodder. Restrict livestock from grazing treated fields. Livestock should be restricted from feeding/grazing of treated cover crops.

In fields previously treated with 2,4-D, plant soybean seed as deep as practical or at least 1.5 to 2.0 inches deep. Adjust the press wheel of the planter, if necessary, to ensure that planted seed is completely covered.

WEEDS & BRUSH ON IRRIGATION CANAL DITCHBANKS

Seventeen Western States: Arizona, California, Colorado, Idaho, Kansas, Montana, Nebraska, New Mexico, Nevada, North Dakota, Oklahoma, Oregon, South Dakota, Texas, Utah, Washington and Wyoming.

For control of annual and perennial broadleaf weeds, apply 1 to 2 quarts of product per acre in approximately 20 to 100 gallons of water per acre. Treat when weeds are young and actively growing before the bud or early bloom stage. For harder-to-control weeds a repeat spray after 3 to 4 weeks using the same rates may be needed for maximum results. Apply no more than two treatments per season.

For woody brush and perennial broadleaf weeds, mix 1 gallon of product in 150 gallons of water. Wet foliage thoroughly using about one gallon of solution per square rod.

Spraying Instructions: Apply with low pressure (10 to 40 psi) power spray equipment mounted on a truck, tractor or boat. Apply while traveling upstream to avoid accidental concentration of chemical into water. Spray when the air is fairly calm; 5 mph or less. Do not use on small canals (less than 10 cfs) where water will be used for drinking purposes.

Boom spraying onto water surface must be held to a minimum and no cross-stream spraying to opposite banks should be permitted. When spraying shoreline weeds, allow no more than two-foot overspray to prevent introduction of greater than negligible amounts of chemical into the water.

Do not allow dairy animals to graze on treated areas for at least 7 days after spraying. Water within treated banks should not be fished.

CONTROL OF WOODY PLANTS OR BRUSH AND BROADLEAF WEEDS ON ROADSIDES, DRAINAGE DITCHBANKS, RIGHTS-OF-WAY, RAILROADS, FIREBREAKS, FENCEROWS, INDUSTRIAL SITES AND OTHER SIMILAR NONCROPLAND AREAS

DIRECTIONS, RESTRICTIONS AND LIMITATIONS FOR USE IN NONCROPLAND

Broadcast applications to annual and perennial weeds: Apply to emerged weeds. For best results, treat when weeds are young and actively growing.

The maximum application rate to general noncropland sites is 0.5 gallon (4 pints) of product per acre per application per site.

When multiple applications of up to 2.0 lbs. acid equivalent per acre are utilized to reach the maximum seasonal use rate, do not make a repeat application within 30 days of the previous application.

Minimum spray volume: Use 2 or more gallons of spray solution per acre.

Number of applications: Limited to 2 applications per year.

Broadcast applications to woody plants: Apply to trees and brush when foliage is fully expanded and plants are actively growing.

Up to 1.0 gallon of product per acre (4.0 lbs. acid equivalent per acre) may be applied in a single application to rights-of-way, including electrical power lines, communication lines, pipelines, highways and railroads that intersect wooded areas or stands of trees, brush and woody plants.

The maximum noncropland application rate for trees, brush and woody plant control is 1.0 gallon of product per acre per application per site.

Target species	Application schedule	Maximum application rate, Gallons of product per acre	Maximum application rate, Pounds of acid equivalent per acre per application	Maximum number of applications per year	Minimum days between applications	Minimum spray volume, gallons per acre
Annual and perennial weeds	Broadcast	0.5 gal./A or 4 pints/A	2.0 #/A	2	30 days	2
Woody plants	Broadcast and high volume foliar	1.0 gal./A or 8 pints/A	4.0 #/A	1	N/A	See Tables 1-2.

High volume foliar applications (100-400 gallons per acre): Apply 0.25-1.0 gallon of product per acre with adequate water or apply a 0.25-1.0% vol/vol spray solution as a full cover spray with high volume equipment. Use the lower spray concentrations in the range for susceptible species and use the higher spray concentrations within the range for hard-to-control species, for mature plants during the late summer or under adverse environmental conditions (e.g. drought).

Spray broadleaf weeds, woody plants or mixed brush uniformly and thoroughly by wetting all leaves, stems, bark and root collars. The total volume of spray solution required for adequate coverage of solid stands of mixed brush can range from 100-400 gallons of spray solution per treated acre. The spray preparation chart for applications on a spray-to-wet basis is shown below in Table 1.

Table 1. Instructions for preparing 100-400 gallons of spray solution at 0.25-1.0% spray concentration with water for high volume foliar applications.

Spray solution per acre, Gallons	Amount of Product Needed for Spray Concentration of:			
	0.25%	0.33%	0.5%	1.0%
100	0.25 gal.	0.33 gal.	0.5 gal.	1.0 gal.
200	0.5 gal.	0.67 gal.	1.0 gal.	—
300	0.75 gal.	1.0 gal.	—	—
400	1.0 gal.	—	—	—

Equal measures: 1 gallon = 4 quarts = 8 pints = 128 fl. oz.

The maximum seasonal application rate for trees, brush and woody plant control is 1.0 gallon of product per acre per application per site.

For backpack sprayers, knapsack sprayers, and hand-pressurized pump sprayers**Table 2.** Instructions for preparing 1-3 gallons of spray solution at 0.25-1.0% spray concentration with water for high volume foliar applications.

Gallons Of Water	Amount of Product Needed for Spray Concentration of:			
	0.25%	0.33%	0.5%	1.0%
1	2 teaspoons	3 teaspoons	4 teaspoons	8 teaspoons
2	4 teaspoons	2 tablespoons	3 tablespoons	6 tablespoons
3	2 tablespoons	3 tablespoons	4 tablespoons	8 tablespoons

Equal measures: 1 fl. oz. = 2 Tablespoons (Tbs.) = 6 Teaspoons (tsp.)

The maximum seasonal application rate for trees, brush and woody plant control is 1.0 gallon of product per acre per application per site.

TANK MIXTURES FOR NONCROPLAND

Utility & Pipeline Rights-of-Way — Use ½ to 1 gallon of AMINE 400 2,4-D WEED KILLER in tank mix combinations with other herbicides labeled for rights-of-way sites and apply in a total spray volume of 5 to 30 gallons per acre.

AMINE 400 2,4-D WEED KILLER can be applied as a tank mixture with other recommended herbicides such as Garlon®, Tordon® and Banvel® to broaden the spectrum of control. In order to assure maximum safety and weed control, follow all precautions and limitations on this label and the labels of products used in tank mixtures with AMINE 400 2,4-D WEED KILLER. Where a rate range is given, the rate should be varied according to the density and target species.

PRODUCTS	RATES
AMINE 400 + Garlon® 3A Herbicide	½-1 gal./A + ½-1 gal./A
AMINE 400 + Garlon® 4E Herbicide	½-1 gal./A + 2-4 qts./A
AMINE 400 + Tordon® K Herbicide	½-1 gal./A + ½-4 qts./A
AMINE 400 + Banvel® Herbicide	½-1 gal./A + 1 qt.-2 gal./A

FORESTS (Forest Site Preparation):**AERIAL APPLICATIONS**

Forestry Site Preparation — For use in desiccation/controlled burning programs, use ½ to 1.0 gallon of AMINE 400 2,4-D WEED KILLER per acre in tank mixes with other herbicides labeled for forestry site preparation (e.g. Garlon®, Tordon®, Arsenal®). Use sufficient water to achieve uniform wetting of target brush species. Do not exceed 25 gallon total spray per acre.

Do Not Apply as a stand release or cover spray established conifers as injury may result.

The maximum application rate to forestry site preparation is 4 pounds 2,4-D acid equivalent per acre per application per site. Seasonal: The maximum seasonal application rate to forestry sites is 4 pounds 2,4-D acid equivalent per acre per application per site.

FORESTRY — TREE INJECTION

Make injections as near the root collar as possible using one injection per inch of trunks dbh (4½ feet). For resistant species such as hickory, injections should overlap. For best results, injections should be made during the growing season — May 15 to October 15.

For Dilute Injection: Mix 1 gallon of product in 19 gallons water for dilute injections.

For Concentration Injection: Use 1 to 2 ml of concentrate per injection. The injector bit must penetrate the inner bark.

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to use of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Do not allow people (other than applicator) or pets on treatment area during application. Do not enter treatment area until spray has dried or dust has settled.

HOME LAWNS – FOR USE ON RESIDENTIAL AND OTHER TURF SITES EXCLUDING SOD FARMS

To control weeds in established lawns and other ornamental turfgrasses such as bluegrass, perennial ryegrass, and fescue, mix according to the following dilution chart and apply during active weed growth. Apply in spring, summer or fall when weeds are actively growing. Spray to give a uniform application. Delay mowing for several days before and after treatment. Do not use on newly seeded areas or on grass seedlings. Do not use on new lawns until mowed twice. Creeping grasses such as zoysiagrass, bermudagrass, St. Augustinegrass, dichondra, and clovers may be injured severely by this product; only spot treat weeds in these types of grasses.

Do not use on golf greens, dichondra, or other broadleaf herbaceous groundcovers.

Reseeding of treated areas should be delayed following treatment. With spring application reseed in fall; with fall application reseed in the spring.

Deep rooted perennials may require repeat applications.

The maximum application rate to turf is 1.4 pounds 2,4-D acid equivalent per acre per application per site. The maximum number of broadcast applications per treatment site is 2 per year.

TO PREPARE THE SPRAY

- 1) Fill the sprayer about one-half full with water.
- 2) Add the required amount of AMINE 400 2,4-D WEED KILLER.
- 3) Add the rest of the water and mix well.

FOR BROADCAST TREATMENTS IN LAWNS AND TURF		
AREA TO BE TREATED	AMOUNT OF AMINE 400 2,4-D TO USE	AMOUNT OF WATER TO USE
500 sq. ft.	1 tablespoon (½ fl. oz.)	2 gal.
1,000 sq. ft.	2 tablespoons (1 fl. oz.)	4 gal.
5,000 sq. ft.	9 tablespoons (4½ fl. oz.)	20 gal.

For Spot Treatment Only In Lawns & Turf

For spot treatment only in lawns and turf, mix 4 fl. oz. of AMINE 400 2,4-D WEED KILLER in 2 to 3 gallons of water and spray the weeds when they are small. Be sure to thoroughly wet all foliage.

NEVER USE EATING, DRINKING, OR HOUSEHOLD KITCHEN UTENSILS FOR MIXING OR MEASURING PESTICIDES.

LIMITED WARRANTY AND DISCLAIMER

The manufacturer warrants only that the chemical composition of this product conforms to the ingredient statement given on the label, and that the product is reasonably suited for the labeled use when applied according to the Directions for Use.

THE MANUFACTURER NEITHER MAKES NOR INTENDS ANY OTHER EXPRESS OR IMPLIED WARRANTIES, INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, WHICH ARE EXPRESSLY DISCLAIMED.

This limited warranty does not extend to the use of the product inconsistent with label instructions, warnings or cautions, or to use of the product under abnormal conditions such as drought, excessive rainfall, tornadoes, hurricanes, etc. These factors are beyond the control of the manufacturer or the seller. Any damages arising from a breach of the manufacturer's warranty shall be limited to direct damages, and shall not include indirect or consequential damages such as loss of profits or values, except as otherwise provided by law.

The terms of this Limited Warranty and Disclaimer cannot be varied by any written or verbal statements or agreements. No employee or agent of the seller is authorized to vary or exceed the terms of this Limited Warranty and Disclaimer in any manner.

Garlon® and Tordon® are registered trademarks of Dow AgroSciences LLC.

Arsenal® and Banvel® are registered trademarks of BASF Corporation.

814/10-2006 AP052005
EPA REG. NO. 2217-2
EPA EST. NO. 2217-KS-1

Manufactured By

 An Employee-Owned Company
 1217 West 12th Street
 Kansas City, Missouri 64101
www.pbigordon.com

THIS PAGE BLANK INTENTIONALLY.

THIS PAGE BLANK INTENTIONALLY.

THIS PAGE BLANK INTENTIONALLY.



AMINE 400

2,4-D WEED KILLER

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

DANGER: Corrosive. Causes irreversible eye damage. Causes skin irritation. Do not get in eyes, on skin or clothing. This product is harmful or fatal if swallowed.

Personal Protective Equipment (PPE):

Applicators and other handlers must wear long-sleeved shirt and long pants, waterproof gloves, shoes plus socks, and protective eyewear.

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

Engineering Control Statements:

Containers over 1 gallon and less than 5 gallons: Mixers and loaders who do not use a mechanical system (probe and pump) to transfer the contents of this container must wear coveralls or a chemical-resistant apron in addition to the other required PPE.

When handlers use enclosed cabs or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d) (4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

User Safety Recommendations:

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

FIRST AID

If in eyes:	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
If on skin or clothing:	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice. <p style="text-align: right;"><i>(cont. on next column)</i></p>

FIRST AID (cont.)

If swallowed:	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by a poison control center or doctor. • Do not give anything by mouth to an unconscious person.
If inhaled:	<ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. • Call a poison control center or doctor for treatment advice.
<p>Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact 1-877-800-5556 for emergency medical treatment advice.</p> <p>NOTE TO PHYSICIAN: Probable mucosal damage may contraindicate the use of gastric lavage.</p>	

ENVIRONMENTAL HAZARDS: Do not apply when weather conditions favor drift from target area. Do not contaminate domestic or irrigation waters.

This product is toxic to aquatic invertebrates. Drift or runoff may adversely affect aquatic invertebrates and nontarget plants. For terrestrial uses, do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwater.

Most cases of groundwater contamination involving phenoxy herbicides such as 2,4-D have been associated with mixing/loading and disposal sites. Caution should be exercised when handling 2,4-D pesticides at such sites to prevent contamination of groundwater supplies. Use of closed systems for mixing or transferring this pesticide will reduce the probability of spills. Placement of the mixing/loading equipment on an impervious pad to contain spills will help prevent groundwater contamination.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application.

For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

Manufactured By



An Employee-Owned Company
1217 West 12th Street
Kansas City, Missouri 64101

www.pbigordon.com

814/10-2006 AP052005
EPA REG. NO. 2217-2
EPA EST. NO. 2217-KS-1

PEEL HERE TO OPEN