



STATE OF HAWAII
Department of Agriculture

ACCEPTED

LICENSE NO. **9554.84**

GORDON'S®

TRIMEC®

**READY SPRAY
LAWN WEED KILLER**

**Just Connect To Hose
And Spray!**

ACTIVE INGREDIENTS:

2,4-D, dimethylamine salt	5.56%
Mecoprop-p, dimethylamine salt	1.34%
Dicamba, dimethylamine salt	0.62%

INERT INGREDIENTS: 92.48%

TOTAL 100.00%

THIS PRODUCT CONTAINS:

4.62% 2,4-dichlorophenoxyacetic acid equivalent.
 1.11% (+)-(R)-2-(2-methyl-4-chlorophenoxy) propionic acid equivalent.
 0.51% 3,6-dichloro-o-anisic acid equivalent.

Isomer Specific by AOAC Method.

Contains the single isomer form of Mecoprop-p.

2,4-D dimethylamine salt (CAS Registry No. 2008-39-1)

Mecoprop-p, dimethylamine salt (CAS Registry No. 66423-09-4)

Dicamba, dimethylamine salt (CAS Registry No. 2300-66-5)

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

See attached booklet for additional Precautionary Statements.

827/10-2006



**NET CONTENTS:
ONE QUART (32 FL. OZ.)**

GORDON'S®
TRIMEC®
READY SPRAY
LAWN WEED KILLER

**JUST CONNECT TO GARDEN HOSE
AND SPRAY! COVERS UP TO
5,000 SQ. FT. OF NORTHERN GRASSES.**

**Peel open for
Precautionary Statements
and Full Directions for Use.**

827/10-2006 AP100906
EPA REG. NO. 2217-540
EPA EST. NO. 2217-KS-2

Manufactured By

G **pbi / GORDON**
CORPORATION

An Employee-Owned Company
1217 West 12th Street
Kansas City, Missouri 64101

www.pbigordon.com



32 FL. OZ.

GORDON'S®
TRIMEC®
READY SPRAY
LAWN WEED KILLER

**JUST CONNECT TO GARDEN HOSE
 AND SPRAY! COVERS UP TO
 5,000 SQ. FT. OF NORTHERN GRASSES.**

ACTIVE INGREDIENTS:

2,4-D, dimethylamine salt	5.56%
Mecoprop-p, dimethylamine salt	1.34%
Dicamba, dimethylamine salt	0.62%

INERT INGREDIENTS: 92.48%

TOTAL 100.00%

THIS PRODUCT CONTAINS:

- 4.62% 2,4-dichlorophenoxyacetic acid equivalent.
- 1.11% (+)-(R)-2-(2-methyl-4-chlorophenoxy) propionic acid equivalent.
- 0.51% 3,6-dichloro-o-anisic acid equivalent.

Isomer Specific by AOAC Method.

Contains the single isomer form of Mecoprop-p.

2,4-D dimethylamine salt (CAS Registry No. 2008-39-1)

Mecoprop-p, dimethylamine salt (CAS Registry No. 66423-09-4)

Dicamba, dimethylamine salt (CAS Registry No. 2300-66-5)

KEEP OUT OF REACH OF CHILDREN
CAUTION

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not allow people (other than applicator) or pets on treatment area during application. Do not enter treatment area until spray has dried or dust has settled.

STORAGE: Keep from freezing. Store in original container in a locked storage area inaccessible to children and pets. Do not store near food or feedstuffs.

DISPOSAL:

If Empty – Do not reuse this container. Place in trash or offer for recycling if available.

If Partly Filled – Call your local solid waste agency or toll free number (1-800-CLEANUP) for disposal instructions. Never place unused product down any indoor or outdoor drain.

This product is designed for use through this sprayer for controlling of broadleaf weeds in bahiagrass, bermudagrass, Kentucky bluegrass, fescue, perennial ryegrass, zoysiagrass, centipedegrass, and St. Augustinegrass lawns, vacant lots, playgrounds and other similar areas.

Temporary injury may occur on bentgrass, bermudagrass, and red fescue lawn grasses. Do not apply to carpetgrass, dichondra, Lippia, and lawns where desirable clovers are present.

Water lawn thoroughly 1 to 2 days before application.

Spray when air is calm to avoid spray drift that might injure desirable ornamental plants. Adjust water pressure at faucet to give coarse spray which is less likely to drift. Do not apply when temperatures are expected to exceed 85°F within 24 hours following application.

Newly seeded lawns may be treated after they have been mowed at least three times.

WHERE TO USE:

Trimec® Ready Spray Lawn Weed Killer may be used on residential and ornamental lawns, vacant lots, playgrounds and other similar areas where broadleaf weeds are present. Established turf of bahiagrass, bermudagrass, Kentucky bluegrass, centipedegrass, fescues, perennial ryegrass, St. Augustinegrass, and zoysiagrass may be treated. This product is especially effective on all hard-to-kill weeds such as poison ivy, poison oak, sheep sorrel and spurge. In addition, Trimec® Ready Spray Lawn Weed Killer controls most other broadleaf weeds, including sensitive weeds such as chickweed, clover, dandelion, knotweed and plantain. A partial list is given on a following panel.

OTHER USE PRECAUTIONS:

Do not allow drift onto vegetables, flowers, ornamental plants, shrubs, trees and other desirable plants since injury may result. Do not pour spray solutions near these plants.

Whether using a hose end sprayer or pressure sprayer, do not use sprays that are fine because they can drift and cause damage to nontarget plants. Coarse droplet sprays are less likely to drift. Use only lawn-type sprayers.

Do not spray on carpetgrass, dichondra, Lippia, nor on lawns or turf where desirable clovers are present. Do not spray roots of trees and ornamentals.

Do not exceed specified dosages for any area; be particularly careful within dripline of tree and other ornamental species.

This product may be injurious to St. Augustinegrass and centipedegrass if misapplications occur. Do not apply this product to 'Floritam' St. Augustinegrass in Florida.

The maximum application rate to turf is 2.0 pounds 2,4-D acid equivalent per acre per application per site. The maximum number of broadcast applications per treatment site is 2 per year.

WHEN TO USE:

1. Spray at any time during the growing season when weeds are actively growing.
2. On new lawns, wait until the grass has reached a height of 2 inches before making an overall spray treatment. However, a spot treatment may be made at any time with a small pressure sprayer. Wait 3 weeks after treatment before seeding grass seed or planting sprigs or stolons.
3. Poor weed control may result if spray is applied during drought or just before rain.
4. For best results, mow before treatment.
5. Do not water lawn within 24 hours after treatment.

HOSE END SPRAYERS**HOW TO APPLY:**

Shake well before applying. Connect spray nozzle to hose. Remove bottle cap and screw bottle onto nozzle. Turn the "ON/OFF" switch on top of the nozzle right to left to "OFF". Turn on the water at the tap. When ready to treat, turn the "ON/OFF" switch to its center "ON" position. The water mixes automatically with the product at the proper ratio. Walk and spray at a steady pace. Each mark on the side of bottle covers 1,000 to 1,200 square feet.

Use a hose end sprayer designed for home lawn care. Refer to Charts 1 and 2 for the proper amounts of Trimec® Ready Spray Lawn Weed Killer. Determine the area of turf to be treated. Pour the recommended amount

of product (Column 1) into the bottle, add water to the indicated level (Column 2), and spray uniformly on turf area shown in Column 3.

CHART 1: Amounts of Trimec® Ready Spray Lawn Weed Killer for bahiagrass, bermudagrass, Kentucky bluegrass, fescues, perennial ryegrass, and zoysiagrass lawns applied with hose end sprayers.

Column 1		Column 2	Column 3
Amount of Lawn Weed Killer to Use		Fill Bottle to, Gallons	Area to Treat, Square Feet
(Tablespoons)	(Fluid Ounces)		
10 to 12	5 to 6	5	950
20 to 24	10 to 12	10	1,900
30 to 36	15 to 18	15	2,850
40 to 48	20 to 24	20	3,800

Note: For these grasses apply the entire contents of this container (32 fluid ounces) on 5,000 to 6,000 square feet of turf.

CHART 2: Amounts of Trimec® Ready Spray Lawn Weed Killer for centipedegrass and St. Augustinegrass lawns applied with hose end sprayers.

Column 1		Column 2	Column 3
Amount of Lawn Weed Killer to Use		Fill Bottle to, Gallons	Area to Treat, Square Feet
(Tablespoons)	(Fluid Ounces)		
6	3	5	950
12	6	10	1,900
18	9	15	2,850
24	12	20	3,800

Note: For St. Augustinegrass and centipedegrass apply the entire contents of this container (32 fluid ounces) on 10,000 square feet of turf.

PRESSURE SPRAYERS, BACKPACK SPRAYERS, OR OTHER MANUALLY OPERATED SPRAYERS:

Prepare the appropriate amounts of spray mixtures according to Chart 3. Measure the recommended amount of Trimec® Ready Spray Lawn Weed Killer into the spray tank and dilute with the proper amount of water. Use a spray pattern with small droplets and treat the area indicated in Chart 3.

Chart 3.

A. For bahiagrass, bermudagrass, Kentucky bluegrass, fescues, perennial ryegrass, and zoysiagrass lawns.			
Amount of Lawn Weed Killer to Use		Amount of Water in Sprayer, Gallons	Area to Treat, Square Feet
(Tablespoons)	(Fluid Ounces)		
2 to 2.4	1 to 1.2	¾	190
4 to 4.8	2 to 2.4	1½	380
8 to 9.6	4 to 4.8	3	760

B. For centipedegrass and St. Augustinegrass lawns.			
Amount of Lawn Weed Killer to Use		Amount of Water in Sprayer, Gallons	Area to Treat, Square Feet
(Tablespoons)	(Fluid Ounces)		
2 to 2.4	1 to 1.2	¾	380
4 to 4.8	2 to 2.4	1½	760
8 to 9.6	4 to 4.8	3	1,520

SPOT SPRAYING:

To spray individual weeds, do not use a hose end sprayer. Mix 1 fluid ounce (2 Tablespoons) of Trimec® Ready Spray Lawn Weed Killer in ½ gallon of water in a pressure sprayer. Spray to wet individual weeds.

GENERAL BRUSH CONTROL:

For control of brush along roads and other similar noncrop areas, mix 10 to 14 fluid ounces of Trimec® Ready Spray Lawn Weed Killer in 3 gallons of water and apply over 1,000 square feet. Use the higher rate of product for larger plants or more dense brush stands. Refer to Table 1 for additional instructions for spray preparations with water.

Table 1. Mixing instructions for preparing 1 to 9 gallons of spray solution with water for foliar application.

Spray Solution Desired, Gallons	Amount of Trimec® Ready Spray Lawn Weed Killer Required	
	Normal Rate	High Rate (For larger plants or more dense brush)
0.25 (32 fl.oz.)	0.83 fl.oz.	1.16 fl.oz.
1	3½ fl.oz.	4¾ fl.oz.
2	6¾ fl.oz.	9½ fl.oz.
3	10 fl.oz.	14 fl.oz.
4	13½ fl.oz.	18¾ fl.oz.
5	16¾ fl.oz.	23½ fl.oz.
6	20 fl.oz.	28 fl.oz.
7	23½ fl.oz.	32¾ fl.oz.
8	26¾ fl.oz.	37½ fl.oz.
9	30 fl.oz.	42 fl.oz.
Equal measures: 1.0 fl.oz. = 2 tablespoons of product 1.0 quart = 32 fl.oz. = 2 pints = 4 cups		

Best results are obtained when brush is young and actively growing. Do not cut brush until the herbicide has translocated throughout the plant causing root death. Keep out of lakes, streams and ponds.

CONTROLS:**BRUSH AND VINES:**

Alder	Bush morningglory	Giant ragweed
Ash	Buttonbush	Gooseberry
Aspen	Cedar	Hackberry
Bindweed	Chamise	Hazel
Birch	Cherry, wild	Hemp
Black cherry	Coastal sage	Himalayan blackberry
Black locust	Cottonwood	Honeylocust
Blackberry	Cucumber tree	Honeysuckle
Brambles	Dogwood	Honeysuckle, Japanese
Buckbrush	Elderberry	Kudzu
Buffalo berry	Elm	Lilac, common

Marijuana	Poison ivy	Sumac, dwarf black
Manzanita	Poison oak	Sycamore
Matrimony vine	Poplar	Tamarisk
Moonseed vine	Raspberry	Thistle
Mountain ash	Redvine	Trumpet Vine
Multiflora rose	Sagebrush	Trumpet creeper
Oak	Saltbush	Virginia creeper
Olive, Russian	Serviceberry	Wild plum
Peppervine	Skunkbrush	Willow
Persimmon	Snowberry	Winterberry & other brush, trees & vines
Pine	Spruce	
Pine, shortleaf	Sumac	

HARDY WEEDS:

For hardy weeds such as clover, ground ivy, oxalis, sheep sorrel, dock, thistles, and morningglory, and Virginia buttonweed in lawns and other areas, repeat application in two to three weeks if necessary.

CONTROLS:**SENSITIVE WEEDS:**

Bedstraw	Dandelion	Purslane
Black medic	Knotweed	Ragweed
Buckhorn	Lambsquarters	Shepherdspurse
Chickweed	Peppergrass	Wild lettuce
Clover	Plantain	

HARD-TO-KILL WEEDS:

Dock	Oxalis	Spurge
Ground ivy	Pigweed	Virginia buttonweed
Healall	Poison ivy	Wild carrot
Lespedeza	Poison oak	Wild onion
Mallow	Sheep sorrel	Yarrow & many other broadleaf weeds
Morningglory	Speedwell	

OTHER WEEDS CONTROLLED:

Aster	Bird vetch	Black mustard
Austrian fieldcress	Bitter wintercress	Blackseed plantain
Bedstraw	Bittercress, hairy	Blessed thistle
Beggarticks	Bitterweed	Bloodflower milkweed
Betony, Florida	Black-eyed Susan	Blue lettuce
Bindweed, field	Black medic	Blue vervain

Bracted plantain	Dogfennel	Oriental cocklebur
Bristly oxtongue	Dollarweed	Oxalis
Broadleaf dock	False dandelion	Parsley-piert
Broadleaf plantain	Falseflax	Parsnip
Broomweed	False sunflower	Pearlwort
Buckhorn plantain	Fiddleneck	Pennycress
Bulbous buttercup	Florida pusley	Pennywort
Bull thistle	Frenchweed	Pepperweed
Bullnettle	Galinsoga	Pigweed
Burclover	Goldenrod	Pineywoods bedstraw
Burdock	Ground ivy	Plains coreopsis
Burning nettle	Gumweed	Plantain
Burweed	Hairy fleabane	Poison ivy
Buttercup	Hawkweed	Poison oak
Buttonweed	Healall	Pokeweed
Canada thistle	Heartleaf drymary	Poorjoe
Carolina geranium	Heath aster	Prairie sunflower
Carpetweed	Hedge bindweed	Prickly lettuce
Catchweed bedstraw	Hedge mustard	Prickly sida
Catnip	Henbit	Prostrate knotweed
Catsear	Hoary cress	Prostrate pigweed
Chickweed, common	Hoary plantain	Prostrate spurge
Chickweed, mouseear	Hoary vervain	Prostrate vervain
Chicory	Horsenettle	Puncturevine
Cinquefoil	Jimsonweed	Purslane, common
Clover, crimson	Knawel	Ragweed
Clover, hop	Knotweed	Red sorrel
Clover, red	Kochia	Redroot pigweed
Clover, strawberry	Lambsquarters	Redstem filaree
Clover, sweet	Lespedeza	Rough cinquefoil
Clover, white	Mallow	Rough fleabane
Cockle	Matchweed	Russian pigweed
Cocklebur	Mexicanweed	Russian thistle
Common mullein	Milk vetch	Scarlet pimpernel
Creeping jenny	Morningglory	Scotch thistle
Cudweed	Mouseear hawkweed	Sheep sorrel
Curly dock	Mugwort	Shepherdspurse
Daisy, English	Musk thistle	Slender plantain
Daisy, fleabane	Mustard	Smallflower galinsoga
Daisy, oxeye	Narrowleaf plantain	Smooth dock
Dandelion	Narrowleaf vetch	Smooth pigweed
Dichondra	Nettle	Sorrel
	Orange hawkweed	Sowthistle

Spanishneedles	Velvetleaf	Wild parsnip
Speedwell	Venice mallow	Wild radish
Spiny amaranth	Virginia buttonweed	Wild rape
Spiny cocklebur	Virginia creeper	Wild strawberry
Spiny sowthistle	Virginia pepperweed	Wild sweet potato
Spotted catsear	Wavyleaf bullthistle	Wild vetch
Spotted spurge	Western clematis	Woodsorrel
Spurweed	Western salsify	Woolly croton
Stinging nettle	White mustard	Woolly morningglory
Strawberry, India mock	Wild aster	Woolly plantain
Tall nettle	Wild buckwheat	Wormseed
Tall vervain	Wild carrot	Yarrow
Tansy ragwort	Wild four-o'clock	Yellowflower
Tansy mustard	Wild garlic	pepperweed
Tanweed	Wild geranium	Yellow rocket and other
Thistle	Wild lettuce	broadleaf weeds
Trailing crownvetch	Wild marigold	
Tumble mustard	Wild mustard	
Tumble pigweed	Wild onion	

PRECAUTIONARY STATEMENTS

Hazards to Humans & Domestic Animals

CAUTION: Harmful if swallowed. Avoid contact with eyes, skin or clothing. Wear face shield or goggles when mixing, loading or applying this product. When using this product, wear long-sleeved shirt, long pants, socks, shoes and rubber gloves.

After using this product, rinse gloves before removing, remove clothing and launder separately before reuse, and promptly and thoroughly wash hands and exposed skin with soap and water. Remove saturated clothing as soon as possible and shower.

FIRST AID

If inhaled:

- Move person to fresh air.
- If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible.
- Call a poison control center or doctor for further treatment advice.

(Continued on next page)

FIRST AID (continued)

If on skin or clothing:	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.
If in eyes:	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
If swallowed:	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by a poison control center or doctor. • Do not give anything by mouth to an unconscious person.
<p>Have the product container or label with you when calling a poison control center or doctor or going for treatment.</p> <p>You may also call toll-free 1-877-800-5556 for emergency medical treatment information.</p>	

ENVIRONMENTAL HAZARDS:

This product is toxic to aquatic invertebrates. Drift or runoff may adversely affect aquatic invertebrates and nontarget plants. Do not apply directly to water. Do not contaminate water when disposing of equipment washwater. When cleaning equipment, do not pour the washwater on the ground; spray or drain over a large area away from wells and other water sources.

Most cases of groundwater contamination involving phenoxy herbicides such as 2,4-D and MCP-p have been associated with mixing/loading and disposal sites. Caution should be exercised when handling 2,4-D and MCP-p pesticides at such sites to prevent contamination of groundwater supplies. Use of closed systems for mixing and transferring this pesticide will reduce the probability of spills. Placement of the mixing/loading equipment on an impervious pad to contain spills will help prevent groundwater contamination.

LIMITED WARRANTY AND DISCLAIMER

The manufacturer warrants only that the chemical composition of this product conforms to the ingredient statement given on the label, and that the product is reasonably suited for the labeled use when applied according to the Directions for Use.

THE MANUFACTURER NEITHER MAKES NOR INTENDS ANY OTHER EXPRESS OR IMPLIED WARRANTIES, INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, WHICH ARE EXPRESSLY DISCLAIMED. This limited warranty does not extend to the use of the product inconsistent with label instructions, warnings or cautions, or to use of the product under abnormal conditions such as drought, excessive rainfall, tornadoes, hurricanes, etc. These factors are beyond the control of the manufacturer or the seller. Any damages arising from a breach of the manufacturer's warranty shall be limited to direct damages, and shall not include indirect or consequential damages such as loss of profits or values, except as otherwise provided by law.

The terms of this Limited Warranty and Disclaimer cannot be varied by any written or verbal statements or agreements. No employee or agent of the seller is authorized to vary or exceed the terms of this Limited Warranty and Disclaimer in any manner.

TRIMEC® is a registered trademark of PBI/Gordon Corporation.

827/10-2006 AP100906
EPA REG. NO. 2217-540
EPA EST. NO. 2217-KS-2

Manufactured By



An Employee-Owned Company

1217 West 12th Street
Kansas City, Missouri 64101

www.pbigordon.com

GORDON'S
TRIMEC® *Ready-Spray Lawn Weed Killer*

**STOP! READ THE ENTIRE LABEL FIRST.
OBSERVE ALL PRECAUTIONS AND FOLLOW DIRECTIONS CAREFULLY.**

PRECAUTIONARY STATEMENTS

Hazards to Humans & Domestic Animals

CAUTION: Harmful if swallowed. Avoid contact with eyes, skin or clothing. Wear face shield or goggles when mixing, loading or applying this product. When using this product, wear long-sleeved shirt, long pants, socks, shoes and rubber gloves.

After using this product, rinse gloves before removing, remove clothing and launder separately before reuse, and promptly and thoroughly wash hands and exposed skin with soap and water. Remove saturated clothing as soon as possible and shower.

ENVIRONMENTAL HAZARDS:

This product is toxic to aquatic invertebrates. Drift or runoff may adversely affect aquatic invertebrates and nontarget plants. Do not apply directly to water. Do not contaminate water when disposing of equipment washwater. When cleaning equipment, do not pour the washwater on the ground; spray or drain over a large area away from wells and other water sources.

Most cases of groundwater contamination involving phenoxy herbicides such as 2,4-D and MCPP-p have been associated with mixing/loading and disposal sites. Caution should be exercised when handling 2,4-D and MCPP-p pesticides at such sites to prevent contamination of groundwater supplies. Use of closed systems for mixing and transferring this pesticide will reduce the probability of spills. Placement of the mixing/loading equipment on an impervious pad to contain spills will help prevent groundwater contamination.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not allow people (other than applicator) or pets on treatment area during application. Do not enter treatment area until spray has dried or dust has settled.

STORAGE: Keep from freezing. Store in original container in a locked storage area inaccessible to children and pets. Do not store near food or feedstuffs.

DISPOSAL:

If Empty – Do not reuse this container. Place in trash or offer for recycling if available.

If Partly Filled – Call your local solid waste agency or toll free number (1-800-CLEANUP) for disposal instructions. Never place unused product down any indoor or outdoor drain.

827/10-2006 AP100906
EPA REG. NO. 2217-540
EPA EST. NO. 2217-KS-2

Manufactured By

G **pbi / GORDON**
CORPORATION

An Employee-Owned Company
1217 West 12th Street
Kansas City, Missouri 64101

www.pbigordon.com

▲
PEEL HERE TO OPEN