



STATE OF HAWAII
Department of Agriculture
ACCEPTED
LICENSE NO.
9554.85

Trimec[®]

LAWN WEED KILLER

▶ **KILLS OVER 230 LISTED WEEDS***

▶ **WON'T HARM LAWNST[†]**



C O N C E N T R A T E

Treats up to 32,000 sq. ft. (Northern grasses) 64,000 sq. ft. (Southern grasses)

ACTIVE INGREDIENTS:

2,4-D, dimethylamine salt	7.59%
Mecoprop-p, dimethylamine salt	1.83%
Dicamba, dimethylamine salt	0.84%
OTHER INGREDIENTS:	89.74%
TOTAL	100.00%

THIS PRODUCT CONTAINS:
 0.94 lb. 2,4-dichlorophenoxyacetic acid equivalent per gallon or 6.30%,
 0.13 lb. (+)-(R)-2-(2-methyl-4-chlorophenoxy)propionic acid equivalent
 per gallon or 1.51%, 0.06 lb. 3,6-dichloro-o-anisic acid equivalent
 per gallon or 0.70%. Isomer Specific by AOAC Method.
 2,4-D, dimethylamine salt (CAS Registry No. 2008-39-1)
 Mecoprop-p, dimethylamine salt (CAS Registry No. 66423-09-4)
 Dicamba, dimethylamine salt (CAS Registry No. 2300-66-5)

*See attached booklet for complete
Precautionary Statements and
Directions for Use (including First Aid,
and Storage and Disposal).*

*See inside booklet for a complete list of weeds controlled
†When used as directed

**NET CONTENTS:
ONE GALLON**

**KEEP OUT OF
REACH OF CHILDREN
CAUTION**



Trimec[®]

LAWN WEED KILLER

→ Getting Started

- **Kills:** Dandelion, clover, thistle, and over 230 other weeds listed.
- **Where To Use:** On established lawns without harming lawn grasses when used as directed (do not use on carpetgrass, improved varieties of St. Augustinegrass, such as 'Floritam', dichondra or desirable clovers).
- **Amount To Use:**
Bahiagrass, Bermudagrass, Fescues, Kentucky Bluegrass, Perennial Ryegrass & Zoysiagrass:
Entire Lawn Treatments: Mix 4 fl. oz. in 1 gallon of water to treat 1,000 sq. ft.
Centipedegrass & St. Augustinegrass:
Entire Lawn Treatments: Mix 2 fl. oz. in 1 gallon of water to treat 1,000 sq. ft.
- **Coverage:** This one gallon bottle treats 32,000 sq. ft. of Bahiagrass, Bermudagrass, Fescues, Kentucky Bluegrass, Perennial Ryegrass & Zoysiagrass or 64,000 sq. ft. of Centipedegrass & St. Augustinegrass.



For questions or comments call toll-free **800.884.3179**
Mon.-Fri. 8:00 a.m. - 4:30 p.m. CST



or visit: **GordonsUSA.com**



pbi / Gordon
CORPORATION

An Employee-Owned Company

809/2-2015 AP061214
EPA REG. NO. 2217-539-33955
EPA EST. NO. 2217-KS-1

See inside pages for complete
Precautionary Statements
and Directions for Use
(including First Aid and
Storage and Disposal).



ONE GALLON



Trimec[®]

LAWN WEED KILLER

ACTIVE INGREDIENTS:

2,4-D, dimethylamine salt	7.59%
Mecoprop-p, dimethylamine salt	1.83%
Dicamba, dimethylamine salt	0.84%
OTHER INGREDIENTS:	89.74%
	TOTAL 100.00%

THIS PRODUCT CONTAINS:

- 0.54 lb 2,4-dichlorophenoxyacetic acid equivalent per gallon or 6.30%
- 0.13 lb (+)-(R)-2-(2-methyl-4-chlorophenoxy)propionic acid equivalent per gallon or 1.51%
- 0.06 lb 3,6-dichloro-o-anisic acid equivalent per gallon or 0.70%

Isomer Specific by AOAC Method.

- 2,4-D, dimethylamine salt (CAS Registry No. 2008-39-1)
- Mecoprop-p, dimethylamine salt (CAS Registry No. 66423-09-4)
- Dicamba, dimethylamine salt (CAS Registry No. 2300-66-5)

KEEP OUT OF REACH OF CHILDREN CAUTION



**READ THE ENTIRE LABEL FIRST.
OBSERVE ALL PRECAUTIONS AND
FOLLOW DIRECTIONS CAREFULLY.**

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION: Harmful if absorbed through skin. Harmful if swallowed. Avoid contact with skin, eyes, or clothing. Causes moderate eye irritation. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet. Wear long-sleeved shirt, long pants, socks, shoes and gloves. Remove and wash contaminated clothing before reuse.

First Aid	
If in eyes:	<ul style="list-style-type: none">• Hold eye open and rinse slowly and gently with water for 15 - 20 minutes.• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.• Call a poison control center or doctor for treatment advice.
If swallowed:	<ul style="list-style-type: none">• Call a poison control center or doctor immediately for treatment advice.• Have person sip a glass of water if able to swallow.• Do not induce vomiting unless told to do so by the poison control center or doctor.• Do not give anything by mouth to an unconscious person.
If on skin or on clothing:	<ul style="list-style-type: none">• Take off contaminated clothing.• Rinse skin immediately with plenty of water for 15 - 20 minutes.• Call a poison control center or doctor for treatment advice.
If inhaled:	<ul style="list-style-type: none">• Move person to fresh air.• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible.• Call a poison control center or doctor for treatment advice.
HOT LINE NUMBER: Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact 1-877-800-5556 for emergency medical treatment information.	

Environmental Hazards

This pesticide is toxic to fish and aquatic invertebrates and may adversely affect non-target plants. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Rinsing application equipment over the treated area will help avoid run off to water bodies or drainage systems.

This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistern or well may result in contamination of drinking water or groundwater.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

1. Use Restrictions

- Do not apply this product in a way that will contact any person or pet, either directly or through drift. Keep people and pets out of the area during application. Do not allow people or pets to enter the treated area until sprays have dried.
- Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.
- Use on established lawns. Will not harm lawn grasses when used as directed (Do not use on carpetgrass, improved varieties of St. Augustinegrass, such as 'Floritam,' dichondra or desirable clovers)
- The turf tolerance of centipedegrass and St. Augustinegrass can range from sensitive to tolerant. Do not apply this product to 'Floritam' St. Augustinegrass in Florida.
- Do not apply as a fine mist because of potential for injury to desirable plants.
- Do not use this product on or near desirable plants, including contact of spray on exposed root systems or adventitious shoots within the drip line of desirable trees and shrubs, since injury may result.
- Do not apply directly to or near water, storm drains, gutters, sewers, or drainage ditches. Do not apply within 25 feet of rivers, fish ponds, lakes, streams, reservoirs,

marshes, estuaries, bays, and oceans. Do not apply when windy. To prevent product run-off, do not over water the treated area(s) to the point of runoff or apply when raining or when rain is expected that day. Rinse sprayer over lawn area only.

2. How Much To Apply

Bahiagrass, Bermudagrass, Fescues, Kentucky Bluegrass, Perennial Ryegrass & Zoysiagrass:

Spot Treatments: Mix 1 fl.oz. (2 Tbsp) in 1 gallon of water to treat 250 sq.ft.

Entire Lawn Treatments:

- Pump Style Sprayer: Mix 4 fl.oz. in 0.5 to 1 gallon of water to treat 1,000 sq.ft.
- Hose-End Sprayer: Apply 4 fl.oz. per 1,000 sq.ft.
- 12 Volt & Tow-Behind Sprayers: Mix 4 fl.oz. in 0.5 to 1 gallon of water to treat 1,000 sq.ft.

Table 1: Spot Treatment of Bahiagrass, Bermudagrass, Fescues, Kentucky Bluegrass, Perennial Ryegrass, & Zoysiagrass

Tank Size	Concentrate to add to tank	Coverage Area
1 gal	1 fl.oz.	250 sq.ft.

Table 2: Entire Lawn Treatment of Bahiagrass, Bermudagrass, Fescues, Kentucky Bluegrass, Perennial Ryegrass, & Zoysiagrass

Tank Size	Concentrate to add to tank	Coverage Area
1 gal	4 fl.oz.	1,000 sq.ft.
15 gal	60 fl.oz.	15,000 sq.ft.
25 gal	100 fl.oz.	25,000 sq.ft.

Centipedegrass & St. Augustinegrass:

Spot treatments: Mix 0.5 fl.oz. (1 Tbsp) in 1 gallon of water to treat 250 sq.ft.

Entire Lawn Treatments:

- Pump Style Sprayer: Mix 2 fl.oz. in 0.5 to 1 gallon of water to treat 1,000 sq.ft.
- Hose-End Sprayer: Apply 2 fl.oz. per 1,000 sq.ft.
- 12 Volt & Tow-Behind Sprayers: Mix 2 fl.oz. in 0.5 to 1 gallon of water to treat 1,000 sq.ft.

Table 3: Spot Treatment of Centipedegrass & St. Augustinegrass		
Tank Size	Concentrate to add to tank	Coverage Area
1 gal	1 fl.oz.	500 sq.ft.

Table 4: Entire Lawn Treatment of Centipedegrass & St. Augustinegrass		
Tank Size	Concentrate to add to tank	Coverage Area
1 gal	4 fl.oz.	2,000 sq.ft.
15 gal	60 fl.oz.	30,000 sq.ft.
25 gal	100 fl.oz.	50,000 sq.ft.

3. How To Apply

3.1 Pump Style Sprayers

Ideal for spraying individual weeds

1. Add the appropriate amount of water and concentrate to the sprayer tank.
2. Close sprayer, shake well, and pump handle to pressurize.
3. Adjust nozzle to deliver a coarse spray pattern.
4. To broadcast over large areas: apply evenly to treated area.
5. To treat individual weeds: spray to wet weed leaf surfaces.
6. Re-pressurize the sprayer as needed to maintain a good spray pattern.

3.2 12 Volt & Tow-Behind Boom Sprayers

Ideal for treating large country and suburban lawns

1. To calibrate your application equipment consult your equipment owner's manual for speed and pressure settings required to deliver 0.5 to 1 gallon of spray solution to treat 1,000 sq.ft.. For example, at 4 mph a sprayer might deliver 1/2 gallon per 1,000 sq.ft. at 40 psi.
2. To apply evenly, drive at a constant speed that delivers the prescribed amount of spray required for the area to be treated.
3. Mix the appropriate amount of product and water in the sprayer using the following tables as a guide.

Table 5: 12 Volt & Pull Behind Sprayers: Application Guide For Bahiagrass, Bermudagrass, Fescues, Kentucky Bluegrass, Perennial Ryegrass, and Zoysiagrass

Tank Size	Spray volume	Coverage Area	Concentrate to add to tank
15 gal	0.5 gal/1000 sq.ft.	30,000 sq.ft.	120 fl.oz.
	1 gal/1000 sq.ft.	15,000 sq.ft.	60 fl.oz.
25 gal	0.5 gal/1000 sq.ft.	50,000 sq.ft.	200 fl.oz.
	1 gal/1000 sq.ft.	25,000 sq.ft.	100 fl.oz.

Table 6: 12 Volt & Pull Behind Sprayers: Application Guide For Centipedegrass and St. Augustinegrass

Tank Size	Spray volume	Coverage Area	Concentrate to add to tank
15 gal	0.5 gal/1000 sq.ft.	30,000 sq.ft.	60 fl.oz.
	1 gal/1000 sq.ft.	15,000 sq.ft.	30 fl.oz.
25 gal	0.5 gal/1000 sq.ft.	50,000 sq.ft.	100 fl.oz.
	1 gal/1000 sq.ft.	25,000 sq.ft.	50 fl.oz.

For spot treatments with 12 Volt & Pull Behind Sprayers

See rates under How Much To Apply.

3.3 Dial Style Hose-End Sprayers

Ideal for treating medium-sized lawns

1. Fill sprayer with sufficient concentrate to cover the total area to be treated. DO NOT ADD WATER.
2. Close sprayer and set dial to 1 oz.
3. Connect sprayer to hose
4. Extend hose to furthest point from faucet. Walking back towards the faucet, begin spraying, avoiding contact with the treated area.
5. Pour any unused product back into this container.

3.4 Concentrate Plus Water Hose-End Sprayers

Ideal for treating medium-sized lawns

1. Fill sprayer with sufficient concentrate to cover the total area to be treated. Pour product into sprayer jar to the “Fluid Oz.” level desired (See sprayer jar for markings)
2. Add the appropriate amount of water to the “Gallons” level indicated on the side of the sprayer jar. Close sprayer, connect to garden hose, shake well.
3. Extend hose to furthest point from faucet. Walking back towards the faucet, begin spraying, avoiding contact with the treated area.

4. Use Precautions

- Apply when weeds are young and actively growing.
- Apply 1 to 3 days after mowing.
- If the lawn is dry, water 24 hours before using this product.
- Do not water lawn within 24 hours after treatment.
- Certain hard-to-kill weeds may need to be spot treated in 30 days.
- For new lawns, apply only when grass has reached a height of at least 2 inches.
- Bare spots may be seeded 3 to 4 weeks after application.
- Spray when air temperatures are between 60° and 90°F with good soil moisture.
- Avoid applications during the spring transition of the warm season grasses and during periods of extremely high temperatures.

5. Limitations

Broadcast applications to residential turfgrass:

The maximum application rate is 4.0 fl.oz. of product per 1,000 sq.ft. per application. The maximum number of broadcast applications is limited to 2 per year with a minimum of 30 days between applications. The maximum seasonal rate is 8.0 fl.oz. of product per 1,000 sq.ft.

Spot treatments to residential turfgrass:

Spot treatment is defined as a treatment area no greater than 1,000 sq.ft. per acre. The maximum application rate is 4.0 fl.oz. per 1,000 sq.ft. per application. The maximum number of spot treatments is limited to 2 per year with a minimum of 30 days between applications.

6. Weeds Killed

Amaranth, purple	Carpetweed	Dove weed	Matchweed
Amaranth, slender	Catnip	Elderberry	Mexicanweed
Aster, heath	Catsear, spotted (false dandelion)	Falseflax	Morningglory, bigroot
Aster, white heath	Chickweed, common	False sunflower	Morningglory, ivyleaf
Aster, purple	Chickweed, mouseear	Fiddleneck	Morningglory, mexican
Aster, white prairie	Chickweed, sticky	Filaree, redstem	Morningglory, red
Austrian fieldcress	Chicory	Filaree, whitestem	Morningglory, tall
Bedstraw, Catchweed	Cinquefoil	Fleabane, daisy (annual)	Morningglory, wolly
Bedstraw, smooth	Clover, crimson	Fleabane, rough	Mugwort
Bedstraw, Northern	Clover, alyce	Galinsoga, hairy	Mustard, ball
Bedstraw, pineywoods	Clover, hop	Galinsoga, smallflower	Mustard, black
Beggarticks, hairy	Clover, rabbit foot	Goldenrod	Mustard, blue
Betony, Florida	Clover, red	Ground ivy	Mustard, hedge
Bindweed, Field	Clover, strawberry	Gumweed	Mustard, Indian,
Bindweed, Hedge	Clover, yellow sweet	Hairy fleabane	Mustard, tansy
Bitter Wintercress	Clover, white sweet	Hawkweed, meadow	Mustard, tumble
Bittercress, hairy	Clover, white	Hawkweed, mouseear	Mustard, white
Bitterweed	Cockle	Hawkweed, orange	Mustard, wormseed
Black-eyed Susan	Cocklebur	Hawkweed, yellow	Nettle, stinging
Black Medic	Cocklebur, oriental	Healall	(burning)
Blood flower milkweed	Cocklebur, spiny	Heartleaf drymary	Nettle, tall
Blue lettuce	Common mullein	Hemp	Nettle, wood
Brassbuttons	Creeping Jenny (Creeping Charlie)	Henbit	Parsley-piert
Bristly oxtongue	Cudweed	Hoary cress	Parsnip
Broomweed	Daisy, English	Horsenettle	Pearlwort
Buckhorn	Daisy, oxeye	Jimsonweed	Pennycress
Bullnettle	Dandelion, common	Knawel	(frenchweed)
Burclover	Dandelion, false	Knotweed, prostrate	Pepperweed,
Burdock	Deadnettle	Kochia	perennial
Burweed, lawn	Dichondra	Lambsquarters	Pepperweed, virginia
(spurweed)	Dock, broadleaf	Lespedeza	(Peppergrass)
Buttercup, bulbous	Dock, curly	Mallow, alkali	Pigweed, prostrate
Buttercup, creeping	Dogbane	Mallow, bristly	Pigweed, redroot
Buttonweed, common	Dogfennel	Mallow, common	Pigweed, russian
(poorjoe)	Dollarweed	Mallow, dwarf	Pigweed, smooth
Buttonweed, virginia	(Pennywort)	Mallow, venice	Pigweed, tumble
Carolina geranium (wild)			(cont. on next page)

6. Weeds Killed (cont.)

Pineapple weed	Ragweed, western	Tanweed	Western salsify
Plains coreopsis (tickseed)	Redstem filaree	Thistle, blessed	Wild aster
Plantain, blackseed	Rough cinquefoil	Thistle, bull	Wild buckweed
Plantain, bracted	Scarlet pimpernel	Thistle, canada	Wild carrot
Plantain, broadleaf	Shepherd's purse	Thistle, flodman (prairie)	Wild four-o'clock
Plantain, buckhorn	Smooth chaff-flower	Thistle, musk	Wild garlic
Plantain, hoary	Smoothdock	Thistle, platte	Wild lettuce
Plantain, narrowleaf	Sorrel, red (sheep)	Thistle, plumeless	Wild marigold
Plantain, slender	Sowthistle, annual	Thistle, russian	Wild mustard (yellow rocket)
Plantain, woolly	Sowthistle, spiny	Thistle, scotch	Wild onion
Poison ivy	Spanish needles	Thistle, wavyleaf	Wild parsnip
Poison oak	Speedwell, bi-lobed	Thistle, yellow (horrible)	Wild radish
Pokeweed	Speedwell, common	Trailing crownvetch	Wild rape
Prairie Sunflower	Speedwell, corn	Velvetleaf	Wild strawberry
Prickly lettuce (compass plant)	Speedwell, ivyleaf	Vervain, blue	Wild sweet potato
Prickly sida	Speedwell, persian	Vervain, hoary	Wild violet
Prostrate knotweed	Speedwell, purslane	Vervain, prostrate	Woodsorrel, creeping
Puncturevine	Speedwell, snow	Vervain, tall	Woodsorrel, violet
Purslane, common	Speedwell, thymeleaf	Vetch, bird	Woodsorrel, yellow (Oxalis)
Pusley, brazil	Spiny amaranth	Vetch, hairy	Woolly croton
Pusley, florida	Spurge, spotted	Vetch, milk	Wormseed
Pusley, large flower	Spurge, prostrate	Vetch, narrowleaf	Yarrow, common
Ragweed, bur	Spurweed	Vetch, wild	Yarrow, western
Ragweed, common	Strawberry, India mock	Virginia creeper	Yarrow, yellow
Ragweed, lancleaf	Tansy ragwort	Western clematis	

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage and disposal.

PESTICIDE STORAGE: Keep from freezing. Store in original container in a locked storage area inaccessible to children and pets.

PESTICIDE DISPOSAL AND CONTAINER HANDLING: Nonrefillable container. Do not reuse or refill this container. **If empty:** Place in trash or offer for recycling, if available. **If partly filled:** Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

LIMITED WARRANTY AND DISCLAIMER

FOR USE ONLY AS DIRECTED. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE MANUFACTURER NEITHER MAKES NOR INTENDS ANY EXPRESS OR IMPLIED WARRANTIES, INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, WHICH ARE HEREBY EXPRESSLY DISCLAIMED. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO CASE SHALL THE MANUFACTURER BE LIABLE FOR INCIDENTAL, CONSEQUENTIAL, OR SPECIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. If these terms are not acceptable, return this product unopened immediately to the point of purchase, and the purchase price will be refunded in full. The terms of this LIMITED WARRANTY AND DISCLAIMER cannot be varied by any written or verbal statements or agreements at the point of sale or elsewhere.

TRIMEC® is a registered trademark of PBI-Gordon Corporation.

809/2-2015 AP061214
EPA REG. NO. 2217-539-33955
EPA EST. NO. 2217-KS-1

MANUFACTURED AND DISTRIBUTED BY
PBI/GORDON CORPORATION
1217 WEST 12TH STREET
KANSAS CITY, MISSOURI 64101
www.GordonsUSA.com

THIS PAGE IS BLANK INTENTIONALLY.



**READ THE ENTIRE LABEL FIRST.
OBSERVE ALL PRECAUTIONS AND
FOLLOW DIRECTIONS CAREFULLY.**

PEEL HERE TO OPEN

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION: Harmful if absorbed through skin. Harmful if swallowed. Avoid contact with skin, eyes, or clothing. Causes moderate eye irritation. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet. Wear long-sleeved shirt, long pants, socks, shoes and gloves. Remove and wash contaminated clothing before reuse.

First Aid	
If in eyes:	<ul style="list-style-type: none">• Hold eye open and rinse slowly and gently with water for 15 - 20 minutes.• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.• Call a poison control center or doctor for treatment advice.
If swallowed:	<ul style="list-style-type: none">• Call a poison control center or doctor immediately for treatment advice.• Have person sip a glass of water if able to swallow.• Do not induce vomiting unless told to do so by the poison control center or doctor.• Do not give anything by mouth to an unconscious person.
If on skin or on clothing:	<ul style="list-style-type: none">• Take off contaminated clothing.• Rinse skin immediately with plenty of water for 15 - 20 minutes.• Call a poison control center or doctor for treatment advice.
If inhaled:	<ul style="list-style-type: none">• Move person to fresh air.• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible.• Call a poison control center or doctor for treatment advice.
HOT LINE NUMBER: Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact 1-877-800-5556 for emergency medical treatment information.	

809/2-2015 AP061214
EPA REG. NO. 2217-539-33955
EPA EST. NO. 2217-KS-1

**MANUFACTURED AND DISTRIBUTED BY
PBI/GORDON CORPORATION**
1217 WEST 12TH STREET
KANSAS CITY, MISSOURI 64101
www.GordonsUSA.com