

2006-2008

9561.135

egrs

GREEN LIGHT[®] LIGHT

**CONTROLS
TAKE-ALL PATCH
& BROWN PATCH**

Fung-Away[®] Systemic Lawn Fungicide



**A Systemic Protectant and Curative Fungicide
for the Control of Diseases in Established Lawns**



LICENSED

PERIOD 2015-2017 LIC. NO.

**9561.135
Disc.**

TREATS UP TO 3,333 SQ. FT.

**KEEP OUT OF REACH OF CHILDREN
CAUTION** See Back Panel For Additional
Precautionary Statements.

EPA Reg. No. 62719-461-869

NET WT. 8 LBS. (3.6kg)

ACTIVE INGREDIENT:
myclobutanil: a-butyl-a-(4-chlorophenyl)-
1-H-1,2,4 triazole-propanenitrile 0.39%
OTHER INGREDIENTS: 99.61%
TOTAL 100.00%

EPA Est. No. 73273-TX-1

FUNG-AWAY[®]



LICENSED

PERIOD 2012-2014 LIC. NO.



LICENSED

PERIOD 2009-2011 LIC. NO.

GREEN LIGHT[®] LIGHT

Fung-Away[®] Systemic Lawn Fungicide

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

Causes Moderate Eye Irritation • Harmful if Absorbed Through The Skin • Avoid Contact with Skin, Eyes or Clothing.
PERSONAL PROTECTIVE EQUIPMENT (PPE): User Must Wear: Long-sleeved shirt and long pants, chemical-resistant gloves made of any waterproof material, shoes plus socks.
Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS: Users should wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
FIRST AID: IF SWALLOWED: IF IN EYES: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.
IF ON SKIN OR CLOTHING: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

HOT LINE NUMBER: Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-992-5994 day or night, for emergency treatment information.

ENVIRONMENTAL HAZARDS: This pesticide is toxic to fish. Drift and runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment wash-waters or disposing of wastes.

NOTICE: Read the entire label. Use only according to label directions. Before using this product, read Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies at end of Directions for Use. If terms are unacceptable, return at once unopened.

In case of emergency endangering health or the environment involving this product, call 1-800-992-5994.
Agricultural Chemical: Do not ship or store with food, feeds, drugs or clothing.

DIRECTIONS FOR USE: IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

NON-AGRICULTURAL USE REQUIREMENTS: The requirements in this box apply to uses of this product that are not within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries or greenhouses. Seed treatments and professional applications to lawn grasses, golf courses, industrial (office park), municipal and residential lawns are not within the scope of the Worker Protection Standard. KEEP UNPROTECTED PERSONS OUT OF TREATED AREA UNTIL DUSTS HAVE SETTLED.

STORAGE AND DISPOSAL: Do not contaminate water, food or feed by storage and disposal.
Storage: Keep away from fire and sparks. Store in a cool, dry, well-ventilated area. In case of spill, sweep up material and dispose of material according to "Pesticide Disposal" instructions.

Pesticide Disposal: Wastes resulting from the use of this product may be disposed of on site according to label directions or at an approved waste disposal facility.

Container Disposal: Do not reuse empty containers. Completely empty bag into application equipment. Dispose of emptied bag(s) in a sanitary landfill or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

GENERAL INFORMATION: GREEN LIGHT FUNG-AWAY SYSTEMIC LAWN FUNGICIDE is a systemic, protectant and curative fungicide recommended for the control of listed diseases in established lawns. Optimum disease control is achieved when this product is applied to established turfgrasses in a regularly scheduled preventative program.

USE DIRECTIONS FOR TURFGRASSES

• GREEN LIGHT FUNG-AWAY SYSTEMIC LAWN FUNGICIDE should be used in conjunction with turf management practices that promote good plant health and optimum disease control. The key to selecting a fungicide is the proper diagnosis of the organism causing the disease. Diagnostic kits, extension experts, or other identification methods should be used when developing disease control strategies.

• Optimum disease control is achieved when GREEN LIGHT FUNG-AWAY SYSTEMIC LAWN FUNGICIDE is applied in a preventative disease control program at a rate of 4 to 8 lbs. per 1000 square feet. See the following table for specific application rates for various diseases.

• For the control of root and crown rot, this product must be watered in using a minimum of 3 to 4 gallons of water per 1000 square feet.

• Under conditions that are favorable for disease development, the interval between applications of GREEN LIGHT FUNG-AWAY SYSTEMIC LAWN FUNGICIDE should be reduced.

• Unless otherwise specified, when disease pressure is high or when used as a curative, use higher rates of GREEN LIGHT FUNG-AWAY SYSTEMIC LAWN FUNGICIDE and shorter application interval. Under light to moderate disease pressure, apply GREEN LIGHT FUNG-AWAY SYSTEMIC LAWN FUNGICIDE at the low use rate and/or longer application intervals.

• Apply GREEN LIGHT FUNG-AWAY SYSTEMIC LAWN FUNGICIDE uniformly using an appropriate fertilizer spreader. Calibrate the spreader before application as calibration may change with spreader age and use, walking speed and evenness of terrain. To calibrate the spreader, weigh out the amount of product necessary to treat an area of 800 square feet. Using the suggested spreader setting provided by the manufacturer or from the table below, apply to a pre-measured area of 500 square feet (5 x 100 square feet) area. Weigh back product remaining in application equipment and adjust the spreader as necessary to give the correct rate of delivery for the measured area.

• To avoid pick-up, lightly irrigate treated areas soon after application.

• On short cut bentgrass, (1/2" or less) when temperatures are above 80°F, apply only to dry foliage.

Apply GREEN LIGHT FUNG-AWAY SYSTEMIC LAWN FUNGICIDE with drop or rotary spreaders designed to apply granules. Avoid the use of spreaders that would apply this product in narrow rows or concentrated bands. Before each application, calibrate the spreader according to the equipment manufacturer's directions such that the spreader delivers the recommended application rate. Apply this product uniformly over the lawn or ornamental turf area. A more uniform application can be made by spreading half of the required amount over the area in a given direction and then applying the remaining half at a right angle to the previous direction. Avoid streaking, skips or overlaps during application. Check equipment frequently to ensure equipment is functioning properly and applying uniform distribution of granules.

Disease	FUNG-AWAY (lb/1000 sq. ft.)	Application Interval/Timing	Instructions	Restrictions
Anthraxnose	4	14-21 days	Apply when conditions are favorable for disease development.	Do not apply more than 46 pounds product per 1,000 sq. ft. per year.
Red Thread				
Septoria Leaf Spot				For Nassau and Suffolk Counties in New York State, use of this product is limited to 11.5 pounds per 1000 sq. ft. per year (1.95 lb. active ingredient per acre).
Brown Patch	4 - 8	14 days	Begin applications when conditions are favorable for disease development and before disease symptoms are apparent.	
Copper Spot	4	14 days	Apply when conditions are favorable for disease development.	For professional use only in the State of New York.
Zonate Leaf Spot				
Dollar Spot	3 - 8	14 - 28 days	Apply when conditions are favorable for disease development. Make no more than 3 consecutive applications for dollar spot control before rotating to a registered fungicide with a different mode of action.	
Fusarium Blight	4 - 8	14 - 21 days	Apply when conditions are favorable for disease development.	
Fusarium Patch (Pink Snow mold)	4 - 8	Fall-Winter	Apply prior to snow cover.	
Leaf Spot	4	14 days	Apply when conditions are favorable for disease development. For crown rot, water in with 3 to 4 gallons of water per 1000 square feet to increase penetration to crown and roots.	
Melting-Out				
Crown Rot				
Leaf Smuts	4	14 days	Apply in the fall after grass enters dormancy and/or in the spring prior to the initiation of growth.	
Necrotic Ring Spot	4 - 8	Spring: 28 days	Make applications on a preventative basis in early to mid-spring.	
Powdery Mildew	4	14 - 28 days	Apply when conditions are favorable for disease development.	
Rusts				

Disease	FUNG-AWAY (lb/1000 sq. ft.)	Application Interval/Timing	Instructions	Restrictions
Necrotic Ring Spot	4 - 8	Fall: 28 days	Make 2 applications beginning in August before the turf goes dormant. Apply 8 lbs./1000 sq. ft. followed by a second application one month later.	
Summer Patch	4 - 8	14 - 28 days	Begin applications in the spring when conditions are favorable for disease development. Make 2 to 4 applications depending on recommendations from local turfgrass extension experts. Water in with at least 3 to 4 gallons of water per 1000 square feet to increase spray penetration to crown and roots.	
Take-All Patch	8	Spring/Fall: 28 days	To reduce the severity of take-all patch, make 1 to 2 fall applications in September and October or when night temperatures drop to 55°F and 1 to 2 spring applications in April and May depending on local recommendations.	
Zoysia Large Patch	8	Fall: 28 days	Make applications in fall before turf becomes dormant. Water in with at least 3 to 4 gallons of water per 1000 square feet to increase penetration to crown and roots.	

APPROXIMATE SPREADER SETTINGS

MAKE	MODEL	LBS. PER 1,000 SQ. FT.			SPREAD WIDTH
Broadcast Spreaders		2	4	8	
CYCLONE	B1/B55	4 3/4	5 1/2	7 1/2	5 ft.
ORTHO	600	3	6	12	5 ft.
SCOTTS	ER-2	27	29	33 1/2	3 1/2 ft.
SEARS 1100	671.1987	3	4 1/2	6	5 ft.
SCOTTS	Speedy Green	5	8	14	6 ft.
Drop Spreaders					
ORTHO	5000	7 1/2	9 1/2	12	18 in.
SCOTTS	PF-3	7	9	12	21 in.

TERMS AND CONDITIONS OF USE

If terms of the following Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies are not acceptable, return unopened package at once to the seller for a full refund of purchase price paid. Otherwise, use by the buyer or any other user constitutes acceptance of the terms under Warranty Disclaimer, Inherent Risks of Use and Limitations of Remedies.

WARRANTY DISCLAIMER

Seller warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions, subject to the inherent risks set forth below. Seller MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.

INHERENT RISKS OF USE

It is impossible to eliminate all risks associated with use of this product. Plant injury, lack of performance, or other unintended consequences may result because of such factors as use of the product contrary to label instructions (including conditions noted on the label, such as unfavorable temperature, soil conditions, etc.), abnormal conditions (such as excessive rainfall, drought, tornadoes, hurricanes), presence of other materials, the manner of application, or other factors, all of which are beyond the control of the Seller. All such risks shall be assumed by buyer.

Limitation of Remedies: The exclusive remedy for losses or damages resulting from this product (including claims based on contract, negligence, strict liability, or other legal theories), shall be limited to, at Seller's election, one of the following:

1. Refund of purchase price paid by buyer or user for product bought, or
2. Replacement of amount of product used

Seller shall not be liable for losses or damages resulting from handling or use of this product unless Seller is promptly notified of such loss or damage in writing. In no case shall Seller be liable for consequential or incidental damages or losses.

The terms of the Warranty Disclaimer and Inherent Risks of Use above and this Limitation of Remedies cannot be varied by any written or verbal statements or agreements. No employee or sales agent of Seller or the seller is authorized to vary or exceed the terms of the Warranty Disclaimer or this Limitation of Remedies in any manner.

MANUFACTURED FOR GREEN LIGHT COMPANY
P. O. BOX 17985, SAN ANTONIO, TX 78217-0985

www.greenlightco.com Tel. (210) 494-3481

Fung-Away[®] is a registered trademark of Green Light Company



100%
EMPLOYEE
OWNED

Rev.
7-17-2002

#49424-ADN
FOR POSITION ONLY

Fung-Away[®]

Fung-Away[®]

Fung-Away[®]