

# 12455-117-3240

Printed at 100% on 11x17



LICENSED

PERIOD 2012-2014 LIC. NO.



LICENSED

PERIOD 2015-2017 LIC. NO.

9566.49

**Agriid<sub>3</sub>®**  
**PELLETED BAIT**

**KILLS RATS & MICE**

- Can Be Used Indoors or Out
- Can Be Used for Burrow Baiting
- Palatable Formula
- Kills Anticoagulant Resistant Norway Rats

**MOTOMCO**

**KEEP OUT OF REACH OF CHILDREN CAUTION**

See back panels for First Aid and additional precautionary statements.

ACTIVE INGREDIENT: Cholecalciferol (CAS #67-97-0): . . . 0.075%  
OTHER INGREDIENTS: . . . . . 99.925%  
TOTAL 100.000%

**Net Weight: 10 lbs (4.5 kg)**

OMRI Listed

This product may only be used inside and within 100 feet of agricultural buildings and man-made agricultural structures and for all burrow baiting.

Photo is not representative of a label placement.

FIRST AID	
Have label with you when obtaining treatment advice	
If swallowed	<ul style="list-style-type: none"> <li>Call a poison control center, doctor, or 1-877-854-2494 immediately for treatment advice.</li> <li>Have person sip a glass of water if able to swallow.</li> <li>Do not induce vomiting unless told to do so by the poison control center or doctor.</li> </ul>

**NOTE TO PHYSICIAN**  
If serum calcium levels are elevated, treatment with Calcitonin is effective in reducing calcium to normal levels. Continue monitoring serum calcium and treat as necessary for hypercalcaemia (Reference, AMA Drug Evaluations, Third Edition (1977), Chapter 16, pp. 248-251).

**TREATMENT FOR PET POISONING**  
If animal eats bait, call veterinarian or 1-877-854-2494 at once.

**NOTE TO VETERINARIAN**  
**Calcium Mobilizer Cholecalciferol:** For animals ingesting bait, induce vomiting by using hydrogen peroxide or administration of activated charcoal with a cathartic. If clinical signs develop, treatment consisting of saline diuresis combined with the use of furosemide, corticosteroids, and phosphate binders are recommended. Calcitonin or pamidronate may be needed for animals that remain hypercalcaemic despite symptomatic treatment.

**DIRECTIONS FOR USE**  
It is a violation of Federal law to use this product in a manner inconsistent with its labeling.  
**READ THIS LABEL:** Read this entire label and follow all use directions and use precautions. Use only for the sites, pests, and application methods described on this label.

- IMPORTANT:** Do not expose children, pets, or nontarget animals to rodenticides. To help to prevent exposure:
- Store unused product out of reach of children and pets.
  - Apply bait in locations out of reach of children, pets, domestic animals and nontarget wildlife, or in tamper-resistant bait stations. These stations must be resistant to destruction by dogs and by children under six years of age, and must be used in a manner that prevents such children from reaching into bait compartments and obtaining bait. If bait can be shaken from bait stations when they are lifted, units must be secured or otherwise immobilized. Stronger bait stations are needed in areas open to hoofed livestock, raccoons, bears, or other potentially destructive animals, or in areas prone to vandalism.
  - Dispose of product container and unused, spoiled, or unconsumed bait as specified on this label.

**Note:** Bait stations are mandatory for outdoor, above-ground use. Tamper-resistant bait stations must be used if children, pets, non-target mammals, or birds may access the bait.

**USE RESTRICTIONS:** This product can only be used to control Norway rats, roof rats and house mice in and within 100 feet of agricultural buildings and man-made agricultural structures constructed in a manner so as to be vulnerable to commensal rodent invasions and/or to harboring or attracting rodent infestations. Fence and perimeter baiting, beyond 100 feet from an agricultural structure as defined above, is prohibited. This product must not be applied directly to food or feed crops.

This product may be applied to active rodent burrows to control Norway rats, and roof rats within or beyond 100 feet of agricultural buildings and man-made agricultural structures, provided that infestations of these rodents have been confirmed. Bait must be placed no less than 6 inches into active

6080B-2

**Agriid<sub>3</sub>®**  
**PELLETED BAIT**  
**FOR ORGANIC PRODUCTION**  
**KILLS RATS & MICE**

**KILLS NORWAY RATS, ROOF RATS AND HOUSE MICE**

ACTIVE INGREDIENT: Cholecalciferol (CAS #67-97-0): . . . 0.075%  
OTHER INGREDIENTS: . . . . . 99.925%  
TOTAL 100.000%

**KEEP OUT OF REACH OF CHILDREN CAUTION**

See side panels for First Aid and additional precautionary statements.

**STORAGE AND DISPOSAL**

Do not contaminate water, food or feed by storage or disposal.

**Pesticide Storage:** Store only in original container in a cool, dry place inaccessible to children and pets. Keep containers closed and away from other chemicals.

**Pesticide Disposal:** Wastes resulting from the use of this product may be placed in trash or delivered to an approved waste disposal facility.

**Container Handling:** Nonrefillable container. Do not reuse or refill this container.

**[Plastic:]** Offer for recycling or reconditioning; or puncture and dispose of in a sanitary landfill; or by incineration.

**WARRANTY:** To the extent consistent with applicable law, seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.

**Net Weight: 10 lbs (4.5 kg)**

EPA REG. NO. 12455-117-3240

EPA EST. NO. 12455-WI-1

Available Exclusively From:



**The World Leader In Rodent Control Technology®**

3699 Kinsman Blvd.  
Madison, WI 53704 U.S.A.  
www.motomco.com

**DIRECTIONS FOR USE** (Continued from other panel)

**Norway/roof rat burrows.** Do not broadcast bait. Because Norway/roof rat infestations may occur in areas farther than 100 feet from buildings and man-made structures when the rodents have ample supplies of food and cover, efforts should be made to remove food trash, garbage, clutter and debris.

Do not place near or inside ventilation duct openings. Do not contaminate water, food, feedstuffs, food or feed handling equipment, or milk or meat handling equipment or surfaces that come into direct contact with food. When used in USDA inspected facilities, this product must be applied in tamper-resistant bait stations.

Do not sell this product in individual containers holding less than 4 pounds of bait.

**SELECTION OF TREATMENT AREAS:** Determine areas where rats or mice will most likely find and consume the bait. Generally, these areas are along walls, by gnawed openings, in or beside burrows, in corners and concealed places, between floors and walls, or in locations where signs of rats or house mice have been seen. Protect bait from rain and snow. Remove as much alternative food as possible.

**APPLICATION DIRECTIONS:**  
**Norway Rats and Roof Rats:** Use 2 to 8 oz. of bait per placement. Space placements at intervals of 15 to 30 feet in infested areas. Maintain an uninterrupted supply of fresh bait for 10 days or until there no longer are signs of recent feeding on bait by rats.

**House Mice:** Use 1/4 to 1/2 oz. (1 to 2 level tablespoons) of bait per placement. Space placements at intervals of 8 to 12 feet in infested areas. Up to 1 oz. of bait may be used at points of very high mouse activity. Maintain an uninterrupted supply of fresh bait for 15 days or until there no longer are signs of recent feeding on bait by house mice.

**FOLLOW-UP:** Replace contaminated or spoiled bait immediately. Wearing gloves, collect and dispose of all dead, exposed animals and leftover bait. To prevent reinfestation, limit sources of rodent food, water, and harborage as much as possible. If reinfestation does occur, repeat treatment. Where a continuous source of infestation is present, establish permanent bait stations and replenish as needed.

**PRECAUTIONARY STATEMENTS**

**HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

**CAUTION:** Harmful if swallowed. Keep away from children, domestic animals and pets. All handlers (including applicators) must wear shoes, socks, and gloves. Any person who retrieves carcasses or unused bait following application of this product must wear gloves.

**User Safety Requirements**

Follow manufacturer's instruction for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry. Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash hands thoroughly after applying bait and before eating, drinking, chewing gum, using the toilet and change into clean clothing.

**ENVIRONMENTAL HAZARDS**

This product is toxic to fish, birds and other wildlife. Dogs and other predatory and scavenging mammals and birds might be poisoned if they feed upon animals that have eaten the bait. Do not apply this product directly to water or to areas where surface water is present or to intertidal areas below the mean high water mark.

050912/04-12

