



LICENSED

PERIOD 2012-2014 LIC. NO.



LICENSED

PERIOD 2015-2017 LIC. NO. 9591.72

Special Instructions for Cleaning and Decontamination Against HIV-1, HBV and HCV of Surfaces or Objects Soiled With Blood or Body Fluids:

Personal Protection: When using this disinfecting towelette, wear appropriate barrier protection such as disposable protective gloves, protective gown, face mask and eye covering.

Cleaning Procedure: Blood and other body fluids must be thoroughly cleaned from surfaces and objects before application of this disinfecting towelette. A **MetriWipes** towelette can be used for this purpose.

Contact Time: Thoroughly wet surface with a new **MetriWipes** towelette. Repeated use of the product may be required to ensure that the surface remains visibly wet for 2 minutes. This contact time will not be sufficient for some of the organisms listed on the label. Refer to label for contact times against other organisms.

Disposal Of Infectious Materials: Cleaning materials used that may contain blood or other body fluids must be autoclaved and disposed of in accordance with Federal, State and local regulations for infectious materials disposal.

This product is not to be used as a terminal sterilant/high-level disinfectant on any surface or instrument that (1) is introduced directly into the human body, either into or in contact with the bloodstream or normally sterile areas of the body, or (2) contacts intact mucous membranes but which does not ordinarily penetrate the blood barrier or otherwise enter normally sterile areas of the body. This product may be used to pre-clean or decontaminate critical or semi-critical medical devices prior to sterilization/high-level disinfection.

PRECAUTIONARY STATEMENTS
Hazards to Humans and Domestic Animals
CAUTION

Harmful if absorbed through the skin. Causes moderate eye irritation. Avoid contact with eyes, skin, or clothing. Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet. Remove contaminated clothing and wash clothing before reuse.

FIRST AID

If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing. Call a poison control center or doctor for further advice.

If on skin: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

For Chemical Emergency ONLY (spill, leak, fire, exposure or accident), call CHEMTREC at 1-800-424-9300. For ALL other inquiries about this product, call Metrex Customer Service at 1-800-841-1428, Monday through Friday, between 6 a.m. and 4 p.m. Pacific Time.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment.

STORAGE AND DISPOSAL

Towelette - Do not reuse towelette. Dispose of used towelette in trash.

Dispenser - Nonrefillable container. Do not reuse or refill this container. Offer for recycling, if available.

For product or technical information, contact Metrex at **800-841-1428** or visit our website at **www.metrex.com**.

Packed For:
METREX® RESEARCH
28210 Wick Road
Romulus, MI 48174

MADE IN USA

REORDER No. XX-XXX

70-xxxx-1
xxxx-xx

MetriWipes®

Disinfecting Towelettes

- TUBERCULOCIDAL
- BACTERICIDAL
- VIRUCIDAL*
- FUNGICIDAL



Net Contents: X lb. X oz.
120 Pre-Saturated Towelettes (6" x 9")

MetriWipes Disinfecting Towelettes are non-woven disposable towelettes. **MetriWipes** are intended for use in health care settings such as hospitals, dental offices, laboratories, clinics, ophthalmic offices, veterinary facilities and in kitchens** and bathrooms. **MetriWipes** are effective against the following microorganisms on hard, non-porous surfaces when used as directed:

- *Mycobacterium tuberculosis* var: *bovis* (BCG)
- *Staphylococcus aureus*
- *Pseudomonas aeruginosa*
- *Salmonella enterica*
- *Trichophyton mentagrophytes*
- Methicillin Resistant *Staphylococcus aureus* (MRSA)
- Vancomycin Resistant *Enterococcus faecalis* (VRE)
- *Staphylococcus aureus* with Reduced Susceptibility to Vancomycin
- Hepatitis B Virus (HBV)*
- Hepatitis C Virus (HCV)*
- Herpes Simplex Virus Types 1 and 2*
- Human Immunodeficiency Virus (HIV-1)*
- Influenza A2 Virus*

ACTIVE INGREDIENTS:

Diisobutylphenoxyethoxyethyl dimethyl benzyl ammonium chloride	0.28%
Isopropanol.....	17.20%
INERT INGREDIENTS†:	82.52%
TOTAL	100.00%

† Does not include the weight of the dry towelette.



KEEP OUT OF REACH OF CHILDREN CAUTION

See other panel for additional precautionary statements.

EPA Reg. No. 46781-9
EPA Est. No. 56952-WI-001

DO NOT USE AS A BABY WIPE. Not For Use On Skin.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

MetriWipes Disinfecting Towelettes are non-woven disposable towelettes. **MetriWipes** are intended for use on hard, non-porous surfaces in health care settings such as hospitals, emergency medical settings, neonatal units, surgical centers, isolation areas, out-patient surgical centers, nursing homes, dental offices, ophthalmic offices, eye surgical centers, patient care areas, dialysis centers, donation centers (blood, plasma, semen, milk, aphaeresis), veterinary facilities, salon settings, emergency vehicles, correctional facilities, schools, health club facilities, tanning salons, kitchens**, non-food contact surfaces in food preparation areas**, bathrooms, day care centers, police and fire vehicles, cadaver processing areas, laundry rooms, and other critical care areas where control of cross contamination between treated surfaces is required.

MetriWipes will clean and disinfect, when used as directed, such items as interior and exterior surfaces of infant incubators, bassinets, cribs and warmers, infant/child care equipment surfaces, diaper changing stations, oxygen hoods, operating room tables and lights, laboratory equipment and surfaces, physical therapy equipment surfaces, ambulance equipment surfaces, slit lamps, hard, non-porous exterior surfaces of ultrasound transducers or probes, exam tables, gurneys, IV poles, dental unit instrument trays, operatory light switches, exterior surfaces of amalgamators and dental curing lights, exterior surfaces of anesthesia machines and respiratory therapy equipment, SilkPeel™ with DermalInfusion™, workstations, dental countertops, loupes, exterior surfaces of endodontic equipment, such as apex locators, pulp testers and motors, stools, nail/hair care implements, tanning beds, salon surfaces, non-food contact surfaces**, exterior surfaces of refrigerator units**, exterior surfaces of appliances such as refrigerators, microwaves, food processors and stoves**, vanity tops**, floors, walls, handrails, door knobs, bed railings, tables, chairs, bathroom fixtures, exterior toilet surfaces, garbage cans, keyboards, carts, shopping carts, weight-lifting surfaces, drained whirlpool tank surfaces, cabinets, lights, telephones, which are made of plastic (i.e., polycarbonate, polypropylene, polyvinylchloride, polystyrene), and use sites listed on this label consisting of non-porous vinyl, stainless steel, painted surfaces, Plexiglas®, Formica® and glass.

Kitchen and Bath Wipes for Hard, Non-Porous Surfaces Cleans, Deodorizes and Kills Common Household Germs**

Dispenser Directions:

To start feed, remove cover and discard seal. From center of towelette roll, pull up a towelette corner, twist it into a point and thread it through the hole in the container cover. Pull through about one inch. Replace cover. Pull out first towelette and tear off at a 90° angle. Remaining towelettes feed automatically, ready for next use. When not in use, keep center cap closed to prevent solution loss.

Cleaning Instructions:

Use one **MetriWipes** towelette to completely preclean surface of all gross debris. **For use as a virucide*, including HIV-1, HBV and HCV applications and effectiveness against Methicillin Resistant Staphylococcus aureus (MRSA), Vancomycin Resistant Enterococcus faecalis (VRE) and Staphylococcus aureus with reduced susceptibility to vancomycin:** Use a second **MetriWipes** towelette to thoroughly wet the surface. Repeated use of the product may be required to ensure that the surface remains visibly wet for 2 minutes at room temperature (68 °F/20 °C).

For use as a disinfectant:

Use a second **MetriWipes** towelette to thoroughly wet the surface. Repeated use of the product may be required to ensure that the surface remains visibly wet for 3 minutes at room temperature (68 °F/20 °C) for Tuberculocidal activity and effectiveness against *Staphylococcus aureus*, *Pseudomonas aeruginosa*, *Trichophyton mentagrophytes* and *Salmonella enterica*.

To disinfect external surfaces of ultrasound transducers:

Pre-clean ultrasound gel from external transducer to be disinfected. Then thoroughly wet the probe with a **MetriWipes** towelette. Repeated use of the product may be required to ensure that the surface remains visibly wet for 3 minutes at 20 °C.

**A thorough rinse with potable water is required for surfaces in direct contact with food.

MetriWipes effectively kills Human Immunodeficiency Virus Type 1 (HIV), Hepatitis B Virus (HBV) and Hepatitis C Virus (HCV) on pre-cleaned environmental surfaces/objects previously soiled with blood/body fluids in healthcare settings or other settings in which there is an expected likelihood of soiling of inanimate surfaces/objects with blood/body fluids, and in which the surfaces/objects can be associated with the potential for transmission of HIV, HBV and HCV.



LICENSED

PERIOD 2012-2014 LIC. NO.



LICENSED

PERIOD 2015-2017 LIC. NO.

9591.72

Special Instructions for Cleaning and Decontamination Against HIV-1, HBV and HCV of Surfaces or Objects Soiled With Blood or Body Fluids:

Personal Protection: When using this disinfecting towelette, wear appropriate barrier protection such as disposable protective gloves, protective gown, face mask and eye covering.

Cleaning Procedure: Blood and other body fluids must be thoroughly cleaned from surfaces and objects before application of this disinfecting towelette. A MetriWipes towelette can be used for this purpose.

Contact Time: Thoroughly wet surface with a new MetriWipes towelette. Repeated use of the product may be required to ensure that the surface remains visibly wet for 2 minutes. This contact time will not be sufficient for some of the organisms listed on the label. Refer to label for contact times against other organisms.

Disposal Of Infectious Materials: Cleaning materials used that may contain blood or other body fluids must be autoclaved and disposed of in accordance with Federal, State and local regulations for infectious materials disposal.

This product is not to be used as a terminal sterilant/high-level disinfectant on any surface or instrument that (1) is introduced directly into the human body, either into or in contact with the bloodstream or normally sterile areas of the body, or (2) contacts intact mucous membranes but which does not ordinarily penetrate the blood barrier or otherwise enter normally sterile areas of the body. This product may be used to pre-clean or decontaminate critical or semi-critical medical devices prior to sterilization/high-level disinfection.

PRECAUTIONARY STATEMENTS Hazards to Humans and Domestic Animals CAUTION

Harmful if absorbed through the skin. Causes moderate eye irritation. Avoid contact with eyes, skin, or clothing. Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet. Remove contaminated clothing and wash clothing before reuse.

FIRST AID

If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing. Call a poison control center or doctor for further advice.

If on skin: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

For Chemical Emergency ONLY (spill, leak, fire, exposure or accident), call CHEMTREC at 1-800-424-9300. For ALL other inquiries about this product, call Metrex Customer Service at 1-800-841-1428, Monday through Friday, between 6 a.m. and 4 p.m. Pacific Time.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment.

STORAGE AND DISPOSAL

Towelette - Do not reuse towelette. Dispose of used towelette in trash.

Dispenser - Nonrefillable container. Do not reuse or refill this container. Offer for recycling, if available.

For product or technical information, contact Metrex at 800-841-1428 or visit our website at www.metrex.com.

Packed For: METREX® RESEARCH 28210 Wick Road Romulus, MI 48174

MADE IN USA

REORDER No. XX-XXX

70-xxxx-1
xxxx-xx

MetriWipes®

Disinfecting Towelettes

- TUBERCULOCIDAL
- BACTERICIDAL
- VIRUCIDAL*
- FUNGICIDAL



Net Contents: X lb. X oz.
160 Pre-Saturated Towelettes (6" x 6.75")

MetriWipes Disinfecting Towelettes are non-woven disposable towelettes. MetriWipes are intended for use in health care settings such as hospitals, dental offices, laboratories, clinics, ophthalmic offices, veterinary facilities and in kitchens** and bathrooms. MetriWipes are effective against the following microorganisms on hard, non-porous surfaces when used as directed:

- *Mycobacterium tuberculosis* var: *bovis* (BCG)
- *Staphylococcus aureus*
- *Pseudomonas aeruginosa*
- *Salmonella enterica*
- *Trichophyton mentagrophytes*
- Methicillin Resistant *Staphylococcus aureus* (MRSA)
- Vancomycin Resistant *Enterococcus faecalis* (VRE)
- *Staphylococcus aureus* with Reduced Susceptibility to Vancomycin
- Hepatitis B Virus (HBV)*
- Hepatitis C Virus (HCV)*
- Herpes Simplex Virus Types 1 and 2*
- Human Immunodeficiency Virus (HIV-1)*
- Influenza A2 Virus*

ACTIVE INGREDIENTS:

Diisobutylphenoxyethoxyethyl dimethyl benzyl ammonium chloride 0.28%
Isopropanol 17.20%
INERT INGREDIENTS† 82.52%
TOTAL 100.00%

† Does not include the weight of the dry towelette.



KEEP OUT OF REACH OF CHILDREN CAUTION

See other panel for additional precautionary statements.

EPA Reg. No. 46781-9
EPA Est. No. 56952-WI-001

DO NOT USE AS A BABY WIPE. Not For Use On Skin.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

MetriWipes Disinfecting Towelettes are non-woven disposable towelettes. MetriWipes are intended for use on hard, non-porous surfaces in health care settings such as hospitals, emergency medical settings, neonatal units, surgical centers, isolation areas, out-patient surgical centers, nursing homes, dental offices, ophthalmic offices, eye surgical centers, patient care areas, dialysis centers, donation centers (blood, plasma, semen, milk, aphaeresis), veterinary facilities, salon settings, emergency vehicles, correctional facilities, schools, health club facilities, tanning salons, kitchens**, non-food contact surfaces in food preparation areas**, bathrooms, day care centers, police and fire vehicles, cadaver processing areas, laundry rooms, and other critical care areas where control of cross contamination between treated surfaces is required.

MetriWipes will clean and disinfect, when used as directed, such items as interior and exterior surfaces of infant incubators, bassinets, cribs and warmers, infant/child care equipment surfaces, diaper changing stations, oxygen hoods, operating room tables and lights, laboratory equipment and surfaces, physical therapy equipment surfaces, ambulance equipment surfaces, slit lamps, hard, non-porous exterior surfaces of ultrasound transducers or probes, exam tables, gurneys, IV poles, dental unit instrument trays, operatory light switches, exterior surfaces of amalgamators and dental curing lights, exterior surfaces of anesthesia machines and respiratory therapy equipment, SilkPeel™ with DermalInfusion™, workstations, dental countertops, loupes, exterior surfaces of endodontic equipment, such as apex locators, pulp testers and motors, stools, nail/hair care implements, tanning beds, salon surfaces, non-food contact surfaces**, exterior surfaces of refrigerator units**, exterior surfaces of appliances such as refrigerators, microwaves, food processors and stoves**, vanity tops**, floors, walls, handrails, door knobs, bed railings, tables, chairs, bathroom fixtures, exterior toilet surfaces, garbage cans, keyboards, carts, shopping carts, weight-lifting surfaces, drained whirlpool tank surfaces, cabinets, lights, telephones, which are made of plastic (i.e., polycarbonate, polypropylene, polyvinylchloride, polystyrene), and use sites listed on this label consisting of non-porous vinyl, stainless steel, painted surfaces, Plexiglas®, Formica® and glass.

Kitchen** and Bath Wipes for Hard, Non-Porous Surfaces Cleans, Deodorizes and Kills Common Household Germs

Dispenser Directions:

To start feed, remove cover and discard seal. From center of towelette roll, pull up a towelette corner, twist it into a point and thread it through the hole in the container cover. Pull through about one inch. Replace cover. Pull out first towelette and tear off at a 90° angle. Remaining towelettes feed automatically, ready for next use. When not in use, keep center cap closed to prevent solution loss.

Cleaning Instructions:

Use one MetriWipes towelette to completely preclean surface of all gross debris. For use as a virucide*, including HIV-1, HBV and HCV applications and effectiveness against Methicillin Resistant *Staphylococcus aureus* (MRSA), Vancomycin Resistant *Enterococcus faecalis* (VRE) and *Staphylococcus aureus* with reduced susceptibility to vancomycin: Use a second MetriWipes towelette to thoroughly wet the surface. Repeated use of the product may be required to ensure that the surface remains visibly wet for 2 minutes at room temperature (68 °F/20 °C).

For use as a disinfectant:

Use a second MetriWipes towelette to thoroughly wet the surface. Repeated use of the product may be required to ensure that the surface remains visibly wet for 3 minutes at room temperature (68 °F/20 °C) for Tuberculocidal activity and effectiveness against *Staphylococcus aureus*, *Pseudomonas aeruginosa*, *Trichophyton mentagrophytes* and *Salmonella enterica*.

To disinfect external surfaces of ultrasound transducers:

Pre-clean ultrasound gel from external transducer to be disinfected. Then thoroughly wet the probe with a MetriWipes towelette. Repeated use of the product may be required to ensure that the surface remains visibly wet for 3 minutes at 20 °C.

**A thorough rinse with potable water is required for surfaces in direct contact with food.

MetriWipes effectively kills Human Immunodeficiency Virus Type 1 (HIV), Hepatitis B Virus (HBV) and Hepatitis C Virus (HCV) on pre-cleaned environmental surfaces/objects previously soiled with blood/body fluids in healthcare settings or other settings in which there is an expected likelihood of soiling of inanimate surfaces/objects with blood/body fluids, and in which the surfaces/objects can be associated with the potential for transmission of HIV, HBV and HCV.



LICENSED

PERIOD 2012-2014 LIC. NO.



LICENSED

PERIOD 2015-2017 LIC. NO.

9591.72

Special Instructions for Cleaning and Decontamination Against HIV-1, HBV and HCV of Surfaces or Objects Soiled With Blood or Body Fluids:

Personal Protection: When using this disinfecting towelette, wear appropriate barrier protection such as disposable protective gloves, protective gown, face mask and eye covering.

Cleaning Procedure: Blood and other body fluids must be thoroughly cleaned from surfaces and objects before application of this disinfecting towelette. A MetriWipesXL towelette can be used for this purpose.

Contact Time: Thoroughly wet surface with a new MetriWipesXL towelette. Repeated use of the product may be required to ensure that the surface remains visibly wet for 2 minutes. This contact time will not be sufficient for some of the organisms listed on the label. Refer to label for contact times against other organisms.

Disposal Of Infectious Materials: Cleaning materials used that may contain blood or other body fluids must be autoclaved and disposed of in accordance with Federal, State and local regulations for infectious materials disposal.

This product is not to be used as a terminal sterilant/high-level disinfectant on any surface or instrument that (1) is introduced directly into the human body, either into or in contact with the bloodstream or normally sterile areas of the body, or (2) contacts intact mucous membranes but which does not ordinarily penetrate the blood barrier or otherwise enter normally sterile areas of the body. This product may be used to pre-clean or decontaminate critical or semi-critical medical devices prior to sterilization/high-level disinfection.

PRECAUTIONARY STATEMENTS Hazards to Humans and Domestic Animals CAUTION

Harmful if absorbed through the skin. Causes moderate eye irritation. Avoid contact with eyes, skin, or clothing. Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet. Remove contaminated clothing and wash clothing before reuse.

FIRST AID

If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing. Call a poison control center or doctor for further advice.

If on skin: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

For Chemical Emergency ONLY (spill, leak, fire, exposure or accident), call CHEMTREC at 1-800-424-9300. For ALL other inquiries about this product, call Metrex Customer Service at 1-800-841-1428, Monday through Friday, between 6 a.m. and 4 p.m. Pacific Time.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment.

STORAGE AND DISPOSAL

Towelette - Do not reuse towelette. Dispose of used towelette in trash.

Dispenser - Nonrefillable container. Do not reuse or refill this container. Offer for recycling, if available.

For product or technical information, contact Metrex at 800-841-1428 or visit our website at www.metrex.com.

Packed For: METREX® RESEARCH 28210 Wick Road Romulus, MI 48174

MADE IN USA

REORDER No. XX-XXX

70-xxxx-1
xxxx-xx



MetriWipes®

Extra Large Disinfecting Towelettes

- TUBERCULOCIDAL
- BACTERICIDAL
- VIRUCIDAL*
- FUNGICIDAL



Net Contents: X lbs. x oz.
65 Pre-Saturated Towelettes (9" x 12")

MetriWipes Disinfecting Towelettes are non-woven disposable towelettes. MetriWipes are intended for use in health care settings such as hospitals, dental offices, laboratories, clinics, ophthalmic offices, veterinary facilities and in kitchens** and bathrooms. MetriWipes are effective against the following microorganisms on hard, non-porous surfaces when used as directed:

- *Mycobacterium tuberculosis* var: *bovis* (BCG)
- *Staphylococcus aureus*
- *Pseudomonas aeruginosa*
- *Salmonella enterica*
- *Trichophyton mentagrophytes*
- Methicillin Resistant *Staphylococcus aureus* (MRSA)
- Vancomycin Resistant *Enterococcus faecalis* (VRE)
- *Staphylococcus aureus* with Reduced Susceptibility to Vancomycin
- Hepatitis B Virus (HBV)*
- Hepatitis C Virus (HCV)*
- Herpes Simplex Virus Types 1 and 2*
- Human Immunodeficiency Virus (HIV-1)*
- Influenza A2 Virus*

ACTIVE INGREDIENTS:

Diisobutylphenoxyethoxyethyl dimethyl benzyl ammonium chloride 0.28%
Isopropanol 17.20%
INERT INGREDIENTS†: 82.52%
TOTAL 100.00%

† Does not include the weight of the dry towelette.



KEEP OUT OF REACH OF CHILDREN CAUTION

See other panel for additional precautionary statements.

EPA Reg. No. 46781-9
EPA Est. No. 56952-WI-001

DO NOT USE AS A BABY WIPE. Not For Use On Skin.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

MetriWipes Disinfecting Towelettes are non-woven disposable towelettes. MetriWipes are intended for use on hard, non-porous surfaces in health care settings such as hospitals, emergency medical settings, neonatal units, surgical centers, isolation areas, out-patient surgical centers, nursing homes, dental offices, ophthalmic offices, eye surgical centers, patient care areas, dialysis centers, donation centers (blood, plasma, semen, milk, aphaeresis), veterinary facilities, salon settings, emergency vehicles, correctional facilities, schools, health club facilities, tanning salons, kitchens**, non-food contact surfaces in food preparation areas**, bathrooms, day care centers, police and fire vehicles, cadaver processing areas, laundry rooms, and other critical care areas where control of cross contamination between treated surfaces is required.

MetriWipes will clean and disinfect, when used as directed, such items as interior and exterior surfaces of infant incubators, bassinets, cribs and warmers, infant/child care equipment surfaces, diaper changing stations, oxygen hoods, operating room tables and lights, laboratory equipment and surfaces, physical therapy equipment surfaces, ambulance equipment surfaces, slit lamps, hard, non-porous exterior surfaces of ultrasound transducers or probes, exam tables, gurneys, IV poles, dental unit instrument trays, operatory light switches, exterior surfaces of amalgamators and dental curing lights, exterior surfaces of anesthesia machines and respiratory therapy equipment, SilkPeel™ with DermalInfusion™, workstations, dental countertops, loupes, exterior surfaces of endodontic equipment, such as apex locators, pulp testers and motors, stools, nail/hair care implements, tanning beds, salon surfaces, non-food contact surfaces**, exterior surfaces of refrigerator units**, exterior surfaces of appliances such as refrigerators, microwaves, food processors and stoves**, vanity tops**, floors, walls, handrails, door knobs, bed railings, tables, chairs, bathroom fixtures, exterior toilet surfaces, garbage cans, keyboards, carts, shopping carts, weight-lifting surfaces, drained whirlpool tank surfaces, cabinets, lights, telephones, which are made of plastic (i.e., polycarbonate, polypropylene, polyvinylchloride, polystyrene), and use sites listed on this label consisting of non-porous vinyl, stainless steel, painted surfaces, Plexiglas®, Formica® and glass.

Kitchen** and Bath Wipes for Hard, Non-Porous Surfaces Cleans, Deodorizes and Kills Common Household Germs

Dispenser Directions:

To start feed, remove cover and discard seal. From center of towelette roll, pull up a towelette corner, twist it into a point and thread it through the hole in the container cover. Pull through about one inch. Replace cover. Pull out first towelette and tear off at a 90° angle. Remaining towelettes feed automatically, ready for next use. When not in use, keep center cap closed to prevent solution loss.

Cleaning Instructions:

Use one MetriWipes towelette to completely preclean surface of all gross debris. For use as a virucide*, including HIV-1, HBV and HCV applications and effectiveness against Methicillin Resistant *Staphylococcus aureus* (MRSA), Vancomycin Resistant *Enterococcus faecalis* (VRE) and *Staphylococcus aureus* with reduced susceptibility to vancomycin: Use a second MetriWipes towelette to thoroughly wet the surface. Repeated use of the product may be required to ensure that the surface remains visibly wet for 2 minutes at room temperature (68 °F/20 °C).

For use as a disinfectant:

Use a second MetriWipes towelette to thoroughly wet the surface. Repeated use of the product may be required to ensure that the surface remains visibly wet for 3 minutes at room temperature (68 °F/20 °C) for Tuberculocidal activity and effectiveness against *Staphylococcus aureus*, *Pseudomonas aeruginosa*, *Trichophyton mentagrophytes* and *Salmonella enterica*.

To disinfect external surfaces of ultrasound transducers:

Pre-clean ultrasound gel from external transducer to be disinfected. Then thoroughly wet the probe with a MetriWipes towelette. Repeated use of the product may be required to ensure that the surface remains visibly wet for 3 minutes at 20 °C.

**A thorough rinse with potable water is required for surfaces in direct contact with food.

MetriWipes effectively kills Human Immunodeficiency Virus Type 1 (HIV), Hepatitis B Virus (HBV) and Hepatitis C Virus (HCV) on pre-cleaned environmental surfaces/objects previously soiled with blood/body fluids in healthcare settings or other settings in which there is an expected likelihood of soiling of inanimate surfaces/objects with blood/body fluids, and in which the surfaces/objects can be associated with the potential for transmission of HIV, HBV and HCV.