

CAVICIDE AF 8.OZ BOTTLE US
File Name: 70-5080-1_CCAF_8.OZ_BOTTLE_BASE_US_P9

SPECS:

Dimensions: 4-1/4" Height x 6-1/8" Length (+/-1/8")

BOOKLET SIZE (flat): 4-1/4" x 5-7/16"

BOOKLET SIZE (folded): 4-1/4" x 2-3/4"

Color: 100% Black, PMS 220, UV Varnish

43802



Department of Agriculture
STATE OF HAWAII

LICENSED

PERIOD 2015-2017 LIC. NO.

9591.77

PEEL HERE

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

KEEP OUT OF REACH OF CHILDREN. Causes moderate eye irritation. Avoid contact with eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet.

FIRST AID

Have the product container or label with you when calling a poison control center or doctor, or going for treatment.

If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

If swallowed: Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor. Do not give anything by mouth to an unconscious person.

If inhaled: Move person to fresh air. If person is not breathing, call 911 or an ambulance then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control center or doctor for further treatment advice.

NOTE TO PHYSICIAN

Probable mucosal damage may contraindicate the use of gastric lavage.

STORAGE AND DISPOSAL

DO NOT CONTAMINATE WATER, FOOD, OR FEED BY STORAGE OR DISPOSAL.

PESTICIDE STORAGE - Store in a dry place no lower in temperature than 50°F or higher than 120°F.

CONTAINER DISPOSAL - Nonrefillable container. Do not reuse or refill this container. Clean container promptly after emptying. Triple rinse as follows: Fill container ¼ full with water and recap. Agitate vigorously. Follow Pesticide Disposal instructions for rinsate disposal. Drain for 10 seconds after flow begins to drip. Repeat procedure two more times. Then offer for recycling or reconditioning. If not available, puncture and dispose in a sanitary landfill.

SPILL OR LEAK PROCEDURES - Small spills may be mopped up, flushed away with water or absorbed on some absorbent material and incinerated. Large spills should be contained, the material then moved into containers and disposed of by approved methods for hazardous wastes.

PESTICIDE DISPOSAL - Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

MADE IN USA

70-5080-1

CaviCide™
AF
disinfectant
cleaner

Ready to Use

- Tuberculocidal
- Bactericidal
- *Virucide

ACTIVE INGREDIENTS:

n-Alkyl (60% C₁₆, 30% C₁₈, 5% C₁₇, 5% C₁₈)
dimethyl benzyl ammonium chlorides 0.105%

n-Alkyl (68% C₁₇, 32% C₁₈)
dimethyl ethylbenzyl ammonium chlorides 0.105%

INERT INGREDIENTS: 99.790%
TOTAL 100.000%

KEEP OUT OF REACH OF CHILDREN

CAUTION

See side panels for additional precautionary statements.

Metrex™

Net Contents: 8 fl. oz. (239 mL)

Reorder No. 13-8008

EPA Reg. No. 1839-225-46781

EPA Est. No. 46781-MF-1

Manufactured For:
METREX™ RESEARCH
28210 Wick Road., Romulus, Michigan 48174 • 1-800-841-1428

LEAFLET PLACEMENT

CAVICIDE AF 8.OZ BOTTLE US

File Name: 70-5080-1_CCAF_8.OZ_BOTTLE_FOLD_US_P9

SPECS:
Dimensions: 4-1/4" Height x 6-1/8" Length (+/-1/8")
BOOKLET SIZE (flat): 4-1/4" x 5-7/16"
BOOKLET SIZE (folded): 4-1/4" x 2-3/4"
Color: 100% Black, PMS 220, Semi Gloss paper

112-301-944

to remain wet for **3 minutes**.
***VIRUCIDAL ACTIVITY** – CaviCide AF, when used on environmental, inanimate hard surfaces exhibits virucidal activity against: **Human Coronavirus, SARS associated Coronavirus, Avian Influenza A virus strain H3N2 and Avian Influenza A virus strain H9N2 (Avian Influenza virus strains H3N2 and H9N2) for 2 minutes; Influenza A virus strain H1N1, Paramyxovirus, Rhinovirus type 39 and Rotavirus for 3 minutes; HIV-1 (associated with AIDS) for 1 minute (60 seconds); and Norovirus (Norwalk virus), Feline Calicivirus and Rabies Virus for 30 seconds.**

KILLS HIV-1 ON PRE-CLEANED ENVIRONMENTAL SURFACES/OBJECTS PREVIOUSLY SOILED WITH BLOOD/BODY FLUIDS in healthcare settings (Hospitals, Nursing Homes) or other settings in which there is an expected likelihood of soiling of inanimate surfaces/objects with blood or body fluids, and in which the surfaces/objects likely to be soiled with blood or body fluids can be associated with the potential for transmission of Human Immunodeficiency Virus Type 1 (HIV-1) (associated with AIDS).

SPECIAL INSTRUCTIONS FOR CLEANING AND DECONTAMINATION AGAINST HIV-1 ON SURFACES/OBJECTS SOILED WITH BLOOD/BODY FLUIDS:
PERSONAL PROTECTION: When handling items soiled with blood or body fluids use disposable latex gloves, gowns, masks, and eye coverings.
CLEANING PROCEDURES: Blood and other body fluids must be thoroughly cleaned from surfaces and objects before application of this product.
CONTACT TIME: Allow surface to remain wet for **1 minute**. This contact time will not control other common types of viruses and bacteria.
DISPOSAL OF INFECTIOUS MATERIALS: Blood and other body fluids must be autoclaved and disposed of according to local regulations for infectious waste disposal.

EFFICACY TESTS HAVE DEMONSTRATED THAT CAVICIDE AF IS AN EFFECTIVE BACTERICIDE, FUNGICIDE, AND VIRUCIDE IN THE PRESENCE OF ORGANIC SOIL (5% BLOOD SERUM).

MILDEWSTAT - To control mold and mildew on pre-cleaned, hard, non-porous surfaces spray surface to be treated making sure to wet completely. Let air dry. Repeat application at weekly intervals or when mildew growth appears.

See expanded label for instructions and additional information.

Reorder No. 13-8008

Manufactured For:
METREX™ RESEARCH
28210 Wick Road., Romulus,
Michigan 48174
1-800-841-1428

For Chemical Emergency ONLY (spill, leak, fire, exposure or accident), call CHEMTREC at 1-800-424-9300. For ALL other inquiries about this product, call Metrex Customer Service at 1-800-841-1428, Monday through Friday, between 6 a.m. and 4 p.m. Pacific Time.

For product or technical information, contact Metrex at **800-841-1428** or visit our website at www.metrex.com.

MADE IN USA

70-5080-1



+H97113100846

CaviCide AF is designed specifically as a general ready-to-use, nonabrasive cleaner and disinfectant for use in homes, hospitals, nursing homes, patient rooms, operating rooms, ICU areas, shower rooms, locker rooms, public restrooms, schools, food processing plants, food service establishments, transportation terminals, office buildings, manufacturing facilities, lodging establishments, retail businesses, veterinary clinics, pet shops, animal life science laboratories and athlete/recreational facilities where housekeeping is of prime importance in controlling the hazard of cross contamination. It is formulated to disinfect hard, non-porous, inanimate environmental surfaces: floors, walls, metal surfaces, stainless steel surfaces, glazed porcelain, glazed ceramic tile, plastic surfaces, chrome, brass, copper, laminated surfaces, baked enamel surfaces, bathrooms, shower stalls, bathtubs, and cabinets. **For plastic and painted surfaces, spot test on an inconspicuous area before use.** May be used in the kitchen on counters, sinks, appliances, and stovetops. A rinse with potable water is required for surfaces in direct contact with food. In addition, **CaviCide AF** deodorizes those areas that generally are hard to keep fresh smelling, such as garbage storage areas, empty garbage bins and cans, basements, restrooms and other areas which are prone to odors caused by microorganisms.

CaviCide AF is a one-step hospital-use germicidal disinfectant cleaner and deodorant designed for general cleaning, and disinfecting of hard, non-porous inanimate surfaces. Quickly removes dirt, grime, mold, mildew, food residue, blood and other organic matter commonly found in hospitals and other healthcare facilities on medical surfaces. It also eliminates odors leaving surfaces smelling clean and fresh. Use where odors are a problem.

DIRECTIONS FOR USE
It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. This product is not for use on critical or semi-critical medical device surfaces.
DISINFECTION, DEODORIZING AND CLEANING: Remove gross filth or heavy soil prior to application of the product. Hold container six to eight inches from surface to be treated. Spray area until it is covered with the solution. Allow product to penetrate and the surface to remain wet for **3 minutes** (for HIV-1, allow surface to remain wet for **1 minute**). No scrubbing is necessary. Wipe off with a clean cloth, mop or sponge. The product will not leave grit or soap scum.

To disinfect food processing premises: floors, walls and storage areas, hold container six to eight inches from the surface to be treated and spray area until it is covered with the solution. Allow **CaviCide AF** to penetrate and remain wet for **3 minutes** (for HIV-1 allow surface to remain wet for **1 minute**). Wipe off with a clean cloth, mop or sponge and thoroughly rinse all surfaces in the area with potable water. Heavily soiled areas must be pre-cleaned before using this product. Food products and packaging materials must be removed or carefully protected prior to using this product.

To disinfect food service establishment food contact surfaces: countertops, appliances, tables, hold container six to eight inches from the surface to be treated and spray area until it is covered with the solution. Allow **CaviCide AF** to penetrate and remain wet for **3 minutes** (for HIV-1 allow surface to remain wet for **1 minute**). Wipe off with a clean cloth, mop or sponge and thoroughly rinse all surfaces in the area with potable water. Heavily soiled areas must be pre-cleaned before using this product. Do not use on eating utensils, glassware and cookware.

Disinfection of animal quarters and kennels: For disinfection of pre-cleaned animal quarters and kennels, apply **CaviCide AF**. Remove all animals and feed from premises. Remove all litter and droppings from floors, walls and surfaces of facilities occupied or traversed by animals. Empty all troughs, racks, and other feeding and watering appliances. Thoroughly clean all surfaces with soap or detergent and rinse with water. Saturate the surfaces with the disinfecting solution for a period of **3 minutes** (for Feline Calicivirus and Rabies virus **30 seconds**). Ventilate building and other closed spaces. Do not house animals or employ equipment until treatment has been absorbed, set or dried. All treated equipment that will contact feed or drinking water must be rinsed with potable water before reuse.

This product **CaviCide AF** has demonstrated effectiveness against Influenza A virus and is expected to inactivate all Influenza A viruses including Pandemic 2009 H1N1 Influenza A virus.

Using approved AOAC test methods (GLPs), in the presence 5% blood serum and a **3 minute** contact time, unless otherwise noted, **CaviCide AF** kills the following organisms on hard non-porous inanimate surfaces:

BACTERICIDAL ACTIVITY (BACTERIA):
Pseudomonas aeruginosa, Staphylococcus aureus, Salmonella (choleraesuis) enterica, Escherichia coli, Escherichia coli O157:H7, Streptococcus pyogenes (Necrotizing Fasciitis-Group A), Listeria monocytogenes, Yersinia enterocolitica, Enterococcus faecium, Corynebacterium ammoniagenes, Salmnella (typhi) enterica

ANTIBIOTIC-RESISTANT BACTERICIDAL ACTIVITY:
Methicillin resistant *Staphylococcus aureus* (MRSA), Vancomycin resistant *Enterococcus faecalis* (VRE), Vancomycin intermediate resistant *Staphylococcus aureus* (VISA), Methicillin resistant *Staphylococcus epidermidis* (MRSE), Community Associated Methicillin resistant *Staphylococcus aureus* (CA-MRSA) (NRS 123), Genotype USA400, Community Associated Methicillin resistant *Staphylococcus aureus* (CA-MRSA) (NRS 384), Genotype USA300

TUBERCULOCIDAL ACTIVITY: **CaviCide AF** exhibits disinfectant efficacy against *Mycobacterium tuberculosis* BCG (*Mycobacterium bovis*) at 20°C when the treated surface is allowed

3

4

FPO
BARCODE
TO COME

1

2

CAVICIDE AF 24 OZ. BOTTLE US

File Name: 70-5082-1_CCAF_24.OZ_BOTTLE_US_P7



LICENSED

PERIOD 2015-2017 LIC. NO.

9591.77

SPECS:

Dimensions: 3.875" x 10" with 0.125" radious corners

Color: 100% Black, PMS 220, UV Varnish

CaviCide™ AF

disinfectant
cleaner

Ready-to-Use

• TUBERCULOCIDAL • BACTERICIDAL • *VIRUCIDAL

ACTIVE INGREDIENTS:

n-Alkyl (60% C ₁₄ , 30% C ₁₆ , 5% C ₁₂ , 5% C ₁₈) dimethyl benzyl ammonium chlorides	0.105%
n-Alkyl (68% C ₁₂ , 32% C ₁₄) dimethyl ethylbenzyl ammonium chlorides	0.105%
INERT INGREDIENTS:	99.790%
TOTAL	100.000%



Net Contents: 24 fl. oz. (709 mL)

EPA Reg. No. 1839-225-46781 EPA Est. No. 46781-MI-1

KEEP OUT OF REACH OF CHILDREN
CAUTION

See side panels for additional precautionary statements.

REORDER No. 13-8024

This product **CaviCide™ AF** has demonstrated effectiveness against Influenza A virus and is expected to inactivate all Influenza A viruses including Pandemic 2009 H1N1 Influenza A virus.

CaviCide AF is designed specifically as a general ready-to-use, nonabrasive cleaner and disinfectant for use in homes, hospitals, nursing homes, patient rooms, operating rooms, ICU areas, shower rooms, locker rooms, public restrooms, schools, food processing plants, food service establishments, transportation terminals, office buildings, manufacturing facilities, lodging establishments, retail businesses, veterinary clinics, pet shops, animal life science laboratories and athletic/recreational facilities where housekeeping is of prime importance in controlling the hazard of cross contamination. It is formulated to disinfect hard, non-porous, inanimate environmental surfaces: floors, walls, metal surfaces, stainless steel surfaces, glazed porcelain, glazed ceramic tile, plastic surfaces, chrome, brass, copper,

laminated surfaces, baked enamel surfaces, bathrooms, shower stalls, bathtubs, and cabinets. For plastic and painted surfaces, spot test on an inconspicuous area before use. A rinse with potable water is required for surfaces in direct contact with food. In addition, **CaviCide AF** deodorizes those areas that generally are hard to keep fresh smelling, such as garbage storage areas, empty garbage bins and cans, basements, restrooms and other areas which are prone to odors caused by microorganisms. **CaviCide AF** is a one-step hospital-use germicidal disinfectant cleaner and deodorant designed for general cleaning, and disinfecting of hard, non-porous inanimate surfaces. Quickly removes dirt, grime, mold, mildew, food residue, blood and other organic matter commonly found in hospitals and other healthcare facilities on medical surfaces. It also eliminates odors leaving surfaces smelling clean and fresh. Use where odors are a problem.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. This product is not for use on critical or semi-critical medical device surfaces.

DISINFECTION, DEODORIZING AND CLEANING: Remove gross filth or heavy soil prior to application of the product. Hold container six to eight inches from surface to be treated. Spray area until it is covered with the solution. Allow product to penetrate and the surface to remain wet for **3 minutes** (for HIV-1, allow surface to remain wet for **1 minute**). No scrubbing is necessary. Wipe off with a clean cloth, mop or sponge. The product will not leave grit or soap scum.

To disinfect food processing premises: floors, walls and storage areas, hold container six to eight inches from the surface to be treated and spray area until it is covered with the solution. Allow **CaviCide AF** to penetrate and remain wet for **3 minutes** (for HIV-1 allow surface to remain wet for **1 minute**). Wipe off with a clean cloth, mop or sponge and thoroughly rinse all surfaces in the area with potable water. Heavily soiled areas must be pre-cleaned before using this product. Food products and packaging materials must be removed or carefully protected prior to using this product.

To disinfect food service establishment food contact surfaces: countertops, appliances, tables, hold container six to eight inches from the surface to be treated and spray area until it is covered with the solution. Allow **CaviCide AF** to penetrate and remain wet for **3 minutes** (for HIV-1 allow surface to remain wet for **1 minute**). Wipe off with a clean cloth, mop or sponge and thoroughly rinse all surfaces in the area with potable water. Heavily soiled areas must be pre-cleaned before using this product. Do not use on eating utensils, glassware and cookware.

Disinfection of animal quarters and kennels: For disinfection of pre-cleaned animal quarters and kennels, apply **CaviCide AF**. Remove all animals and feed from premises. Remove all litter and droppings from floors, walls and surfaces of facilities occupied or traversed by animals. Empty all troughs, racks, and other feeding and watering appliances. Thoroughly clean all surfaces with soap or detergent and rinse with water. Saturate the surfaces with the disinfecting solution for a period of **3 minutes** (for Feline Calicivirus and Rabies virus **30 seconds**). Ventilate building and other closed spaces. Do not house animals or employ equipment until treatment has been absorbed, set or dried. All treated equipment that will contact feed or drinking water must be rinsed with potable water before reuse.

Using approved AOAC test methods (under Good Laboratory Practices, [GLPs]), in the presence of 5% blood serum and a **3 minute** contact time, unless otherwise noted, **CaviCide AF** kills the following organisms on hard non-porous inanimate surfaces:

BACTERICIDAL ACTIVITY (BACTERIA):

Pseudomonas aeruginosa, *Staphylococcus aureus*, *Salmonella (choleraesuis) enterica*, *Escherichia coli*, *Escherichia coli* O157:H7, *Streptococcus pyogenes* (Necrotizing Fasciitis-Group A), *Listeria monocytogenes*, *Yersinia enterocolitica*, *Enterococcus faecium*, *Corynebacterium ammoniagenes*, *Salmonella (typhi) enterica*,

ANTIBIOTIC-RESISTANT BACTERICIDAL ACTIVITY:

Methicillin resistant *Staphylococcus aureus* (MRSA), Vancomycin resistant *Enterococcus faecalis* (VRE), Vancomycin intermediate resistant *Staphylococcus aureus* (VISA), Methicillin resistant *Staphylococcus epidermidis* (MRSE), Community Associated Methicillin resistant *Staphylococcus*

aureus (CA-MRSA), (NRS 123) Genotype USA400, Community Associated Methicillin resistant *Staphylococcus aureus* (CA-MRSA), (NRS 384) Genotype USA300

TUBERCULOCIDAL ACTIVITY: **CaviCide AF** exhibits disinfectant efficacy against *Mycobacterium tuberculosis* BCG (*Mycobacterium bovis*) at 20°C when the treated surface is allowed to remain wet for **3 minutes**.

***VIRUCIDAL ACTIVITY – CaviCide AF**, when used on environmental, inanimate hard surfaces exhibits virucidal activity against: **Human Coronavirus SARS associated Coronavirus**, **Avian Influenza A virus strain H3N2 and Avian Influenza A virus strain H9N2 (Avian Influenza virus strains H3N2 and H9N2) for 2 minutes**; **Influenza A virus strain H1N1**, **Parvovirus**, **Rhinovirus type 39 and Rotavirus for 3 minutes**; **HIV-1 (associated with AIDS) for 1 minute (60 seconds)**; and **Norovirus (Norwalk virus), Feline Calicivirus and Rabies Virus for 30 seconds**.

KILLS HIV-1 ON PRE-CLEANED ENVIRONMENTAL SURFACES/OBJECTS PREVIOUSLY SOILED WITH BLOOD/BODY FLUIDS in healthcare settings (Hospitals, Nursing Homes) or other settings in which there is an expected likelihood of soiling of inanimate surfaces/objects with blood or body fluids, and in which the surfaces/objects likely to be soiled with blood or body fluids can be associated with the potential for transmission of Human Immunodeficiency Virus Type 1 (HIV-1) (associated with AIDS).

SPECIAL INSTRUCTIONS FOR CLEANING AND DECONTAMINATION AGAINST HIV-1 ON SURFACES/OBJECTS SOILED WITH BLOOD/BODY FLUIDS: PERSONAL PROTECTION: When handling items soiled with blood or body fluids use disposable latex gloves, gowns, masks, and eye coverings.

CLEANING PROCEDURES: Blood and other body fluids must be thoroughly cleaned from surfaces and objects before application of this product.

CONTACT TIME: Allow surface to remain wet for **1 minute**. This contact time will not control other common types of viruses and bacteria.

DISPOSAL OF INFECTIOUS MATERIALS: Blood and other body fluids must be autoclaved and disposed of according to local regulations for infectious waste disposal.

EFFICACY TESTS HAVE DEMONSTRATED THAT CAVICIDE AF IS AN EFFECTIVE BACTERICIDE, FUNGICIDE, AND VIRUCIDE IN THE PRESENCE OF ORGANIC SOIL (5% BLOOD SERUM).

MILDEWSTAT - To control mold and mildew on pre-cleaned, hard, non-porous surfaces spray surface to be treated making sure to wet completely. Let air dry. Repeat application at weekly intervals or when mildew growth appears.

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS CAUTION

KEEP OUT OF REACH OF CHILDREN. Causes moderate eye irritation. Avoid contact with eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet.

FIRST AID

Have the product container or label with you when calling a poison control center or doctor, or going

for treatment.

If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

If swallowed: Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor. Do not give anything by mouth to an unconscious person.

If inhaled: Move person to fresh air. If person is not breathing, call 911 or an ambulance then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control center or doctor for further treatment advice.

NOTE TO PHYSICIAN

Probable mucosal damage may contraindicate the use of gastric lavage.

STORAGE AND DISPOSAL

DO NOT CONTAMINATE WATER, FOOD, OR FEED BY STORAGE OR DISPOSAL.

PESTICIDE STORAGE - Store in a dry place no lower in temperature than 50°F or higher than 120°F. **CONTAINER DISPOSAL -** Nonrefillable container. Do not reuse or refill this container. Clean container promptly after emptying. Triple rinse as follows: Fill container ¼ full with water and recap. Agitate vigorously. Follow Pesticide Disposal instructions for rinsate disposal. Drain for 10 seconds after flow begins to drip. Repeat procedure two more times. Then offer for recycling or reconditioning. If not available, puncture and dispose in a sanitary landfill.

SPILL OR LEAK PROCEDURES - Small spills may be mopped up, flushed away with water or absorbed on some absorbent material and incinerated. Large spills should be contained, the material then moved into containers and disposed of by approved methods for hazardous wastes.

PESTICIDE DISPOSAL - Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

Manufactured For:
METREX™ RESEARCH, 28210 Wick Road, Romulus, Michigan 48174 • 1-800-841-1428

For Chemical Emergency ONLY (spill, leak, fire, exposure or accident), call CHEMTREC at 1-800-424-9300. For ALL other inquiries about this product, call Metrex Customer Service at 1-800-841-1428, Monday through Friday, between 6 a.m. and 4 p.m. Pacific Time.

For product or technical information, contact Metrex at 800-841-1428 or visit our website at www.metrex.com.

MADE IN USA



+79113802407

70-5082-1

CAVICIDE AF 1 GALLON US
File Name: 70-5084-1_CCAF_1_GALLON_US_P11



LICENSED

PERIOD 2015-2017 LIC. NO.

9591.77

SPECS:
Dimensions: 4.75" x 17.8125" with 0.125" radius corners
Color: 100% Black, PMS 220, UV Varnish

This product **CaviCide™ AF** has demonstrated effectiveness against Influenza A virus and is expected to inactivate all Influenza A viruses including Pandemic 2009 H1N1 Influenza A virus.

CaviCide AF is designed specifically as a general ready-to-use, nonabrasive cleaner and disinfectant for use in homes, hospitals, nursing homes, patient rooms, operating rooms, ICU areas, shower rooms, locker rooms, public restrooms, schools, food processing plants, food service establishments, transportation terminals, office buildings, manufacturing facilities, lodging establishments, retail businesses, veterinary clinics, pet shops, animal life science laboratories and athlete/recreational facilities where housekeeping is of prime importance in controlling the hazard of cross contamination. It is formulated to disinfect hard, non-porous, inanimate environmental surfaces: floors, walls, metal surfaces, stainless steel surfaces, glazed porcelain, glazed ceramic tile, plastic surfaces, chrome, brass, copper, laminated surfaces, baked enamel surfaces, bathrooms, shower stalls, bathtubs, and cabinets. **For plastic and painted surfaces, spot test on an inconspicuous area before use.** May be used in the kitchen on counters, sinks, appliances, and stovetops. A rinse with potable water is required for surfaces in direct contact with food. In addition, **CaviCide AF** deodorizes those areas that generally are hard to keep fresh smelling, such as garbage storage areas, empty garbage bins and cans, basements, restrooms and other areas which are prone to odors caused by microorganisms.

CaviCide AF is a one-step hospital-use germicidal disinfectant cleaner and deodorant designed for general cleaning, and disinfecting of hard, non-porous inanimate surfaces. Quickly removes dirt, grime, mold, mildew, food residue, blood and other organic matter commonly found in hospitals and other healthcare facilities on medical surfaces. It also eliminates odors leaving surfaces smelling clean and fresh. Use where odors are a problem.

DIRECTIONS FOR USE
It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

This product is not to be used as a terminal sterilant/high-level disinfectant on any surface or instrument that (1) is introduced directly into the human body, either into or in contact with the bloodstream or normally sterile areas of the body, or (2) contacts intact mucous membranes but which does not ordinarily penetrate the blood barrier or otherwise enter normally sterile areas of the body.

When used as directed, **CaviCide AF** contains 2100 ppm of active germicide making it highly effective against a wide variety (broad-spectrum) of pathogenic microorganisms including bacteria, antibiotic resistant bacteria, viruses, fungi, mold and mildew.

DISINFECTION, DEODORIZING AND CLEANING: Remove gross filth or heavy soil prior to application of the product. Hold container six to eight inches from surface to be treated. Spray area until it is covered with the solution. Allow product to penetrate and the surface to remain wet for **3 minutes** (for HIV-1, allow surface to remain wet for **1 minute**). No scrubbing is necessary. Wipe off with a clean cloth, mop or sponge. The product will not leave grit or soap scum.

To disinfect food processing premises: floors, walls and storage areas, hold container six to eight inches from the surface to be treated and spray area until it is covered with the solution. Allow **CaviCide AF** to penetrate and remain wet for **3 minutes** (for HIV-1 allow surface to remain wet for **1 minute**). Wipe off with a clean cloth, mop or sponge and thoroughly rinse all surfaces in the area with potable water. Heavily soiled areas must be pre-cleaned before using this

product. Food products and packaging materials must be removed or carefully protected prior to using this product.

To disinfect food service establishment food contact surfaces: countertops, appliances, tables, hold container six to eight inches from the surface to be treated and spray area until it is covered with the solution. Allow **CaviCide AF** to penetrate and remain wet for **3 minutes** (for HIV-1 allow surface to remain wet for **1 minute**). Wipe off with a clean cloth, mop or sponge and thoroughly rinse all surfaces in the area with potable water. Heavily soiled areas must be pre-cleaned before using this product. Do not use on eating utensils, glassware and cookware.

Disinfection of animal quarters and kennels: For disinfection of pre-cleaned animal quarters and kennels, apply **CaviCide AF**. Remove all animals and feed from premises. Remove all litter and droppings from floors, walls and surfaces of facilities occupied or traversed by animals. Empty all troughs, racks, and other feeding and watering appliances. Thoroughly clean all surfaces with soap or detergent and rinse with water. Saturate the surfaces with the disinfecting solution for a period of **3 minutes** (for Feline Calicivirus and Rabies virus **30 seconds**). Ventilate building and other closed spaces. Do not house animals or employ equipment until treatment has been absorbed, set or dried. All treated equipment that will contact feed or drinking water must be rinsed with potable water before reuse.

Using approved AOAC test methods (under Good Laboratory Practices, [GLPs]), in the presence of 5% blood serum and a **3 minute** contact time, unless otherwise noted, **CaviCide AF** kills the following organisms on hard non-porous inanimate surfaces:

- BACTERICIDAL ACTIVITY (BACTERIA):**
Pseudomonas aeruginosa, (ATCC 15442)
Staphylococcus aureus, (ATCC 6538)
Salmonella (choleraesuis) enterica, (ATCC 10708)
Escherichia coli, (ATCC 11229)
Escherichia coli O157:H7, (ATCC 43895)
Streptococcus pyogenes (Necrotizing Fasciitis-Group A) (V.A. Medical Center Isolate 04001)
Listeria monocytogenes, (ATCC 35152)
Yersinia enterocolitica, (ATCC 23715)
Enterococcus faecium, (ATCC 6569)
Corynebacterium ammoniagenes, (ATCC 6871)
Salmonella (typhi) enterica, (ATCC 6539)

- ANTIBIOTIC-RESISTANT BACTERICIDAL ACTIVITY:**
Methicillin resistant *Staphylococcus aureus* (MRSA), (ATCC 33593)
Vancomycin resistant *Enterococcus faecalis* (VRE), (ATCC 51575)
Vancomycin intermediate resistant *Staphylococcus aureus* (VISA), (CDC Isolate 99287)
Methicillin resistant *Staphylococcus epidermidis* (MRSE), (ATCC 51625)
Community Associated Methicillin resistant *Staphylococcus aureus* (CA-MRSA), (NRS 123)
Genotype USA400
Community Associated Methicillin resistant *Staphylococcus aureus* (CA-MRSA), (NRS 384)
Genotype USA300

CaviCide™ AF

disinfectant
cleaner

Ready-to-Use

- TUBERCULOCIDAL • BACTERICIDAL • *VIRUCIDE

ACTIVE INGREDIENTS:

n-Alkyl (60% C ₁₄ , 30% C ₁₆ , 5% C ₁₂ , 5% C ₁₈)	
dimethyl benzyl ammonium chlorides	0.105%
n-Alkyl (68% C ₁₂ , 32% C ₁₄)	
dimethyl ethylbenzyl ammonium chlorides	0.105%
INERT INGREDIENTS:	99.790%
TOTAL	100.000%

KEEP OUT OF REACH OF CHILDREN
CAUTION

See side panels for additional precautionary statements.



Net Contents: 1 Gallon / 3.785 Liters

EPA Reg. No. 1839-225-46781
EPA Est. No. 46781-MI-1



TUBERCULOCIDAL ACTIVITY: **CaviCide AF** exhibits disinfectant efficacy against *Mycobacterium tuberculosis* BCG (*Mycobacterium bovis*) at 20°C when the treated surface is allowed to remain wet for **3 minutes**.

***VIRUCIDAL ACTIVITY – CaviCide AF**, when used on environmental, inanimate hard surfaces exhibits virucidal activity against: **Human Coronavirus** (ATCC VR-740), **SARS associated Coronavirus** (ZeploMetrix), **Avian Influenza A virus strain H3N2** (ATCC VR-2072) and **Avian Influenza A virus strain H9N2** (Charles River Laboratories) (**Avian Influenza virus strains H3N2 and H9N2**) for **2 minutes**; **Influenza A virus strain H1N1** (ATCC VR-1469), **Paramyxovirus** (Mumps) (ATCC VR-1438), **Rhinovirus type 39** (ATCC VR-340) and **Rotavirus** (University of Ottawa) for **3 minutes**; **HIV-1 (associated with AIDS)** (Advanced Biotechnologies, Inc.) for **1 minute (60 seconds)**; and **Norovirus (Norwalk virus)**, **Feline Calicivirus** (University of Ottawa) and **Rabies Virus** (CDC) for **30 seconds**.

KILLS HIV-1 ON PRE-CLEANED ENVIRONMENTAL SURFACES/OBJECTS PREVIOUSLY SOILED WITH BLOOD/BODY FLUIDS in healthcare settings (Hospitals, Nursing Homes) or other settings in which there is an expected likelihood of soiling of inanimate surfaces/objects with blood or body fluids, and in which the surfaces/objects likely to be soiled with blood or body fluids can be associated with the potential for transmission of Human Immunodeficiency Virus Type 1 (HIV-1) (associated with AIDS).

SPECIAL INSTRUCTIONS FOR CLEANING AND DECONTAMINATION AGAINST HIV-1 ON SURFACES/OBJECTS SOILED WITH BLOOD/BODY FLUIDS:

PERSONAL PROTECTION: When handling items soiled with blood or body fluids use disposable latex gloves, gowns, masks, and eye coverings.
CLEANING PROCEDURES: Blood and other body fluids must be thoroughly cleaned from surfaces and objects before application of this product.

CONTACT TIME: Allow surface to remain wet for **1 minute. This contact time will not control other common types of viruses and bacteria.**

DISPOSAL OF INFECTIOUS MATERIALS: Blood and other body fluids must be autoclaved and disposed of according to local regulations for infectious waste disposal.

EFFICACY TESTS HAVE DEMONSTRATED THAT CAVICIDE AF IS AN EFFECTIVE BACTERICIDE, FUNGICIDE, AND VIRUCIDE IN THE PRESENCE OF ORGANIC SOIL (5% BLOOD SERUM).

MILDEWSTAT - To control mold and mildew on pre-cleaned, hard, non-porous surfaces spray surface to be treated making sure to wet completely. Let air dry. Repeat application at weekly intervals or when mildew growth appears.

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
CAUTION**

KEEP OUT OF REACH OF CHILDREN. Causes moderate eye irritation. Avoid contact with eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet.

FIRST AID
Have the product container or label with you when calling a poison control center or doctor, or going for treatment.

If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

If swallowed: Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor. Do not give anything by mouth to an unconscious person.

If inhaled: Move person to fresh air. If person is not breathing, call 911 or an ambulance then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control center or doctor for further treatment advice.

NOTE TO PHYSICIAN
Probable mucosal damage may contraindicate the use of gastric lavage.

STORAGE AND DISPOSAL
DO NOT CONTAMINATE WATER, FOOD, OR FEED BY STORAGE OR DISPOSAL.
PESTICIDE STORAGE - Store in a dry place no lower in temperature than 50°F or higher than 120°F.
CONTAINER DISPOSAL - Nonrefillable container. Do not reuse or refill this container. Clean container promptly after emptying. Triple rinse as follows: Fill container ¼ full with water and recap. Agitate vigorously. Follow Pesticide Disposal instructions for rinsate disposal. Drain for 10 seconds after flow begins to drip. Repeat procedure two more times. Then offer for recycling or reconditioning, if not available, puncture and dispose in a sanitary landfill.
SPILL OR LEAK PROCEDURES – Small spills may be mopped up, flushed away with water or absorbed on some absorbent material and incinerated. Large spills should be contained, the material then moved into containers and disposed of by approved methods for hazardous wastes.
PESTICIDE DISPOSAL – Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

Manufactured For:
METREX™ RESEARCH
28210 Wick Road
Romulus, Michigan 48174
1-800-841-1428

REORDER No. 13-8000

For Chemical Emergency ONLY (spill, leak, fire, exposure or accident), call CHEMTREC at 1-800-424-9300. For ALL other inquiries about this product, call Metrex Customer Service at 1-800-841-1428, Monday through Friday, between 6 a.m. and 4 p.m. Pacific Time.

For product or technical information, contact Metrex at **800-841-1428** or visit our website at www.metrex.com.

MADE IN USA
70-5084-1 00000-22

CAVICIDE AF 2.5 GALLON US

File Name: 70-5094-1 CCAF_2.5_GALLON_US_P5



LICENSED

PERIOD 2015-2017 LIC. NO.

9591.77

SPECS:

Dimensions: 7.75" x 7.5" with 0.125" radius corners
Color: 100% Black, PMS 220, UV Varnish

This product **CaviCide™ AF** has demonstrated effectiveness against Influenza A virus and is expected to inactivate all Influenza A viruses including Pandemic 2009 H1N1 Influenza A virus.

CaviCide AF is designed specifically as a general ready-to-use, nonabrasive cleaner and disinfectant for use in homes, hospitals, nursing homes, patient rooms, operating rooms, ICU areas, shower rooms, locker rooms, public restrooms, schools, food processing plants, food service establishments, transportation terminals, office buildings, manufacturing facilities, lodging establishments, retail businesses, veterinary clinics, pet shops, animal life science laboratories and athlete/recreational facilities where housekeeping is of prime importance in controlling the hazard of cross contamination. It is formulated to disinfect hard, non-porous, inanimate environmental surfaces: floors, walls, metal surfaces, stainless steel surfaces, glazed porcelain, glazed ceramic tile, plastic surfaces, chrome, brass, copper, laminated surfaces, baked enamel surfaces, bathrooms, shower stalls, bathtubs, and cabinets. **For plastic and painted surfaces, spot test on an inconspicuous area before use.** May be used in the kitchen on counters, sinks, appliances, and stovetops. A rinse with potable water is required for surfaces in direct contact with food. In addition, **CaviCide AF** deodorizes those areas that generally are hard to keep fresh smelling, such as garbage storage areas, empty garbage bins and cans, basements, restrooms and other areas which are prone to odors caused by microorganisms.

CaviCide AF is a one-step hospital-use germicidal disinfectant cleaner and deodorant designed for general cleaning, and disinfecting of hard, non-porous inanimate surfaces. Quickly removes dirt, grime, mold, mildew, food residue, blood and other organic matter commonly found in hospitals and other healthcare facilities on medical surfaces. It also eliminates odors leaving surfaces smelling clean and fresh. Use where odors are a problem.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

This product is not to be used as a terminal sterilant/high-level disinfectant on any surface or instrument that (1) is introduced directly into the human body, either into or in contact with the bloodstream or normally sterile areas of the body, or (2) contacts intact mucous membranes but which does not ordinarily penetrate the blood barrier or otherwise enter normally sterile areas of the body.

When used as directed, **CaviCide AF** contains 2100 ppm of active germicide making it highly effective against a wide variety (broad-spectrum) of pathogenic microorganisms including bacteria, antibiotic resistant bacteria, viruses, fungi, mold and mildew.

DISINFECTION, DEODORIZING AND CLEANING: Remove gross filth or heavy soil prior to application of the product. Hold container six to eight inches from surface to be treated. Spray area until it is covered with the solution. Allow product to penetrate and the surface to remain wet for **3 minutes** (for HIV-1, allow surface to remain wet for **1 minute**). No scrubbing is necessary. Wipe off with a clean cloth, mop or sponge. The product will not leave grit or soap scum.

To disinfect food processing premises: floors, walls and storage areas, hold container six to eight inches from the surface to be treated and spray area until it is covered with the solution. Allow **CaviCide AF** to penetrate and remain wet for **3 minutes** (for HIV-1 allow surface to remain wet for **1 minute**). Wipe off with a clean cloth, mop or sponge and thoroughly rinse all surfaces in the area with potable water. Heavily soiled areas must be pre-cleaned before using this product. Food products and packaging materials must be removed or carefully protected prior to using this product.

To disinfect food service establishment food contact surfaces: countertops, appliances, tables, hold container six to eight inches from the surface to be treated and spray area until it is covered with the solution. Allow **CaviCide AF** to penetrate and remain wet for **3 minutes** (for HIV-1 allow surface to remain wet for **1 minute**). Wipe off with a clean cloth, mop or sponge and thoroughly rinse all surfaces in the area with potable water. Heavily soiled areas must be pre-cleaned before using this product. Do not use on eating utensils, glassware and cookware.

Disinfection of animal quarters and kennels: For disinfection of pre-cleaned animal quarters and kennels, apply **CaviCide AF**. Remove all animals and feed from premises. Remove all litter and droppings from floors, walls and surfaces of facilities occupied or traversed by animals. Empty all troughs, racks, and other feeding and watering appliances. Thoroughly clean all surfaces with soap or detergent and rinse with water. Saturate the surfaces with the disinfecting solution for a period of **3 minutes** (for Feline Calicivirus and Rabies virus **30 seconds**). Ventilate building and other closed spaces. Do not house animals or employ equipment until treatment has been absorbed, set or dried. All treated equipment that will contact feed or drinking water must be rinsed with potable water before reuse.

Using approved AOAC test methods (under Good Laboratory Practices, [GLPs]), in the presence of 5% blood serum and a **3 minute** contact time, unless otherwise noted, **CaviCide AF** kills the following organisms on hard non-porous inanimate surfaces:

BACTERICIDAL ACTIVITY (BACTERIA):

- Pseudomonas aeruginosa - Staphylococcus aureus - Salmonella (choleraesuis) enterica - Escherichia coli
- Escherichia coli 0157:H7 - Streptococcus pyogenes (Necrotizing Fasciitis-Group A) - Listeria monocytogenes - Yersinia enterocolitica - Enterococcus faecium - Corynebacterium ammoniogenes
- Salmonella (typhi) enterica

ANTIBIOTIC-RESISTANT BACTERICIDAL ACTIVITY:

- Methicillin resistant *Staphylococcus aureus* (MRSA), (ATCC 33593)
- Vancomycin resistant *Enterococcus faecalis* (VRE), (ATCC 51575)

CaviCide™ AF

disinfectant cleaner

Ready-to-Use

• TUBERCULOCIDAL • BACTERICIDAL • *VIRUCIDE

ACTIVE INGREDIENTS:

n-Alkyl (60% C ₁₄ , 30% C ₁₆ , 5% C ₁₂ , 5% C ₁₈)	
dimethyl benzyl ammonium chlorides	0.105%
n-Alkyl (68% C ₁₂ , 32% C ₁₄)	
dimethyl ethylbenzyl ammonium chlorides	0.105%
INERT INGREDIENTS:	99.790%
TOTAL	100.000%



FPO
BARCODE
TO COME



Net Contents: 2.5 Gallon

Vancomycin intermediate resistant *Staphylococcus aureus* (VISA), (CDC Isolate 99287)
Methicillin resistant *Staphylococcus epidermidis* (MRSE), (ATCC 51625)
Community Associated Methicillin resistant *Staphylococcus aureus* (CA-MRSA), (NRS 123) Genotype USA400
Community Associated Methicillin resistant *Staphylococcus aureus* (CA-MRSA), (NRS 384) Genotype USA300

TUBERCULOCIDAL ACTIVITY: **CaviCide AF** exhibits disinfectant efficacy against *Mycobacterium tuberculosis* BCG (*Mycobacterium bovis*) at 20°C when the treated surface is allowed to remain wet for **3 minutes**.

***VIRUCIDAL ACTIVITY – CaviCide AF**, when used on environmental, inanimate hard surfaces exhibits virucidal activity against: **Human Coronavirus** (ATCC VR-740), **SARS associated Coronavirus** (ZepitoMetrix), **Avian Influenza A virus strain H3N2** (ATCC VR-2072) and **Avian Influenza A virus strain H9N2** (Charles River Laboratories) (**Avian Influenza virus strains H3N2 and H9N2** for **2 minutes**; **Influenza A virus strain H1N1** (ATCC VR-1469), **Paramyxovirus** (Mumps) (ATCC VR-1438), **Rhinovirus type 39** (ATCC VR-340) and **Rotavirus** (University of Ottawa) for **3 minutes**; **HIV-1** (associated with AIDS) (Advanced Biotechnologies, Inc.) for **1 minute (60 seconds)**; and **Norovirus (Norwalk virus)**, **Feline Calicivirus** (University of Ottawa) and **Rabies Virus** (CDC) for **30 seconds**.

KILLS HIV-1 ON PRE-CLEANED ENVIRONMENTAL SURFACES/OBJECTS PREVIOUSLY SOILED WITH BLOOD/BODY FLUIDS in healthcare settings (Hospitals, Nursing Homes) or other settings in which there is an expected likelihood of soiling of inanimate surfaces/objects with blood or body fluids, and in which the surfaces/objects likely to be soiled with blood or body fluids can be associated with the potential for transmission of Human Immunodeficiency Virus Type 1 (HIV-1) (associated with AIDS).

SPECIAL INSTRUCTIONS FOR CLEANING AND DECONTAMINATION AGAINST HIV-1 ON SURFACES/OBJECTS SOILED WITH BLOOD/BODY FLUIDS:

PERSONAL PROTECTION: When handling items soiled with blood or body fluids use disposable latex gloves, gowns, masks, and eye coverings.

CLEANING PROCEDURES: Blood and other body fluids must be thoroughly cleaned from surfaces and objects before application of this product.

CONTACT TIME: Allow surface to remain wet for **1 minute**. **This contact time will not control other common types of viruses and bacteria.**

DISPOSAL OF INFECTIOUS MATERIALS: Blood and other body fluids must be autoclaved and disposed of according to local regulations for infectious waste disposal.

EFFICACY TESTS HAVE DEMONSTRATED THAT CAVICIDE AF IS AN EFFECTIVE BACTERICIDE, FUNGICIDE, AND VIRUCIDE IN THE PRESENCE OF ORGANIC SOIL (5% BLOOD SERUM).

MILDEWSTAT - To control mold and mildew on pre-cleaned, hard, non-porous surfaces spray surface to be treated making sure to wet completely. Let air dry. Repeat application at weekly intervals or when mildew growth appears.

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS CAUTION

KEEP OUT OF REACH OF CHILDREN. Causes moderate eye irritation. Avoid contact with eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet.

EPA Reg. No. 1839-225-46781 • EPA Est. No. 46781-MI-1

FIRST AID

Have the product container or label with you when calling a poison control center or doctor, or going for treatment.

If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

If swallowed: Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor. Do not give anything by mouth to an unconscious person.

If inhaled: Move person to fresh air. If person is not breathing, call 911 or an ambulance then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control center or doctor for further treatment advice.

NOTE TO PHYSICIAN

Probable mucosal damage may contraindicate the use of gastric lavage.

STORAGE AND DISPOSAL

DO NOT CONTAMINATE WATER, FOOD, OR FEED BY STORAGE OR DISPOSAL.

PESTICIDE STORAGE - Store in a dry place no lower in temperature than 50°F or higher than 120°F.

CONTAINER DISPOSAL - Nonrefillable container. Do not reuse or refill this container. Clean container promptly after emptying. Triple rinse as follows: Fill container ¼ full with water and recap. Agitate vigorously. Follow Pesticide Disposal instructions for rinsate disposal. Drain for 10 seconds after flow begins to drip. Repeat procedure two more times. Then offer for recycling or reconditioning. If not available, puncture and dispose in a sanitary landfill.

SPILL OR LEAK PROCEDURES - Small spills may be mopped up, flushed away with water or absorbed on some absorbent material and incinerated. Large spills should be contained, the material then moved into containers and disposed of by approved methods for hazardous wastes.

PESTICIDE DISPOSAL - Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

Manufactured For:

METREX™ RESEARCH
28210 Wick Road
Ramulus, Michigan 48174
1-800-841-1428

REORDER No. 13-8025

For Chemical Emergency ONLY (spill, leak, fire, exposure or accident), call CHEMTREC at 1-800-424-9300. For ALL other inquiries about this product, call Metrex Customer Service at 1-800-841-1428, Monday through Friday, between 6 a.m. and 4 p.m. Pacific Time.

For product or technical information, contact Metrex at 800-841-1428 or visit our website at www.metrex.com.

MADE IN USA

70-5094-1