

Seite 1: Layout aussen  
 Seite 2: Layout innen  
 Seite 3: Lack und Prägung



STATE OF HAWAII  
 Department of Agriculture  
**ACCEPTED**  
 LICENSE NO. 9618.119

PACK  
 1 2 3 4 5 6



advantage® II  
 large dog

81298378

108 x 34 x 167

advantage® II  
 large dog

Once-A-Month Topical Treatment  
 for Fleas and Lice  
 For Use Only on Dogs and Puppies  
 7 Weeks and Older and Weighing  
 21–55 lbs.

READ THE ENTIRE LABEL BEFORE  
 EACH USE

For the Prevention and Treatment of  
 Flea and Lice Infestations on Dogs

- Kills all flea life stages
- Prevents fleas on treated dogs  
 from infesting your home
- Kills fleas within 12 hours  
 of application
- Treats, prevents and controls  
 lice infestations

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS

**CAUTION:** Harmful if swallowed.  
 Causes moderate eye irritation.  
 Avoid contact with eyes or clothing.  
 Wash hands thoroughly with soap  
 and warm water after handling.

**HAZARDS TO DOMESTIC ANIMALS**  
**For external use only.** Do not use on  
 puppies under 7 weeks of age or  
 weighing less than 21 lbs. As with any  
 product, consult your veterinarian  
 before using this product on  
 debilitated, aged, pregnant or nursing  
 dogs. Individual sensitivities, while  
 rare, may occur after using ANY  
 pesticide product for dogs. If signs  
 persist, or become more severe,  
 consult a veterinarian immediately.

If your dog is on medication, consult  
 your veterinarian before using this or  
 any other product.

RESTRICTIONS:

- Use only on dogs 7 weeks and older  
 and weighing 21–55 lbs. Do not use  
 on other animals.
- Do not apply more than one (1) tube  
 per treatment.
- Do not have contact or allow children  
 to have contact with treated area  
 until completely dry.

Side Effects

Monitor your dog after application.  
 Side effects, although very rare, may  
 include signs of skin irritation such as  
 redness, scratching, or other signs of  
 discomfort. Gastrointestinal signs  
 such as vomiting or diarrhea have also  
 been reported. If these or other side  
 effects (such as lethargy) occur,  
 consult your veterinarian or call  
 1-800-422-9874.



For consumer questions call 1-800-255-6826  
 For medical emergencies call 1-800-422-9874

Manufactured for Bayer HealthCare LLC, Animal Health  
 Division, P.O. Box 390, Shawnee Mission, Kansas 66201, USA  
 Bayer, the Bayer Cross and Advantage are registered  
 trademarks of Bayer.

Made in Germany  
 ©2013 Bayer HealthCare LLC

EPA Est. No. 11556-DEU-1  
 EPA Reg. No. 11556-127

Net Contents: 6 Tubes  
 each 0.085 fl oz (2.5 mL)



1 2 3 4 5 6  
 PACK



1 2 3 4 5 6  
 PACK



large dog 21–55 lbs.

advantage® II  
 large dog




8

advantage® II

large dog

21–55 lbs.

Starts  
 working  
 through  
 contact



Once-A-Month Topical Treatment for  
 Fleas and Lice  
 For Use Only on Dogs and Puppies 7 Weeks  
 and Older and Weighing 21–55 lbs.

Active Ingredients % by Weight  
 Imidacloprid..... 9.10 %  
 Pyriproxyfen..... 0.46 %  
 Other Ingredients..... 90.44 %  
 Total..... 100.00 %



EPA Est. No. 11556-DEU-1  
 EPA Reg. No. 11556-127

KEEP OUT OF REACH OF CHILDREN  
**CAUTION**  
 See back panel for Precautionary  
 Statements. For Directions for Use,  
 Storage and Disposal, and First Aid  
 see package insert inside.



Lot No:



L&S GmbH & Co. KG In der Steele 13 · 40599 Düsseldorf Tel.: 0211774849-0 · Fax: -410	PM no.: 81298378 CR no.: 011521 Format: 108 x 34 x 167 mm Font size: 6 pt Drawing no.: 2367	For questions call Technology: phone: +49 431 3820... KVP Tanja Bernsdorf .....128 KVP Holger Schulz .....626 KVP Torsten Gagsch .....107	
	Colours: CMYK Pantone 10165 Pantone 877 gloss varnish matt varnish embossing Stanze	Operator ID: AG Version no.: 2 Date: 26.04.2013	Text + Presentation: phone: +49 2173 38... Christina Höfer.....4724 Rita Felicitas Reich.....4098 Yvonne Sabatini.....4082 Renate Steinacker.....3046
	Colours inside: Pantone 10165 Schwarz		
	THIS PROOF IS NOT A TRUE REPRESENTATION OF THE FINAL COLOURS THAT WILL BE PRINTED.		

 <b>GmbH &amp; Co. KG</b> In der Steele 13 · 40599 Düsseldorf Tel.: 0211/74849-0 · Fax: -410	
<b>PM no.:</b> 81298378 <b>CR no.:</b> 011521 <b>Format:</b> 108 x 34 x 167 mm <b>Font size:</b> 6 pt <b>Drawing no.:</b> 2367	<b>Colours:</b> <b>CMYK</b> Pantone 10165 Pantone 877 matt vernisch embossing Stanze
<b>Colours inside:</b> Pantone 10165 Schwarz	<b>Operator ID:</b> <b>AG</b> <b>Version no.:</b> <b>2</b> <b>Date:</b> <b>26.04.2013</b>
<b>For questions call</b> <b>Technology:</b> phone: +49 431 3820... KVP Tanja Bernsdorf .....128 KVP Holger Schulz .....626 KVP Torsten Göggsch .....107 <b>Text + Presentation:</b> phone: +49 2173 38... Christina Höfer .....4724 Rita Felicitas Reich .....4098 Yvonne Sabatini .....4082 Renate Steinacker .....3046	

**For more information  
 visit [www.petparents.com](http://www.petparents.com)**

**DIRECTIONS FOR USE**


It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

**HOW TO OPEN** 

1. Be sure tube is at bottom of pouch.
2. Using scissors, cut the pouch across the top and remove tube.

**HOW TO APPLY**

1. Remove one applicator tube from the package. See "HOW TO OPEN" section.
2. Hold applicator tube in an upright position facing away from you and your pet's face and eyes. Pull cap off tube.
3. Turn the cap around and place other end of cap back on tube.
4. Twist cap to break seal, then remove cap from tube.
5. The dog should be standing for easy application. Apply the entire contents of the tube evenly to three or four spots on the top of the back from the shoulder to the base of the tail. At each spot, part the hair until the skin is visible. Place the tip of the tube on the skin and gently squeeze to expel a portion of the solution on the skin. Do not apply an excessive amount of solution at any one spot that could cause some of the solution to run off the side of the dog. *Do not get this product in your dog's eyes or allow your dog to ingest this product.*



6. Discard empty tube as described in the Storage and Disposal section.
7. Under normal conditions the product is effective for a month. However, in case of severe flea infestation, retreatment may be necessary earlier than four weeks. Do not retreat more often than once every seven (7) days. After flea control is attained, return to a monthly retreatment schedule.

81356785

PACK  
1 2 3 4



advantage® II

108 x 34 x 120



STATE OF HAWAII  
Department of Agriculture

ACCEPTED

LICENSE NO.

9618.119

advantage® II  
large dog

Once-A-Month Topical Treatment for Fleas and Lice  
For Use Only on Dogs and Puppies 7 Weeks and Older and Weighing 21–55 lbs.

READ THE ENTIRE LABEL BEFORE EACH USE

For the Prevention and Treatment of Flea and Lice Infestations on Dogs

- Kills all flea life stages
- Prevents fleas on treated dogs from infesting your home
- Kills fleas within 12 hours of application
- Treats, prevents and controls lice infestations

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS

**CAUTION:** Harmful if swallowed. Causes moderate eye irritation. Avoid contact with eyes or clothing. Wash hands thoroughly with soap and warm water after handling.

HAZARDS TO DOMESTIC ANIMALS

**For external use only.** Do not use on puppies under 7 weeks of age or weighing less than 21 lbs. As with any product, consult your veterinarian before using this product on debilitated, aged, pregnant or nursing dogs. Individual sensitivities,

Manufactured for Bayer HealthCare LLC, Animal Health Division,  
P.O. Box 390, Shawnee Mission, Kansas 66201, USA  
Bayer, the Bayer Cross and Advantage are registered trademarks of Bayer.

Made in Germany  
©2013 Bayer HealthCare LLC

while rare, may occur after using ANY pesticide product for dogs. If signs persist, or become more severe, consult a veterinarian immediately. If your dog is on medication, consult your veterinarian before using this or any other product.

RESTRICTIONS:

- Use only on dogs 7 weeks and older and weighing 21–55 lbs. Do not use on other animals.
- Do not apply more than one (1) tube per treatment.
- Do not have contact or allow children to have contact with treated area until completely dry.

Side Effects

Monitor your dog after application. Side effects, although very rare, may include signs of skin irritation such as redness, scratching, or other signs of discomfort. Gastrointestinal signs such as vomiting or diarrhea have also been reported. If these or other side effects (such as lethargy) occur, consult your veterinarian or call 1-800-422-9874.



DO NOT USE ON CATS

Keep cats away from treated dogs for 24 hours. If applied to a cat or ingested by a cat, contact your veterinarian.

For consumer questions call 1-800-255-6826  
For medical emergencies call 1-800-422-9874

EPA Est. No. 11556-DEU-1  
EPA Reg. No. 11556-127

Net Contents: 4 Tubes  
each 0.085 fl oz (2.5 mL)

Bayer

1 2 3 4  
PACK



1 2 3 4  
PACK



large dog 21–55 lbs.

advantage® II



advantage® II

large dog  
21–55 lbs.

Starts working through contact

Once-A-Month Topical Treatment for Fleas and Lice  
For Use Only on Dogs and Puppies 7 Weeks and Older and Weighing 21–55 lbs.

Active Ingredients	% by Weight
Imidacloprid.....	9.10 %
Pyriproxyfen.....	0.46 %
Other Ingredients.....	90.44 %
Total .....	100.00 %

EPA Est. No. 11556-DEU-1  
EPA Reg. No. 11556-127

KEEP OUT OF REACH OF CHILDREN  
**CAUTION**  
See back panel for Precautionary Statements. For Directions for Use, Storage and Disposal, and First Aid see package insert inside.



7

24089 20283

3

## DIRECTIONS FOR USE


It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

### HOW TO OPEN

1. Be sure tube is at bottom of pouch.
2. Using scissors, cut the pouch across the top and remove tube.

### HOW TO APPLY

1. Remove one applicator tube from the package. See “HOW TO OPEN” section.
2. Hold applicator tube in an upright position facing away from you and your pet’s face and eyes. Pull cap off tube.
3. Turn the cap around and place other end of cap back on tube.
4. Twist cap to break seal, then remove cap from tube.
5. The dog should be standing for easy application. Apply the entire contents of the tube evenly to three or four spots on the top of the back from the shoulder to the base of the tail. At each spot, part the hair until the skin is visible. Place the tip of the tube on the skin and gently squeeze to expel a portion of the solution on the skin. Do not apply an excessive amount of solution at any one spot that could cause some of the solution to run off the side of the dog. *Do not get this product in your dog’s eyes or allow your dog to ingest this product.*



6. Discard empty tube as described in the Storage and Disposal section.
7. Under normal conditions the product is effective for a month. However, in case of severe flea infestation, retreatment may be necessary earlier than four weeks. Do not retreat more often than once every seven (7) days. After flea control is attained, return to a monthly retreatment schedule.

For more information  
visit [www.petparents.com](http://www.petparents.com)



# advantage® II

## large dog

**Once-A-Month Topical Treatment for Fleas and Lice**  
For Use Only on Dogs and Puppies 7 Weeks and Older and Weighing 21–55 lbs.

### READ THE ENTIRE LABEL BEFORE EACH USE.

#### For the Prevention and Treatment of Flea and Lice Infestations on Dogs

- Kills all flea life stages
- Prevents fleas on treated dogs from infesting your home
- One treatment prevents further flea infestations for a month
- Kills fleas on dogs within 12 hours of application
- Treats, prevents and controls lice infestations
- Waterproof
- Fragrance-free

#### Active Ingredients

% By Weight

Imidacloprid.....	9.10 %
Pyriproxyfen.....	0.46 %
<b>Other Ingredients.....</b>	<u>90.44 %</u>
<b>Total.....</b>	<b>100.00 %</b>

See back for First Aid and Precautionary Statements.

#### DIRECTIONS FOR USE


It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

#### HOW TO OPEN


1. Be sure tube is at bottom of pouch.
2. Using scissors, cut the pouch across the top and remove tube.

#### HOW TO APPLY

1. Remove one applicator tube from the package. See "HOW TO OPEN" section.
2. Hold applicator tube in an upright position facing away from you and your pet's face and eyes. Pull cap off tube.



3. Turn the cap around and place other end of cap back on tube.
4. Twist cap to break seal, then remove cap from tube.
5. The dog should be standing for easy application. Apply the entire contents of the tube evenly to three or four spots on the top of the back from the shoulder to the base of the tail. At each spot, part the hair until the skin is visible. Place the tip of the tube on the skin and gently squeeze to expel a portion of the solution on the skin. Do not apply an excessive amount of solution at any one spot that could cause some of the solution to run off the side of the dog. *Do not get this product in your dog's eyes or allow your dog to ingest this product.*



6. Discard empty tube as described in the Storage and Disposal section.

7. Under normal conditions the product is effective for a month. However, in case of severe flea infestation, retreatment may be necessary earlier than four weeks. Do not retreat more often than once every seven (7) days. After flea control is attained, return to a monthly retreatment schedule.

#### PRODUCT INFORMATION

The successive feeding activity of fleas on dogs frequently elicits a hypersensitivity skin disorder known as flea allergy dermatitis (FAD) or flea bite hypersensitivity. Treatment of dogs with Advantage II Large Dog rapidly kills fleas within 12 hours which may reduce the incidence of this condition.

Advantage II Large Dog kills existing fleas on dogs within 12 hours. Reinfesting fleas are killed within 2 hours with protection against further flea infestation lasting for four (4) weeks. Pre-existing pupae in the environment may continue to emerge for six (6) weeks or longer depending upon the climatic conditions.

Fleas, flea eggs, and flea larvae in the dog's environment are killed following contact with an Advantage II Large Dog treated dog. Advantage II Large Dog provides multi-stage flea control effectively breaking all flea life-cycle stages for quick and lasting control of flea populations.

Advantage II Large Dog kills adult fleas quickly, within 12 hours, inhibits the development of immature flea life stages and prevents them from reaching the biting adult stage.

Advantage II Large Dog kills chewing lice that infest dogs and puppies, and prevents further infestations.

Advantage II Large Dog is waterproof and remains effective following a shampoo treatment, swimming or after exposure to rain or sunlight.

Monthly treatments are required for optimal control and prevention of fleas.

# KEEP OUT OF REACH OF CHILDREN CAUTION

## PRECAUTIONARY STATEMENTS

### HAZARDS TO HUMANS

**CAUTION:** Harmful if swallowed. Causes moderate eye irritation. Avoid contact with eyes or clothing. Wash hands thoroughly with soap and warm water after handling.

FIRST AID	
<b>If Swallowed:</b>	<ul style="list-style-type: none"><li>• Call a poison control center or doctor immediately for treatment advice.</li><li>• Have person sip a glass of water if able to swallow.</li><li>• Do not induce vomiting unless told to do so by the poison control center or doctor.</li><li>• Do not give anything to an unconscious person.</li></ul>
<b>If In Eyes:</b>	<ul style="list-style-type: none"><li>• Hold eye open and rinse slowly and gently with water for 15–20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li><li>• Call a poison control center or doctor for treatment advice.</li></ul>
<b>If On Skin</b>	<ul style="list-style-type: none"><li>• Wash with plenty of soap and water.</li></ul>
HOT LINE NUMBER	
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For medical emergencies call 1-800-422-9874. For consumer questions call 1-800-255-6826.	
NOTE TO PHYSICIAN	
Treat the patient symptomatically.	

### HAZARDS TO DOMESTIC ANIMALS

**For external use only.** Do not use on puppies under 7 weeks of age or weighing less than 21 lbs. As with any product, consult your veterinarian before using this product on debilitated, aged, pregnant or nursing dogs. Individual sensitivities, while rare, may occur after using ANY pesticide product for dogs. If signs persist, or become more severe, consult a veterinarian immediately. If your dog is on medication, consult your veterinarian before using this or any other product.

**Side Effects:** Monitor your dog after application. Side effects, although very rare, may include signs of skin irritation such as redness, scratching, or other signs of discomfort. Gastrointestinal signs such as vomiting or diarrhea have also been reported. If these or other side effects (such as lethargy) occur, consult your veterinarian or call 1-800-422-9874.



For consumer questions call 1-800-255-6826.  
For medical emergencies call 1-800-422-9874.

### RESTRICTIONS:

- Use only on dogs 7 weeks and weighing 21–55 lbs. Do not use on other animals.
- Do not apply more than one (1) tube per treatment.
- Do not have contact or allow children to have contact with treated area until completely dry.

### STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

**Pesticide Storage:** Store in a cool, dry place inaccessible to children and pets.

**Pesticide Disposal and Container Handling:** Nonrefillable container.

**If empty:** Do not reuse or refill this container. Place in trash or offer for recycling if available.

**If partly filled:** Call your local solid waste agency or 1-800-422-9874 for disposal instructions. Never place unused product down any indoor or outdoor drain.

### LIMITED WARRANTY AND LIMITATION OF DAMAGES

Bayer HealthCare LLC, Animal Health Division warrants that this material conforms to the chemical description on the label. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BAYER MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY, INCLUDING ANY OTHER EXPRESS OR IMPLIED WARRANTY OF FITNESS OR MERCHANTABILITY, and no agent of Bayer is authorized to do so except in writing with a specific reference to this warranty. To the extent consistent with applicable law, any damages arising from a breach of this warranty shall be limited to direct damages and shall not include consequential commercial damages such as loss of profits or values, etc.

### Manufactured For

Bayer HealthCare LLC, Animal Health Division,  
P.O. Box 390, Shawnee Mission, Kansas 66201, USA  
Bayer, the Bayer Cross and Advantage are registered trademarks of Bayer.

© 2013 Bayer HealthCare LLC

Made in Germany

Advantage II is protected by the following U.S. patents:  
6,232,328 and 6,001,858

EPA Est. No. 11556-DEU-1  
EPA Reg. No. 11556-127

81298033

# Bayer

**THIS UNIT NOT FOR RETAIL SALE.**

**advantage<sup>®</sup> II**

**large dog**

**For external use only on dogs and puppies 7 weeks and older and weighing 21–55 lbs.**

See leaflet for complete information.

9.10 % Imidacloprid  
0.46 % Pyriproxyfen  
1 – 0.085 fl. oz. (2.5 mL) Tube  
EPA Reg. No. 11556-127



**DO NOT USE  
ON CATS**



81292361

9.1% Imidacloprid  
0.46% Pyriproxyfen  
0.084 fl.oz. (2.5 mL)  
EPA Reg. No. 11556-127

**advantage® II**  
**large dog**

Keep Out of Reach of  
Children

**CAUTION**

Read The Entire  
Label Before Use



8 1 2 7 0 7 7 5 ■

Lot No.: