

81356815



K9 advantix® II

108 x 34 x 120



STATE OF HAWAII
Department of Agriculture
ACCEPTED
LICENSE NO.

9618.122

K9 advantix® II medium dog

Topical Prevention and Treatment of Ticks, Fleas, Mosquitoes, Biting Flies and Lice for Monthly Use Only on Dogs and Puppies 7 Weeks of Age and Older and Weighing 11–20 lbs

READ THE ENTIRE LABEL BEFORE EACH USE
For the Prevention and Treatment of Ticks, Fleas, Mosquitoes, Biting Flies and Lice on Dogs

- Repels and kills ticks including Deer ticks (vector of Lyme disease), American dog ticks (vector of Rocky Mountain spotted fever), Brown dog ticks (vector of Ehrlichiosis) and Lone Star ticks (vector of Ehrlichiosis) for up to four weeks
- Repels and kills all life stages of fleas
- Prevents fleas on treated dogs from infesting your home
- Kills fleas on dogs within 12 hours and continues to prevent infestations for a month
- Prevents development of all flea stages for a month
- Larval flea stages in the dog's surroundings are killed following contact with a K9 Advantix® II treated dog
- Repels and kills mosquitoes for up to four weeks
- Repels and prevents blood-feeding by biting flies
- Kills chewing lice

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS

WARNING. Causes substantial but temporary eye injury. Do not get in eyes or on skin or clothing. Harmful if swallowed. Harmful if absorbed through skin. Avoid contact with skin. Wash thoroughly with soap and water

Manufactured for Bayer HealthCare LLC, Animal Health Division, P.O. Box 390, Shawnee Mission, Kansas 66201 USA
Bayer, the Bayer Cross and K9 Advantix are registered trademarks of Bayer.

Made in Germany
©2013 Bayer HealthCare LLC

after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Remove and wash contaminated clothing before reuse.

HAZARDS TO DOMESTIC ANIMALS

For external use on dogs only. Do not use on animals other than dogs. Do not use on puppies under seven (7) weeks of age. Do not get this product in dog's eyes or mouth. As with any product, consult your veterinarian before using this product on debilitated, aged, pregnant or nursing animals. Individual sensitivities, while rare, may occur after using ANY pesticide product for dogs. If signs persist, or become more severe, consult a veterinarian immediately. If your animal is on medication, consult your veterinarian before using this or any other product.

ENVIRONMENTAL HAZARDS

This pesticide is extremely toxic to aquatic organisms, including fish and invertebrates. Do not add directly to water. Do not contaminate water when disposing of product or packaging.

Side Effects

Monitor your dog after application. Side effects, although very rare, may include signs of skin irritation such as redness, scratching or other signs of discomfort. Gastrointestinal signs such as vomiting or diarrhea have also been reported. If these or other side effects (such as lethargy or agitation) occur, consult your veterinarian or call 1-800-422-9874.

DO NOT USE ON CATS – MAY BE FATAL.
Keep cats away from treated dogs for 24 hours. If applied to a cat or ingested by a cat, contact your veterinarian immediately.



For consumer questions call 1-800-255-6826
For medical emergencies call 1-800-422-9874



EPA Est. No. 11556-DEU-1
EPA Reg No. 11556-142

Net Contents: 4 Tubes
each 0.034 fl oz (1.0 mL)

Bayer



K9 advantix® II



K9 advantix® II



medium dog 11–20 lbs

**REPELS AND kills
ticks, fleas and mosquitoes**

Topical Prevention and Treatment of Fleas, Ticks, Mosquitoes, Biting Flies and Lice for Monthly Use ONLY on Dogs and Puppies 7 Weeks of Age and Older and Weighing 11–20 lbs

Active Ingredients	% by Weight
Imidacloprid.....	8.80%
Permethrin*	44.00%
Pyriproxyfen	0.44%
Other Ingredients	46.76%
Total	100.00%

* cis/trans ratio: max 55% (±) cis and min 45% (±) trans



EPA Est. No. 11556-DEU-1
EPA Reg No. 11556-142

KEEP OUT OF REACH OF CHILDREN
WARNING

See back panel for Precautionary Statements. For Directions for Use, Storage and Disposal, and First Aid, see package insert inside.



DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. Do not contaminate feed or food.


HOW TO OPEN

1. Be sure tube is at bottom of pouch.
2. Using scissors, cut the pouch across the top and remove tube.

HOW TO APPLY

1. Use only on dogs. **DO NOT USE ON CATS** or on other animals.
2. Remove applicator tube from the package. Use scissors and peel down foil.
3. Hold applicator tube in an upright position. Pull cap off tube.
4. Turn the cap around and place other end of cap back on tube.
5. Twist cap to break seal, then remove cap from tube.
6. The dog should be standing for easy application. The entire contents of the K9 Advantix® II Plus tube should be applied evenly to two to three spots on the top of the back from the shoulder to the base of the tail. At each spot, part the hair until the skin is visible. Place the tip of the tube on the skin and gently squeeze to expel a portion of the solution on the skin. Do not apply an excessive amount of solution at any one spot that could cause some of the solution to run off the side of the dog. Do not get this product in your dog's eyes, or allow your dog to ingest this product.

Do not get this product in your dog's eyes or mouth.



7. Discard empty tube as described in Storage and Disposal.
8. In cases of severe pest infestation, retreatment is allowable earlier than four (4) weeks. Do not retreat more often than once every seven (7) days. After pest control is attained, return to a monthly retreatment schedule.

For more information
visit www.petparents.com



K9 advantix® II

medium dog

Topical prevention and treatment of ticks, fleas, mosquitoes, biting flies and lice for monthly use only on dogs and puppies 7 weeks of age and older and weighing **11–20 lbs.**

READ THE ENTIRE LABEL BEFORE EACH USE

DO NOT USE ON CATS

For the Prevention and Treatment of Ticks, Fleas, Mosquitoes, Biting Flies and Lice on Dogs

- Repels and kills ticks including Deer ticks (vector of Lyme disease), American dog ticks (vector of Rocky Mountain spotted fever), Brown dog ticks (vector of Ehrlichiosis), and Lone Star ticks (vector of Ehrlichiosis) for up to four weeks
- Repels and kills all life stages of fleas
- Prevents fleas on treated dogs from infesting your home
- Kills fleas on dogs within 12 hours and continues to prevent infestations for a month
- Prevents development of fleas, flea eggs, pupae and larvae for a month
- Larval flea stages in the dog's surroundings are killed following contact with a K9 Advantix® II treated dog
- Repels and kills mosquitoes for up to four weeks
- Repels and prevents blood-feeding by biting flies
- Waterproof
- 5-way protection against ticks, fleas, mosquitoes, biting flies and lice
- Kills chewing lice

Active Ingredients % By Weight

imidacloprid.....	8.80%
Permethrin*	44.00%
Pyriproxyfen.....	0.44%
Other Ingredients	46.76%
Total	100.00%

* cis/trans ratio: max 55% (±) cis and min 45% (±) trans

KEEP OUT OF REACH OF CHILDREN

WARNING

See back for First Aid and Precautionary Statements.

DIRECTIONS FOR USE


It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. Do not contaminate feed or food.

HOW TO OPEN


1. Be sure tube is at bottom of pouch.
2. Using scissors, cut the pouch across the top and remove tube.

HOW TO APPLY

1. Use only on dogs.
DO NOT USE ON CATS or on other animals.
2. Remove applicator tube from the package. Use scissors and peel down foil.
3. Hold applicator tube in an upright position. Pull cap off tube.



4. Turn the cap around and place other end of cap back on tube.
5. Twist cap to break seal, then remove cap from tube.
6. The dog should be standing for easy application. The entire contents of the K9 Advantix® II tube should be applied evenly to two to three spots on the top of the back from the shoulder to the base of the tail. At each spot, part the hair until the skin is visible. Place the tip of the tube on the skin and gently squeeze to expel a portion of the solution on the skin. Do not apply an excessive amount of solution at any one spot that could cause some of the solution to run off the side of the dog.
Do not get this product in your dog's eyes, or allow your dog to ingest this product.



7. Discard empty tube as described in Storage and Disposal.
8. In cases of severe pest infestation, retreatment is allowable earlier than four (4) weeks. Do not retreat more often than once every seven (7) days. After pest control is attained, return to a monthly retreatment schedule.

PRODUCT INFORMATION

The successive feeding activity of fleas on dogs may elicit a hypersensitivity skin disorder known as flea allergy dermatitis (FAD). Treatment of dogs with K9 Advantix® II rapidly kills fleas and may reduce the incidence of this condition.

K9 Advantix® II kills existing fleas on dogs within 12 hours. Reinfesting fleas are killed within 2 hours with protection against further flea infestation lasting at least four (4) weeks. Pre-existing pupae in the environment may continue to emerge for six (6) weeks or longer depending upon the climatic conditions.

Larval flea stages in the dog's surroundings are killed following contact with a K9 Advantix® II treated dog. K9 Advantix® II is waterproof and remains effective following a shampoo treatment, swimming or after exposure to rain or sunlight.

Monthly treatments are required for optimal control and prevention of fleas and ticks.

KEEP OUT OF REACH OF CHILDREN WARNING

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS

WARNING. Causes substantial but temporary eye injury. Do not get in eyes or on skin or clothing. Harmful if swallowed. Harmful if absorbed through skin. Avoid contact with skin. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Remove and wash contaminated clothing before reuse.

HAZARDS TO DOMESTIC ANIMALS

For external use on dogs only. Do not use on animals other than dogs. Do not use on puppies under seven weeks of age.

Do not get this product in dog's eyes or mouth. As with any product, consult your veterinarian before using this product on debilitated, aged, pregnant or nursing animals. Individual sensitivities, while rare, may occur after using ANY pesticide product for dogs. If signs persist, or become more severe, consult a veterinarian immediately.

If your animal is on medication, consult your veterinarian before using this or any other product.



DO NOT USE ON CATS – MAY BE FATAL.

Keep cats away from treated dogs for 24 hours. If applied to a cat or ingested by a cat, contact your veterinarian immediately.

Side Effects: Monitor your dog after application. Side effects, although very rare, may include signs of skin irritation such as redness, scratching, or other signs of discomfort. Gastrointestinal signs such as vomiting or diarrhea have also been reported. If these or other side effects (such as lethargy or agitation) occur, consult your veterinarian or call 1-800-422-9874.

ENVIRONMENTAL HAZARDS

This pesticide is extremely toxic to aquatic organisms, including fish and invertebrates. Do not add directly to water. Do not contaminate water when disposing of product or packaging.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

Pesticide Storage: Store in a cool, dry place. Protect from freezing.

Pesticide Disposal and Container Handling: Nonrefillable container.

If empty: Do not reuse this container.

Place in trash or offer for recycling, if available.

If partly filled: Call your local solid waste agency or 1-800-422-9874 for disposal instructions.

Never place unused product down any indoor or outdoor drain.

LIMITED WARRANTY AND LIMITATION OF DAMAGES

Bayer HealthCare LLC, Animal Health Division warrants that this material conforms to the chemical description on the label. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BAYER MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY, INCLUDING ANY OTHER EXPRESS OR IMPLIED WARRANTY OF FITNESS OR MERCHANTABILITY, and no agent of Bayer is authorized to do so except in writing with a specific reference to this warranty. To the extent consistent with applicable law, any damages arising from a breach of this warranty shall be limited to direct damages and shall not include consequential commercial damages such as loss of profits or values, etc.

FIRST AID

If In Eyes:	<ul style="list-style-type: none">• Hold eye open and rinse slowly and gently with water for 15–20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.• Call a poison control center or doctor for treatment advice.
If Swallowed:	<ul style="list-style-type: none">• Call a poison control center or doctor immediately for treatment advice.• Have person sip a glass of water if able to swallow.• Do not induce vomiting unless told to do so by the poison control center or doctor.• Do not give anything to an unconscious person.
If On Skin or Clothing:	<ul style="list-style-type: none">• Take off contaminated clothing.• Rinse skin immediately with plenty of water for 15–20 minutes.• Call a poison control center or doctor for treatment advice.

HOT LINE NUMBER

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-422-9874 for emergency medical treatment information. For consumer questions call 1-800-255-6826.

NOTE TO PHYSICIAN

Treat the patient symptomatically.

Manufactured For

Bayer HealthCare LLC, Animal Health Division,
P.O. Box 390, Shawnee Mission,
Kansas 66201 USA

Bayer, the Bayer Cross, and K9 Advantix
are registered trademarks of Bayer

© 2013 Bayer HealthCare LLC

Made in Germany

K9 Advantix® is protected by the following

U.S. patents: 6,232,328, 7,728,011

EPA Est. No. 11556-DEU-1

EPA Reg. No. 11556-142

81294100

Bayer

THIS UNIT NOT FOR RETAIL SALE.

K9 advantix[®] II

medium dog

For external use only on dogs and puppies 7 weeks and older and weighing 11–20 lbs.

See leaflet for complete product information.

8.80% Imidacloprid

44.0% Permethrin

0.44% Pyriproxyfen

1 - 0.034 fl. oz. (1.0 mL) Tube

EPA Reg. No. 11556-142



**DO NOT USE
ON CATS**



81291853

0.034 fl.oz.

8.8% imidacloprid
44.0% permethrin
0.44% pyriproxyfen

K9 advantix® II
medium dog

EPA Reg. No. 11556-142

Keep out of reach of children

WARNING

DO NOT USE ON CATS

Read The Entire
Label Before Use



8 1 2 7 0 8 0 5 ■

Lot No.: