



advantage II

advantage II large cat

Once-A-Month Topical Flea Prevention and Treatment for Cats
For Use ONLY on Cats 8 Weeks and Older and Weighing Over 9 lbs.

READ THE ENTIRE LABEL BEFORE EACH USE

For the Prevention and Treatment of Flea Infestations

- 3-way flea protection kills adults, larvae and eggs
- Controls existing fleas and flea eggs and prevents future flea infestations
- Kills fleas on cats within 12 hours of application

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS

CAUTION: Harmful if swallowed. Causes moderate eye irritation. Avoid contact with eyes or clothing. Wash hands thoroughly with soap and warm water after handling. Keep out of reach of children. Do not contaminate feed or food.

HAZARDS TO DOMESTIC ANIMALS

For external use only. Do not apply to cats or kittens under 8 weeks of age or weighing less

Manufactured for Bayer HealthCare LLC, Animal Health Division,
P.O. Box 390, Shawnee Mission, Kansas 66201, USA
Bayer, the Bayer Cross and Advantage are registered trademarks of Bayer.
Made in Germany
©2013 Bayer HealthCare LLC

EPA Est. No. 11556-DEU-1
EPA Reg. No. 11556-152

Net Contents: 4 Tubes
each 0.027 fl oz (0.8 mL)



than 9 lbs. As with any product, consult your veterinarian before using this product on debilitated, aged, pregnant or nursing cats. Individual sensitivities, while rare, may occur after using ANY pesticide product for cats. If signs persist, or become more severe, consult a veterinarian immediately. If your cat is on medication, consult your veterinarian before using this or any other product.

RESTRICTIONS:

- Use only on cats or kittens 8 weeks and older. Do not apply to cats or kittens weighing less than 9 lbs. Do not use on other animals.
- Do not apply more than one (1) tube per treatment, even for larger cats.
- Do not have contact or allow children to have contact with treated area until completely dry.

Side Effects

Monitor your cat after application. Side effects, although very rare, may include signs of skin irritation such as redness, scratching, or other signs of discomfort. Gastrointestinal signs such as hypersalivation, vomiting or diarrhea have also been reported. If these or other side effects (such as lethargy) occur, consult your veterinarian or call 1-800-422-9874.


For consumer questions call 1-800-255-6826
For medical emergencies call 1-800-422-9874

108 x 34 x 120



large cat over 9 lbs.

advantage II



STATE OF HAWAII
Department of Agriculture
ACCEPTED
LICENSE NO. **9618.127**



advantage II

large cat over 9 lbs.

Starts working through contact

Once-A-Month Topical Flea Prevention and Treatment for Cats
For Use ONLY on Cats 8 Weeks and Older and Weighing Over 9 lbs.

Active Ingredients	% by Weight
Imidacloprid.....	9.10 %
Pyriproxyfen.....	0.46 %
Other Ingredients.....	90.44 %
Total	100.00 %

EPA Est. No. 11556-DEU-1
EPA Reg. No. 11556-152

**KEEP OUT OF REACH OF CHILDREN
CAUTION**
See back panel for Precautionary Statements. For Directions for Use, Storage and Disposal, and First Aid see package insert inside.



DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

HOW TO OPEN

1. Be sure tube is at bottom of pouch.
2. Using scissors, cut the pouch across the top and remove tube.

HOW TO APPLY

1. Remove one applicator tube from the package. See “HOW TO OPEN” section.
2. Hold applicator tube in an upright position facing away from you and your pet’s face and eyes. Pull cap off tube.
3. Turn the cap around and place other end of cap back on tube.
4. Twist cap to break seal, then remove cap from tube.
5. Part the hair on the neck at the base of the skull until the skin is visible. Place the tip of the tube on the skin and squeeze the tube to expel the entire contents directly on the skin. *Do not get this product in your cat’s eyes, or allow your cat to ingest this product. The product is bitter tasting and salivation may occur for a short time if the cat licks the product immediately after treatment. Treatment at the base of the skull will minimize the opportunity for the cat to lick the product. Do not allow the product to run off.*



6. Discard empty tube as described in the Storage and Disposal section.
7. Under normal conditions the product is effective for a month. However, in case of severe flea infestation, retreatment may be necessary earlier than four (4) weeks. Do not retreat more often than once every seven (7) days. After flea control is attained, return to a monthly retreatment schedule.

For more information
visit www.petparents.com



advantage® II

large cat

Once-A-Month Topical Flea Prevention and Treatment for Cats
For Use **ONLY** on Cats 8 Weeks and Older and Weighing Over 9 lbs.

READ THE ENTIRE LABEL BEFORE EACH USE.

For the Prevention and Treatment of Flea Infestations

- Kills all flea life stages
- Controls existing flea infestations on your cat and prevents further infestations
- Kills fleas within 12 hours of application
- Convenient, easy-to-apply topical solution
- Fragrance-free
- Waterproof

Active Ingredients	% By Weight
Imidacloprid.....	9.10 %
Pyriproxyfen.....	0.46 %
Other Ingredients	<u>90.44 %</u>
Total	100.00 %

See back for First Aid and Precautionary Statements.

DIRECTIONS FOR USE


It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

HOW TO OPEN

1. Be sure tube is at bottom of pouch.
2. Using scissors, cut the pouch across the top and remove tube.

HOW TO APPLY

1. Remove one applicator tube from the package. See "HOW TO OPEN" section.
2. Hold applicator tube in an upright position facing away from you and your pet's face and eyes. Pull cap off tube.



3. Turn the cap around and place other end of cap back on tube.
4. Twist cap to break seal, then remove cap from tube.
5. Part the hair on the neck at the base of the skull until the skin is visible. Place the tip of the tube on the skin and squeeze the tube to expel the entire contents directly on the skin. *Do not get this product in your cat's eyes, or allow your cat to ingest this product. The product is bitter tasting and salivation may occur for a short time if the cat licks the product immediately after treatment.*

Treatment at the base of the skull will minimize the opportunity for the cat to lick the product. Do not allow the product to run off.



6. Discard empty tube as described in Storage and Disposal.
7. Under normal conditions this product is effective for a month. However, in case of severe flea infestation, retreatment may be necessary earlier than four (4) weeks. Do not retreat more often than once every seven (7) days. After flea control is attained, return to a monthly retreatment schedule.

PRODUCT INFORMATION

The successive feeding activity of fleas on cats frequently elicits a hypersensitivity skin disorder known as flea allergy dermatitis (FAD) or flea bite hypersensitivity. Treatment of cats with Advantage II Large Cat kills fleas and may reduce the incidence of this condition.

Advantage II Large Cat kills the existing fleas on cats within 12 hours. Reinfesting fleas are killed within 2 hours with protection against further flea infestation lasting for up to four (4) weeks. Pre-existing pupae in the environment may continue to emerge for six (6) weeks or longer depending upon the climatic conditions.

Fleas, eggs and larvae in the cat's surroundings are killed following contact with an Advantage II Large Cat treated cat. Advantage II Large Cat provides multi-stage flea control effectively breaking all flea life-cycle stages for lasting control of flea populations.

Advantage II Large Cat kills adult fleas quickly, within 12 hours, inhibits the development of immature flea life stages and prevents them from reaching the biting adult stage.

Advantage II Large Cat is waterproof and remains effective following a shampoo treatment or after exposure to rain or sunlight.

Apply monthly treatments for optimal control and prevention of fleas.

KEEP OUT OF REACH OF CHILDREN CAUTION

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS

Caution: Harmful if swallowed. Causes moderate eye irritation. Avoid contact with eyes or clothing. Wash hands thoroughly with soap and warm water after handling. Keep out of reach of children. Do not contaminate feed or food.

FIRST AID	
If Swallowed:	<ul style="list-style-type: none">• Call a poison control center or doctor immediately for treatment advice.• Have person sip a glass of water if able to swallow.• Do not induce vomiting unless told to do so by the poison control center or doctor.• Do not give anything to an unconscious person.
If In Eyes:	<ul style="list-style-type: none">• Hold eye open and rinse slowly and gently with water for 15–20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.• Call a poison control center or doctor for treatment advice.
If On Skin:	<ul style="list-style-type: none">• Wash with plenty of soap and water.
HOT LINE NUMBER	
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For medical emergencies call 1-800-422-9874. For customer questions call 1-800-255-6826.	
NOTE TO PHYSICIAN	
Treat the patient symptomatically.	

HAZARDS TO DOMESTIC ANIMALS

For external use only. Do not use on cats or kittens under 8 weeks of age or weighing less than 9 lbs. As with any product, consult your veterinarian before using this product on debilitated, aged, pregnant or nursing cats. Individual sensitivities, while rare, may occur after using ANY pesticide product for cats. If signs persist, or become more severe, consult a veterinarian immediately. If your cat is on medication, consult your veterinarian before using this or any other product.

Side Effects

Monitor your cat after application. Side effects, although very rare, may include signs of skin irritation such as redness, scratching, or other signs of discomfort. Gastrointestinal signs such as hypersalivation, vomiting or diarrhea have also been reported. If these or other side effects (such as lethargy) occur, consult your veterinarian or call 1-800-422-9874.

For consumer questions call 1-800-255-6826.

For medical emergencies call 1-800-422-9874.

RESTRICTIONS:

- Use only on cats or kittens 8 weeks and older. Do not apply to cats or kittens weighing less than 9 lbs. Do not use on other animals.
- Do not apply more than one (1) tube per treatment.
- Do not have contact or allow children to have contact with treated area until completely dry.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

Pesticide Storage: Store in a cool, dry place inaccessible to children and pets.

Pesticide Disposal and Container Handling: Nonrefillable container.

If empty: Do not reuse or refill this container. Place in trash or offer for recycling if available.

If partly filled: Call your local solid waste agency or 1-800-422-9874 for disposal instructions. Never place unused product down any indoor or outdoor drain.

LIMITED WARRANTY AND LIMITATION OF DAMAGES

Bayer HealthCare LLC, Animal Health Division warrants that this material conforms to the chemical description on the label. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BAYER MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY, INCLUDING ANY OTHER EXPRESS OR IMPLIED WARRANTY OF FITNESS OR MERCHANTABILITY, and no agent of Bayer is authorized to do so except in writing with a specific reference to this warranty. Any damages arising from a breach of this warranty shall be limited to direct damages and shall not include consequential commercial damages such as loss of profits or values, etc.

Manufactured For

Bayer HealthCare LLC, Animal Health Division,
P.O. Box 390, Shawnee Mission, Kansas 66201, USA
Bayer, the Bayer Cross and Advantage are registered trademarks of Bayer.

© 2013 Bayer HealthCare LLC

Made in Germany

Advantage II is protected by the following U.S. patents:
6,232,328 and 6,001,858

EPA Est. No. 11556-DEU-1

EPA Reg. No. 11556-152

81297878

Bayer

THIS UNIT NOT FOR RETAIL SALE.

advantage[®] II

81292337

large cat

For external use only on cats and kittens 8 weeks and older and weighing over 9 lbs.

See leaflet for complete information.

9.10 % Imidacloprid

0.46 % Pyriproxyfen

1 – 0.027 fl. oz. (0.8 mL) Tube

EPA Reg. No. 11556-152



Lot:

1 x auf 125 mm (Folienbreite)

4 x auf 420 mm (Abschnittslänge 105 mm)

9.1% Imidacloprid
0.46% Pyriproxyfen

advantage® II
large cat

EPA Reg. No. 11556-152
Keep Out of Reach of
Children

CAUTION

Read The Entire
Label Before Use



81270740 ■

Lot No.:

0.027 fl.oz.