



advantage[®] 18
Topical Solution

8 Weeks and Older
and Over 9 lbs.

6 Pack

105x25x165

79619707

**Over
9 lbs.**



B
A
Y
E
R

Topical Solution
advantage[®] 18

advantage[®] 18
Topical Solution

advantage[®] 18
Topical Solution

**Once-a-Month Topical Flea Treatment
for Cats and Kittens 8 Weeks and Older
and Over 9 lbs.**

- Kills fleas on cats within 12 hours
- Kills fleas before they lay eggs
- Convenient, easy to apply

Active Ingredient	% By Weight
Imidacloprid.....	9.1 %
Other Ingredients	90.9 %
Total	100.0 %



**KEEP OUT OF REACH
OF CHILDREN
WARNING**

See back panel for First Aid.
For Directions for Use, and Storage and
Disposal, see supplemental labeling inside.



LICENSED

PERIOD **2009-2011** LIC. NO.



6 Pack



Department of Agriculture
STATE OF HAWAII

LICENSED

PERIOD **2015-2017** LIC. NO.

9618.91



Department of Agriculture
STATE OF HAWAII

LICENSED

PERIOD **2012-2014** LIC. NO.

advantage[®] 18
Topical Solution

**Once-a-Month Topical Flea Treatment
for Cats and Kittens 8 Weeks and Older
and Over 9 lbs.**

READ THE ENTIRE LABEL BEFORE EACH USE

**For the Prevention and Treatment of
Flea Infestations on Cats**

- Kills fleas on cats within 12 hours
- Kills fleas before they lay eggs

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS:

Causes eye irritation. Harmful if swallowed.
Do not get in eyes or on clothing. Avoid contact
with skin. Wash hands thoroughly with soap and
warm water after handling. Keep out of reach of
children. Do not contaminate feed or food.

HAZARDS TO DOMESTIC ANIMALS

For external use only. Do not use on kittens under
eight weeks of age. As with any product, consult
your veterinarian before using this product on
debilitated, aged, pregnant or nursing animals.
Individual sensitivities, while rare, may occur
after using ANY pesticide product for pets. If
signs persist, or become more severe, consult a
veterinarian immediately. If your animal is on
medication, consult your veterinarian before
using this or any other product.

For consumer questions call 1-800-255-6826.
For medical emergencies call 1-800-422-9874.

FIRST AID

Have the product container or label with you
when calling a poison control center or doctor,
or going for treatment.

If in eyes:

Hold eye open and rinse slowly and gently
with water for 15–20 minutes.
Remove contact lenses, if present, after the
first 5 minutes, then continue rinsing eye.
Call a poison control center or doctor for
treatment advice.

If swallowed:

Call poison control center or doctor immediately
for treatment advice. Have person sip a glass of
water if able to swallow. Do not induce vomiting
unless told to do so by the poison control center
or doctor. Do not give anything by mouth to an
unconscious person.

If on skin or clothing:

Take off contaminated clothing. Rinse skin
immediately with plenty of water for 15–20
minutes. Call a poison control center or doctor for
treatment advice.

To Physician: Treat the patient symptomatically.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product
in a manner inconsistent with its labeling.

HOW TO APPLY

1. Use only on cats. Do not use on other animals.
2. Remove one applicator tube from the package.
3. Hold applicator tube in an upright position.
Pull cap off tube.
4. Turn the cap around and place other end of cap
back on tube.
5. Twist cap to break seal, then remove cap from
tube.



6. Part the hair on the neck at the base of the
skull until the skin is visible. Place the tip of the
tube on the skin and squeeze the tube to *expel
the entire contents of the tube directly on the
skin.*

*Do not get this product in your pet's eyes or
mouth.* The product is bitter tasting and
salivation may occur for a short time if the cat
licks the product immediately after treatment.
Treatment at the base of the skull will minimize
the opportunity for the cat to lick the product.

7. Discard empty tube as described in Storage
and Disposal.
8. In case of severe flea infestation, retreatment
may be necessary earlier than four weeks.
Do not retreat more often than once every
seven days. After flea control is attained, return
to a monthly retreatment schedule.

Manufactured for Bayer HealthCare LLC,
Animal Health Division, P.O. Box 390,
Shawnee Mission, KS, 66201, USA

Bayer, the Bayer Cross and Advantage
are registered trademarks of Bayer.

Made in Germany

© 2009 Bayer HealthCare LLC

01791692-053799, R.7
EPA Est. No. 11556-DEU-1
EPA Reg. No. 11556-118



Lot No.:





advantage[®] 18

Topical Solution

Once-A-Month Topical Flea Treatment for Cats and Kittens 8 Weeks and Older and Over 9 lbs.

READ THE ENTIRE LABEL BEFORE EACH USE.

For the Prevention and Treatment of Flea Infestations on Cats

- One treatment prevents further flea infestation for up to four weeks
- Kills fleas on cats within 12 hours
- Kills fleas before they lay eggs
- Waterproof
- Fragrance-free

Active Ingredient

Imidacloprid..... 9.1%

Other Ingredients..... 90.9%

Total..... 100.0%

KEEP OUT OF REACH OF CHILDREN WARNING

See below for First Aid and Precautionary Statements.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS: Causes eye irritation.

Harmful if swallowed. Do not get in eyes or on clothing. Avoid contact with skin. Wash hands thoroughly with soap and warm water after handling. Keep out of reach of children. Do not contaminate feed or food.

HAZARDS TO DOMESTIC ANIMALS

For external use only. Do not use on kittens under eight weeks of age. As with any product, consult your veterinarian before using this product on debilitated, aged, pregnant or nursing animals. Individual sensitivities, while rare, may occur after using ANY pesticide product for pets.

If signs persist, or become more severe, consult a veterinarian immediately. If your animal is on medication, consult your veterinarian before using this or any other product.

For consumer questions call 1-800-255-6826.

For medical emergencies call 1-800-422-9874.

FIRST AID

Have the product container or label with you when calling a poison control center or doctor, or going for treatment.

If in eyes: Hold eye open and rinse slowly and gently with water for 15–20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.

Call a poison control center or doctor for treatment advice.

If swallowed: Call poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor. Do not give anything by mouth to an unconscious person.

If on skin or clothing: Take off contaminated clothing.

Rinse skin immediately with plenty of water for 15–20 minutes.

Call a poison control center or doctor for treatment advice.


To Physician: Treat the patient symptomatically.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

HOW TO APPLY

1. Use only on cats. Do not use on other animals.
2. Remove one applicator tube from the package.
3. Hold applicator tube in an upright position. Pull cap off tube.



4. Turn the cap around and place other end of cap back on tube.
5. Twist cap to break seal, then remove cap from tube.



6. Part the hair on the neck at the base of the skull until the skin is visible. Place the tip of the tube on the skin and squeeze the tube to expel the entire contents of the tube directly on the skin. Do not get this product in your pet's eyes or mouth. The product is bitter tasting and salivation may occur for a short time if the cat licks the product immediately after treatment. Treatment at the base of the skull will minimize the opportunity for the cat to lick the product.
7. Discard empty tube as described in Storage and Disposal.
8. In case of severe flea infestation, retreatment may be necessary earlier than four weeks. Do not retreat more often than once every seven days. After flea control is attained, return to a monthly retreatment schedule.

GENERAL INFORMATION

The successive feeding activity of fleas on pets may elicit a hypersensitivity skin disorder known as flea allergy dermatitis (FAD). Treatment of cats with Advantage[®] kills fleas and may reduce the incidence of this condition.

Advantage kills fleas on cats within 12 hours. Reinfesting fleas are killed within 2 hours with protection against further flea infestation lasting for up to four (4) weeks. Pre-existing pupae in the environment may continue to emerge for six (6) weeks or longer depending upon the climatic conditions. Larval flea stages in the pet's surroundings are killed following contact with an Advantage treated pet. Advantage is waterproof and remains effective following a shampoo treatment, swimming or after exposure to rain or sunlight. Monthly treatments are required for optimal control and prevention of fleas.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

Storage: Store in a cool, dry place.

Disposal: Nonrefillable container.

If empty: Do not reuse or refill this container. Place in trash or offer for recycling if available.

If partly filled: Call your local solid waste agency or 1-800-422-9874 for disposal instructions. Never place unused product down any indoor or outdoor drain.

LIMITED WARRANTY AND LIMITATION OF DAMAGES

Bayer HealthCare LLC, Animal Health Division warrants that this material conforms to the chemical description on the label. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BAYER MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY, INCLUDING ANY OTHER EXPRESS OR IMPLIED WARRANTY OF FITNESS OR MERCHANTABILITY, and no agent of Bayer is authorized to do so except in writing with a specific reference to this warranty. Any damages arising from a breach of this warranty shall be limited to direct damages and shall not include consequential commercial damages such as loss of profits or values, etc.

Manufactured For

Bayer HealthCare LLC, Animal Health Division,
P.O. Box 390, Shawnee Mission, Kansas 66201 USA
Bayer, the Bayer Cross and Advantage are registered trademarks of Bayer

© 2009 Bayer HealthCare LLC

Made in Germany

Advantage is protected by the following U.S. patents:
6,232,328 and 6,001,858

79619693

00837877-032799/01791692-053799, R.4

EPA Est. No. 11556-DEU-1

EPA Reg. No. 11556-118



Bayer HealthCare

9.1% Imidacloprid
0.027 fl.oz. (0.8 mL)
EPA Reg. No. 11556-118

 **advantage** 18

Keep Out of Reach of
Children

WARNING
Read The Entire
Label Before Use



79618557 ■

Lot No.: