


2003-2005 Key

PROD. #2429  
1097C

BEFORE USING THIS PRODUCT,  
PLEASE READ THIS ENTIRE LABEL.

**SANITIZER**



**CHLORINATED DISINFECTANT AND GERMICIDE**

<b>ACTIVE INGREDIENTS:</b>	
Sodium Hypochlorite .....	5.25%
<b>INERT INGREDIENTS:</b> .....	94.75%
<b>TOTAL</b> .....	<b>100.00%</b>

**KEEP OUT OF REACH OF CHILDREN**

**DANGER**

**STATEMENT OF PRACTICAL TREATMENT (FIRST AID)**

**IF CONTACT WITH EYES OCCURS,** flush with water for at least 15 minutes. Get prompt medical attention.

**IF CONTACT WITH SKIN OCCURS,** wash with plenty of soap and water.

**IF SWALLOWED,** drink large amounts of water. DO NOT induce vomiting. Call a physician or poison control center immediately.

**FOR INDUSTRIAL AND INSTITUTIONAL USE ONLY**

E.P.A. Reg. No. 1270-20004

E.P.A. Est. No. 1270-GA-1

**PRECAUTIONARY STATEMENTS  
HAZARDS TO HUMANS AND DOMESTIC ANIMALS  
DANGER**

**DANGER:** Corrosive, may cause severe skin and eye irritation or chemical burns to broken skin. Causes eye damage. Wear safety glasses or goggles and rubber gloves when handling this product. Wash after handling. Avoid breathing vapors. Vacate poorly-ventilated areas as soon as possible. Do not return until strong odors have dissipated.

**ENVIRONMENTAL HAZARDS**

This pesticide is toxic to fish and aquatic organisms. Do not discharge effluent containing this product into lakes, streams, ponds, estuaries, oceans or other waters unless in accordance with the requirements of a National Discharge Elimination System (NPDES) permit and the permitting authority has been notified in writing prior to discharge. Do not discharge effluent containing this product to sewer systems without previously notifying the local sewage treatment plant authority. For guidance, contact your State Water Board or Regional Office of the EPA.

**PHYSICAL OR CHEMICAL HAZARDS**

**STRONG OXIDIZING AGENT:** Mix only with water, according to label directions. Mixing this product with chemicals (e.g. ammonia, acids, detergents, etc.) or organic matter (e.g. urine, feces, etc.) will release chlorine gas, which is irritating to eyes, lungs and mucous membranes.

**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

**NOTE:** This product degrades with age. Use a chlorine test kit and increase dosage, as necessary, to obtain the required level of available chlorine.

**STORAGE AND DISPOSAL**

**STORAGE:** Store this product in a cool, dry area, away from direct sunlight and heat to avoid deterioration. In case of spill, flood areas with large quantities of water.

**PESTICIDE DISPOSAL:** Product or rinsates that cannot be used should be diluted with water before disposal in a sanitary sewer. Do not contaminate food or feed by storage, disposal or cleaning of equipment.

**CONTAINER DISPOSAL:** Triple rinse (or equivalent), then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill or by other procedures approved by state and local authorities.

**FOOD EGG SANITIZATION IN POULTRY AND EGG PROCESSING PLANTS**

Thoroughly clean all eggs. Thoroughly mix 5 oz. of this product with 10 gallons of warm water to produce a 200 ppm available chlorine solution. The sanitizer temperature should not exceed 130°F. Spray the warm sanitizer so that the eggs are thoroughly wetted. Allow the eggs to thoroughly dry before casing or breaking. Do not apply a potable water rinse. The solution should not be reused to sanitize eggs.

**SANITIZATION OF SURFACES SUCH AS PROCESSING EQUIPMENT, DAIRY EQUIPMENT, FOOD UTENSILS, DISHES, SILVERWARE, GLASSES, SINK TOPS, COUNTER TOPS, REFRIGERATED STORAGE AND DISPLAY EQUIPMENT AND OTHER NON-POROUS FOOD CONTACT SURFACES IN CANNERIES, MEAT PACKING PLANTS, DAIRIES, AND OTHER FOOD INDUSTRIES:** RINSE METHOD: A solution of 100 ppm available chlorine may be used in the sanitizing solution, if a chlorine test kit is available. Solutions containing an initial concentration of 100 ppm available chlorine must be tested and adjusted periodically to insure that the available chlorine does not drop below 50 ppm. Prepare a 100 ppm sanitizing solution by thoroughly mixing 2-1/2 oz. of this product with 10 gallons of water. If no test kit is available, prepare a sanitizing solution by thoroughly mixing 5 oz. of this product with 10 gallons of water, to provide approximately 200 ppm available chlorine by weight. Clean equipment surfaces in the normal manner. Prior to use, rinse all surfaces thoroughly with the sanitizing solution, maintaining contact with the sanitizer for at least 2 minutes. If solution contains less than 50 ppm available chlorine, as determined by a suitable test kit, either discard the solution or add sufficient product to reestablish a 200 ppm residual. Do not rinse equipment with water after treatment and do not soak equipment overnight. Sanitizers used in automated systems may be used for general cleaning but may not be reused for sanitizing purposes.

**IMMERSION METHOD:** A solution of 100 ppm available chlorine may be used in the sanitizing solution if a chlorine test kit is available. Solutions containing an initial concentration of 100 ppm available chlorine must be tested and adjusted periodically to insure that the available chlorine does not drop below 50 ppm. Prepare a 100 ppm sanitizing solution by thoroughly mixing 2-1/2 oz. of this product with 10 gallons of water. If no test kit is available, prepare a sanitizing solution by thoroughly mixing 5 oz. of this product with 10 gallons of water, to provide approximately 200 ppm available chlorine by weight. Clean equipment in the normal manner. Prior to use, immerse equipment in the sanitizing solution for at least 2 minutes and allow the sanitizer to drain. If solution contains less than 50 ppm available chlorine, as determined by a suitable test kit, either discard the solution or add sufficient product to reestablish a 200 ppm residual. Do not rinse equipment with water after treatment. Sanitizers used in automated systems may be used for general cleaning, but may not be reused for sanitizing purposes.

**SANITIZATION OF SURFACES SUCH AS CUTTING BLOCKS, WOOD-TOP TABLES, STEAM TABLE BOARDS, AND OTHER POROUS FOOD CONTACT SURFACES:** RINSE METHOD: Prepare a sanitizing solution by thoroughly mixing 15 oz. of this product with 10 gallons of water, to provide approximately 600 ppm available chlorine by weight. Clean surfaces in the normal manner. Prior to use, rinse all surfaces thoroughly with the sanitizing solution, maintaining contact with the sanitizer for at least 2 minutes. Rinse equipment with water after treatment and do not soak equipment overnight.

**SPRAY/FOG METHOD:** Pre-clean all surfaces after use. Prepare a 600 ppm available chlorine sanitizing solution of sufficient size by thoroughly mixing the product in a ratio of 15 oz. product with 10 gallons of water. Use spray or fogging equipment which can resist hypochlorite solutions. Always empty and rinse spray/fog equipment with potable water after use. Thoroughly spray or fog all surfaces until wet, allowing excess sanitizer to drain. Vacate area for at least 2

hours. Prior to using equipment, rinse all surfaces with a 200 ppm available chlorine solution. Prepare a 200 ppm sanitizing solution by thoroughly mixing 5 oz. of this product with 10 gallons of water.

**DISINFECTION OF SURFACES SUCH AS CONCRETE FLOORS, TILED WALLS, AND OTHER NON-POROUS NON-FOOD CONTACT SURFACES IN MEAT AND POULTRY PROCESSING PLANTS:** RINSE METHOD: Prepare a disinfecting solution by thoroughly mixing 15 oz. of this product with 10 gallons of water, to provide approximately 600 ppm available chlorine by weight. Clean equipment surfaces in the normal manner. Prior to use, rinse all surfaces thoroughly with the disinfecting solution, maintaining contact with the solution for at least 10 minutes. Do not rinse equipment with water after treatment and do not soak equipment overnight.

**COMMERCIAL LAUNDRY SANITIZERS**

Wet fabrics or clothes should be spun dry prior to sanitization. Thoroughly mix 5 oz. of this product with 10 gallons of water, to yield 200 ppm available chlorine. Promptly after mixing the sanitizer, add the solution into the prewash, prior to washing fabrics/clothes in the regular wash cycle with a good detergent. Test the level of available chlorine, if solution has been allowed to stand. Add more of this product if the available chlorine level has dropped below 200 ppm.

**COOLING TOWER/EVAPORATIVE CONDENSER WATER:**

**SLUG FEED METHOD:** Initial Dose: When system is noticeably fouled, apply 128 to 256 oz. of this product per 10,000 gallons of water in the system to obtain from 5 to 10 ppm available chlorine. Repeat until control is achieved. Subsequent Dose: When microbial control is evident, add 25 oz. of this product per 10,000 gallons of water in the system daily, or as needed to maintain control and keep the chlorine residual at 1 ppm. Badly-fouled systems must be cleaned before treatment is begun.

**INTERMITTENT FEED METHOD:** Initial Dose: When system is noticeably fouled, apply 128 to 256 oz. of this product per 10,000 gallons of water in the system to obtain 5 to 10 ppm available chlorine. Apply half (or 1/3, 1/4 or 1/5) of this initial dose when half (or 1/3, 1/4, or 1/5) of the water in the system has been lost by blow down. Subsequent Dose: When microbial control is evident, add 25 oz. of this product per 10,000 gallons of water in the system to obtain a 1 ppm residual. Apply half (or 1/3, 1/4, or 1/5) of this initial dose when half (or 1/3, 1/4, or 1/5) of the water in the system has been lost by blow down. Badly-fouled systems must be cleaned before treatment is begun.

**CONTINUOUS FEED METHOD:** Initial Dose: When system is noticeably fouled, apply 128 to 256 oz. of this product per 10,000 gallons of water in the system to obtain 5 to 10 ppm available chlorine.

Subsequent Dose: Maintain this treatment level by starting a continuous feed of 2-1/2 oz. of this product per 1,000 gallons of water lost by blow down to maintain a 1 ppm residual. Badly-fouled systems must be cleaned before treatment is begun.

American Made Proudly by  
**ZEP MANUFACTURING COMPANY**

Division of National Services Industries, Inc.

ATLANTA, GEORGIA 30301

CLEAN ACROSS AMERICA AND THROUGHOUT THE WORLD™

This product is designed exclusively for industrial and institutional use by trained, professional maintenance personnel. Label directions and precautions must be followed exactly. Zep Manufacturing Company will not be responsible for any injury, loss, or damage if product is used in any manner not in compliance with label directions, or if precautions are not observed.

Net Contents: \_\_\_\_\_



**LICENSED**

PERIOD 2015-2017 LIC. NO.

9625.16



**LICENSED**

PERIOD 2012-2014 LIC. NO.



**LICENSED**

PERIOD 2009-2011 LIC. NO.



**LICENSED**

PERIOD 2006-2008 LIC. NO.