



LICENSED

PERIOD 2015-2017 LIC. NO. **9649.32**



LICENSED

PERIOD 2012-2014 LIC. NO.

**HOSPITAL
INSTITUTIONAL
INDUSTRIAL USE**

CLEAR LEMON

DISINFECTANT, VIRUCIDE* & CLEANER

**FOR HOSPITAL, INSTITUTIONAL
AND INDUSTRIAL USE**

*Human immunodeficiency virus, Type 1 (HIV-1)

ACTIVE INGREDIENT:
Alkyl (C₁₄ 50%, C₁₂ 40%, C₁₆ 10%)
dimethyl benzyl ammonium chloride ... 3.3%

INERT INGREDIENTS: 96.7%

**KEEP OUT OF REACH OF CHILDREN
DANGER**

See back panel for additional precautionary
and first aid statements.

**Net Contents:
3.8 LITERS/1 U.S. GALLON**

CARROLL

**PROFESSIONAL
PRODUCTS**



LICENSED

PERIOD 2009-2011 LIC. NO.



LICENSED

PERIOD 2006-2008 LIC. NO.

This disinfectant has a pleasing lemon fragrance and makes a bright, clear yellow solution when mixed with water. Do not mix with soap or other detergents.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling. For use on precleaned, hard, inanimate surfaces. For heavy soil or organic matter, a precleaning step is required. Rinse with clear water. Apply disinfectant solution of two ounces per gallon of water with cloth or mop to toilets, restroom floors, shower stalls, diaper pails and garbage cans. Wet all surfaces thoroughly. Use four ounces per gallon of water for sick rooms, hospitals, or clinics. Keep surfaces wet for at least 10 minutes. Prepare a fresh solution daily or when use solution becomes visibly dirty.

This product is effective against the following organisms:

Pseudomonas aeruginosa	Staphylococcus aureus
Salmonella choleraesuis	Human immunodeficiency virus, Type 1

This product is not to be used as a terminal sterilant/high level disinfectant on any surface or instrument that (1) is introduced directly into the human body, either into or in contact with the bloodstream or normally sterile areas of the body, or (2) contacts intact mucous membranes but which does not ordinarily penetrate the blood barrier or otherwise enter normally sterile areas of the body. This product may be used to preclean or decontaminate critical or semi-critical medical devices prior to sterilization or high level disinfection.

DISINFECTION FOR TOILET BOWLS: Flush toilet and add 2 ounces of disinfectant solution directly to the bowl water. Swab the bowl completely using a scrub brush or toilet mop, making sure to get under the rim. Keep surfaces wet for 10 minutes and flush.

***KILLS HIV-1 ON PRECLEANED ENVIRONMENTAL SURFACES/OBJECTS PREVIOUSLY SOILED WITH BLOOD/BODY FLUIDS** in health care settings or other settings in which there is an expected likelihood of soiling of inanimate surfaces/objects with blood or body fluids, and in which the surfaces/objects likely to be soiled with blood or body fluids can be associated with the potential for transmission of human immunodeficiency virus, Type 1 (HIV-1) (associated with AIDS).

SPECIAL INSTRUCTIONS FOR CLEANING AND DECONTAMINATION AGAINST HIV OF SURFACES/OBJECTS SOILED WITH BLOOD/BODY FLUIDS: PERSONAL PROTECTION: Cleanup should always be done wearing protective latex gloves, gowns, masks, and eye protection. **CLEANING PROCEDURE:** Blood and other body fluids must be thoroughly cleaned from surfaces and objects

before application of the disinfectant. **DISPOSAL OF INFECTIOUS MATERIALS:** Blood and other body fluids should be autoclaved and disposed of according to federal, state and local regulations for infectious waste disposal. **CONTACT TIME:** Effective against HIV-1 (AIDS Virus) on hard, nonporous surfaces in the presence of a moderate amount of organic soil (5% blood serum) when diluted 1:32. Surfaces must remain wet for 10 minutes at room temperature (20-25°C).

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

DANGER: Corrosive. Keep Out of Reach of Children. Causes irreversible eye damage and skin burns. Harmful if swallowed. Do not get in eyes, on skin or on clothing. Wear protective eyewear (goggles, face shield, or safety glasses), protective clothing and rubber gloves when handling. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse.

FIRST AID

IF IN EYES: Hold eye open and rinse slowly and gently with water for 15 – 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. **IF ON SKIN OR CLOTHING:** Take off contaminated clothing. Rinse skin immediately with plenty of water for 15 – 20 minutes. **IF SWALLOWED:** Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor. Do not give anything by mouth to an unconscious person.

Call a poison control center or doctor for treatment advice. Have the product container or label with you when calling a poison control center or doctor or going for treatment.

NOTE TO PHYSICIAN: Probable mucosal damage may contraindicate the use of gastric lavage. Measures against circulatory shock, respiratory depression and convulsion may be needed.

STORAGE AND DISPOSAL

Store in original container in areas inaccessible to children and persons unfamiliar with its proper use. Do not reuse container. Wrap container and discard in the trash.

776-0903

Manufactured By:

CARROLL[®]
COMPANY

EPA Reg. No. 4313-22

EPA Est. No. 4313-TX-1

2900 W. Kingsley Rd. • Garland, Texas 75041
800.527.5722 • www.carrollco.com



SC9900776A1



4

0