



LICENSED

PERIOD 2015-2017 LIC. NO. 9649.43



LICENSED

PERIOD 2012-2014 LIC. NO.



PC 752

Neutracyde 64

One-Step Disinfectant, Germicidal, Detergent and Deodorant

**Disinfectant - Pseudomonacidal - Staphylocidal - Salmonellacidal
Bactericidal - Fungicidal - Mildewstatic - *Virucidal**

ACTIVE INGREDIENTS:

Didecyl dimethyl ammonium chloride	2.54%
n-Alkyl (C ₁₄ 50%, C ₁₂ 40%, C ₁₆ 10%) dimethyl benzyl ammonium chloride	1.69%
INERT INGREDIENTS:	95.77%
Total	100.00%

**KEEP OUT OF REACH OF CHILDREN
DANGER**

See side panel for Precautionary Statements and First Aid

**Net Contents:
3.8 LITERS/1 U.S. GALLON**



PROFESSIONAL PRODUCTS



LICENSED

PERIOD 2009-2011 LIC. NO.



LICENSED

PERIOD 2006-2008 LIC. NO.

PELIGRO: SI NO PUEDE LEER EN INGLES, PREGUNTE A SU SUPERVISOR SOBRE LAS INSTRUCCIONES DE USO APROPIADAS ANTES DE TRABAJAR CON ESTE PRODUCTO.

A multi-purpose, neutral pH, germicidal detergent and deodorant effective in hard water up to 400 ppm (calculated as CaCO₃) in the presence of a moderate amount of soil (5% organic serum) according to the AOAC Use-dilution Test. Disinfects, cleans, and deodorizes in one labor-saving step.

Recommended for use in hospitals, nursing homes, schools, colleges, commercial and industrial institutions, office buildings, veterinary clinics, animal life science laboratories, zoos, federally inspected meat and poultry establishments, tack shops, pet shops, airports, kennels, hotels, motels, breeding establishments, and grooming establishments.

Disinfects, cleans, and deodorizes the following hard nonporous inanimate surfaces: floors, walls, non-medical metal surfaces, non-medical stainless steel surfaces, glazed porcelain, plastic surfaces (such as polypropylene, polystyrene, etc.)

Recommended for use in: federally inspected meat and poultry plants on all hard, nonporous surfaces in inedible product processing areas, non-processing areas, and/or exterior areas. All surfaces must be thoroughly rinsed with potable water.

Effective against the following pathogens: **Bacteria:** *Pseudomonas aeruginosa*¹, *Staphylococcus aureus*², *Salmonella choleraesuis*, *Acinetobacter calcoaceticus*, *Bordetella bronchiseptica*, *Chlamydia psittaci*, *Enterobacter aerogenes*, *Enterobacter cloacae*, *Enterococcus faecalis*-Vancomycin Resistant (VRE), *Escherichia coli*¹, *Fusobacterium necrophorum*, *Klebsiella pneumoniae*¹, *Legionella pneumophila*, *Listeria monocytogenes*, *Pasteurella multocida*, *Proteus mirabilis*, *Proteus vulgaris*, *Salmonella enteritidis*, *Salmonella typhi*, *Salmonella typhimurium*, *Serratia marcescens*, *Shigella flexneri*, *Shigella sonnei*, *Staphylococcus aureus* - Methicillin Resistant (MRSA), *Staphylococcus aureus* - Vancomycin Intermediate Resistant (VISA), *Staphylococcus epidermidis*¹, *Streptococcus faecalis*¹, *Streptococcus pyogenes*.

Viruses: *Adenovirus* type 4, *Hepatitis B Virus* (HBV), *Hepatitis C Virus* (HCV), *Herpes simplex* type 1, *Herpes simplex* type 2, *Human coronavirus*, *HIV-1* (AIDS virus), *Influenza A/Hong Kong*, *Respiratory syncytial virus* (RSV), *Rubella* (German Measles), *Vaccinia*. **Animal Viruses:** *Avian polyomavirus*, *Canine parvovirus* - at 8oz. per gallon, *Canine distemper*, *Feline leukemia*, *Feline picornavirus*, *Infectious bovine rhinotracheitis*, *Infectious bronchitis* (Avian IBV), *Pseudorabies*, *Rabies*, *Transmissible gastroenteritis virus* (TGE). **Fungi:** *Aspergillus niger*, *Candida albicans*, *Trichophyton mentagrophytes*.

¹ATCC & antibiotic-resistant strain ²antibiotic-resistant strain only

DILUTION: 1:64 (660 ppm quat) 2 OUNCES PER GALLON OF WATER

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

This product is not for use on medical device surfaces.

DISINFECTION/CLEANING/DEODORIZING DIRECTIONS: Remove heavy soil deposits from surface. Then thoroughly wet surface with a use-solution of 2 ounces of the concentrate per gallon of water. The use-solution can be applied with a cloth, mop, sponge, or coarse spray, or by soaking. For sprayer applications, use a coarse spray device. Spray 6-8 inches from the surface, rub with a brush, cloth or sponge. Do not breathe spray. Let solution remain on surface for a minimum of 10 minutes. Rinse or allow to air dry. Rinsing of floors is not necessary unless they are to be waxed or polished. Food contact surfaces must be thoroughly rinsed with potable water. This product must not be used to clean the following food contact surfaces: utensils, glassware and dishes. Prepare a fresh use-solution daily or more often if the solution becomes visibly dirty or diluted.

TOILET BOWLS: Swab bowl with brush to remove heavy soil prior to cleaning or disinfecting. Clean by applying use-solution around the bowl and up under the rim. Stubborn stains may require brushing. To disinfect, first pre-clean to remove heavy soil, then remove or expel over the inner trap the residual bowl water. Pour in three ounces of the diluted solution. Swab the bowl completely using a scrub brush or mop, making sure to get under rim. Let stand for 10 minutes or overnight, then flush.

FUNGICIDAL DIRECTIONS: For use in areas such as locker rooms, dressing rooms, shower and bath areas and exercise facilities follow disinfection directions.

MILDEWSTATIC INSTRUCTIONS: Will effectively control the growth of mold and mildew, plus the odors caused by them when applied to hard, nonporous surfaces such as walls, floors, and table tops. Apply use-solution of 2 ounces per gallon of water when

a cloth, mop, sponge, or coarse spray. Make sure to wet all surfaces completely. Let air dry. Repeat application weekly or when growth reappears.

VETERINARY PRACTICE / ANIMAL CARE / ANIMAL LABORATORY / ZOOS / PET SHOP / KENNELS DISINFECTION DIRECTIONS: For cleaning and disinfecting hard nonporous surfaces: equipment used for animal food or water, utensils, instruments, cages, kennels, stables, catteries, etc. Remove all animals and feeds from premises, animal transportation vehicles, crates, etc. Remove all litter, droppings and manure from floors, walls and surfaces of facilities occupied or traversed by animals. Empty all feeding and watering appliances. Thoroughly clean all surfaces with soap or detergent and rinse with water. Saturate surfaces with a use-solution of 2 oz. of this product per gallon of water for a period of 10 minutes. Wipe or allow to air dry. Ventilate buildings and other closed spaces. Thoroughly scrub all treated surfaces that come into contact with food, including equipment used for feeding or watering, with soap or detergent and rinse with potable water before reuse. Do not house animals or employ equipment until treatment has been absorbed, set or dried.

FOR CANINE PARVOVIRUS ACTIVITY: This product is effective against Canine Parvovirus at 8 oz. per gallon. Follow Veterinary Practice / Animal Care / Animal Laboratory / Zoos / Pet Shop / Kennels Disinfection Directions.

***KILLS HIV-1 (AIDS VIRUS) AND HBV (HEPATITIS B VIRUS) AND HCV (HEPATITIS C VIRUS) OF PRECLEANED ENVIRONMENTAL SURFACES/OBJECTS PREVIOUSLY SOILED WITH BLOOD/BODY FLUIDS** in health care settings or other settings in which there is an expected likelihood of soiling of inanimate surfaces/objects with blood/body fluids, and in which the surfaces/objects likely to be soiled with blood/body fluids can be associated with the potential for transmission of Human Immunodeficiency Virus Type 1 (HIV-1) (associated with AIDS) or Hepatitis B Virus (HBV) or Hepatitis C Virus (HCV).

SPECIAL INSTRUCTIONS FOR CLEANING AND DECONTAMINATION AGAINST HIV-1 (AIDS VIRUS) OR HBV OR HCV OF SURFACES/OBJECTS SOILED WITH BLOOD/BODY FLUIDS: PERSONAL PROTECTION: Disposable protective gloves, gowns, face masks, or eye coverings as appropriate, must be worn during all cleaning of blood/body fluids and during decontamination procedures. **CLEANING PROCEDURES:** Blood/body fluids must be thoroughly cleaned from surfaces/objects before application of disinfectant. **CONTACT TIME:** HIV-1 (AIDS virus) is inactivated after a contact time of 4 minutes at 25°C (77°F) (room temperature). HBV and HCV are inactivated after a 10 minute contact time. Use a 10-minute contact time for other viruses, fungi, and bacteria listed.

DISPOSAL OF INFECTIOUS MATERIALS: Blood/body fluids should be autoclaved and disposed of according to federal, state, and local regulations for infectious waste disposal.

PRECAUTIONARY STATEMENTS

DANGER: HAZARDS TO HUMANS AND DOMESTIC ANIMALS

Corrosive. Causes irreversible eye damage. Do not get in eyes or clothing. Wear protective eyewear (goggles, face shield, or safety glasses). Avoid contact with the skin. Harmful if inhaled. Avoid breathing spray mist. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash clothing before reuse.

FIRST AID

IF IN EYES: Hold eye open and rinse slowly and gently with water for 15 - 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. **IF ON SKIN OR CLOTHING:** Take off contaminated clothing. Rinse skin immediately with plenty of water for 15 - 20 minutes. **IF INHALED:** Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control center or doctor for treatment advice. Have the product container or label with you when calling a poison control center or doctor or going for treatment.

NOTE TO PHYSICIAN: Probable mucosal damage may contraindicate the use of gastric lavage.

STORAGE AND DISPOSAL

Store in original container in areas inaccessible to children. Do not reuse empty container. Wrap and discard in trash or recycle. 752 0505

EPA Reg. No. 47371-131-4313

EPA Est. No. 4313-TX-1

CARROLL
COMPANY

2900 W. Kingsley Rd. Garland, Texas 75041
800.527.5722 www.carrollco.com



PLEASE RECYCLE!



FAX ON DEMAND
888.888.3035
888.888.9964

MADE
in the
U.S.A.

SC9900752A1



6