



Department of Agriculture
STATE OF HAWAII

LICENSED

PERIOD **2015-2017** LIC. NO.

9653.124

activyl[®] TICKPLUS

For Dogs and Puppies

Once-A-Month Topical Flea and Tick Treatment for dogs and puppies 8 weeks and older and over 22 lb and up to 44 lb ONLY
INTENDED TO BE SOLD BY VETERINARIANS

DIRECTIONS FOR USE: Please refer to package insert for complete directions for use.
It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

TO PREVENT HARM TO YOU AND YOUR PET, READ ENTIRE LABEL AND DIRECTIONS BEFORE EACH USE. FOLLOW ALL DIRECTIONS AND PRECAUTIONARY STATEMENTS CAREFULLY.

RESTRICTIONS

- DO NOT USE ON CATS or OTHER ANIMALS.
- DO NOT ALLOW CHILDREN TO APPLY PRODUCT.
- DO NOT APPLY MORE THAN ONCE PER MONTH.
- DO NOT ALLOW CONTACT WITH TREATED AREAS (APPLICATION SITES) UNTIL DRY.

FIRST AID	
If in Eyes	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
If On Skin	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.
If Swallowed	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to by a poison control center or doctor. • Do not give anything to an unconscious person.

HOT LINE NUMBER
Have the product container or label with you when calling a poison control center or doctor or going for treatment.
You may also contact the Rocky Mountain Poison Center at 1-303-595-4869 for emergency medical treatment information.

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS

CAUTION. Causes moderate eye irritation.
Harmful if swallowed. Harmful if absorbed through skin.
Avoid contact with eyes, skin or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet.
Remove and wash contaminated clothing before reuse.

Permethrin-, cyphenothrin- and phenothrin-containing products:

DO NOT USE ON CATS - MAY BE FATAL
Keep cats away from treated dogs for 24 hours.
If applied or ingested, contact your veterinarian immediately.



DO NOT USE ON CATS

HAZARDS CONTINUED ON SIDE PANEL

For Dogs and Puppies
**activyl[®]
TICKPLUS**



Department of Agriculture
STATE OF HAWAII

LICENSED

PERIOD **2012-2014** LIC. NO.

For Dogs and Puppies
8 weeks and older and
over 22 lb and up
to 44 lb ONLY

activyl[®] TICKPLUS

For Dogs and Puppies
INTENDED TO BE SOLD BY
VETERINARIANS

HAZARDS TO DOMESTIC ANIMALS

For external use on dogs and puppies only. Do not use on puppies under 8 weeks of age.

Consult a veterinarian before using this product on debilitated, aged, and medicated animals. Do not use on pregnant or nursing animals. Individual sensitivities, while rare, may occur after using any pesticide product. If signs of sensitivity occur bathe your pet with mild soap and rinse with large amounts of water. If signs continue, consult a veterinarian immediately. Certain medications can interact with pesticides. Consult a veterinarian before using on medicated animals.

DO NOT USE ON CATS.
May be toxic or potentially fatal if applied to or ingested by cats.

Accidental application to cats and/or grooming a recently treated dog may result in tremors and/or uncoordinated muscle movements. If this occurs, immediate veterinary care should be provided. Cats which actively groom or engage in close physical contact with recently treated dogs may be at risk of serious harmful effects.
PHYSICAL OR CHEMICAL HAZARDS
Flammable. Keep away from heat and open flame.

2-D bar code
batching legends printed online

activyl[®] TICKPLUS

For Dogs and Puppies
Once-A-Month Topical Flea and Tick Treatment for dogs and puppies 8 weeks and older and over 22 lb and up to 44 lb ONLY
INTENDED TO BE SOLD BY
VETERINARIANS



Active Ingredients:

Indoxacarb:
(S)-methyl 7-chloro-2,5-dihydro-2-[[[(methoxycarbonyl)[4-trifluoromethoxy]phenyl]amino]-carbonyl]indeno[1,2-e][1,3,4]oxadiazine-4a(3H)-carboxylate 13.01%

Permethrin*:
3-(phenoxybenzyl)-methyl-(±) cis, trans-3-(2,2-dichloroethenyl)-2,2-Dimethylcyclopropanecarboxylate... 44.49%

Other Ingredients:..... 44.50%

Total 100.00%

* cis/trans ratio Min. 35%(±) cis and Max. 65%(±)trans

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

SEE BACK PANEL FOR FIRST AID AND ADDITIONAL PRECAUTIONARY STATEMENTS

MERCK
Animal Health



DO NOT USE ON CATS

For Dogs and Puppies
8 weeks and older and
over 22 lb and up
to 44 lb ONLY

activyl[®] TICKPLUS

For Dogs and Puppies
INTENDED TO BE SOLD BY
VETERINARIANS

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.
Pesticide Storage:
Store in a cool, dry place inaccessible to children and pets.
Pesticide Disposal:
If partly filled: Call your local solid waste agency or 1-800-CLEANUP for disposal instructions.
Never place unused product down any indoor or outdoor drain.
Container Disposal:
If empty: Do not reuse this container. Place in trash or offer for recycling if available.

EPA Reg. No. 773-95
EPA Est. No. 6175-LA-1
Net Contents: 1 applicator(s), each 0.03 fl.oz. (1.0 mL)
[Total 0.03 fl.oz. (1.0 mL)]
Made in the USA
Intervet Inc d/b/a Merck Animal Health, Summit, NJ
Copyright © 2011, Intervet Inc., a subsidiary of Merck & Co., Inc.
All rights reserved.
Rev. 9/11



8 713184 113986 1

116980 R2

BR031_01_R1
47x25,5x134,9mm
C M Y K

PACKAGE INSERT – 22 to 44lb carton

CAT WARNING ICON
Permethrin-, cyphenothrin- and phenothrin-
containing products:

DO NOT USE ON CATS - MAY BE FATAL
Keep cats away from treated dogs for 24 hours. If
applied or ingested, contact your veterinarian
immediately.

Activyl® Tick Plus
For Dogs and Puppies

INTENDED TO BE SOLD BY VETERINARIANS

Once-A-Month Topical Flea and Tick Treatment for dogs and puppies 8 weeks and older and
over 22 lb and up to 44 lb ONLY

Active Ingredients:

Indoxacarb:
(S)-methyl 7-chloro-2,5-dihydro-2-[[[(methoxycarbonyl)[4-trifluoromethoxy]phenyl]amino]
-carbonyl]indeno[1,2-e][1,3,4]oxadiazine-4a(3H)-carboxylate.....13.01%
Permethrin*:
3-(phenoxybenzyl)-methyl-(±) cis, trans-3-(2,2-dichloroethenyl)-2,2-
Dimethylcyclopropanecarboxylate.....42.50%

Other Ingredients:.....44.49%
Total 100.00%

*cis/trans ratio Min. 35%(±) cis and Max. 65%(±)trans


**KEEP OUT OF REACH OF CHILDREN
CAUTION**

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

TO PREVENT HARM TO YOU AND YOUR PET, READ ENTIRE LABEL AND DIRECTIONS BEFORE EACH USE.
FOLLOW ALL DIRECTIONS AND PRECAUTIONARY STATEMENTS CAREFULLY.

HOW TO APPLY- Open one foil pouch and remove applicator.



For DOGS over 22 lb and up to 44 lb ONLY

Step 1. Hold applicator in upright position away from your face and snap tip open by bending it and folding it back on itself.

Step 2. The dog should be standing for easy application. Part the hair and place applicator tip against the skin between shoulder blades.

Step 3. Apply entire contents of tube evenly to two spots on the back from between the shoulder blades to the base of the tail. At each spot part the hair until skin is visible. Place the tip of the applicator on the skin and gently squeeze a portion of the solution on the skin. Do not apply an excessive amount of solution at any one spot. Do not allow runoff.

Use one applicator per treatment. Reapply every 30 days or as recommended by your veterinarian. When application is complete, follow Pesticide Disposal under Storage and Disposal section below.

RESTRICTIONS

- DO NOT USE ON CATS OR OTHER ANIMALS.
- DO NOT ALLOW CHILDREN TO APPLY PRODUCT.
- DO NOT APPLY MORE THAN ONCE PER MONTH.
- DO NOT ALLOW CONTACT WITH TREATED AREAS (APPLICATION SITES) UNTIL DRY.

FIRST AID	
If in Eyes	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
If On Skin	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.
If Swallowed	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to by a poison control center or doctor. • Do not give anything to an unconscious person.
HOT LINE NUMBER	
Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact the Rocky Mountain Poison Center at 1-303-595-4869 for emergency medical treatment information.	

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS

CAUTION. Causes moderate eye irritation. Harmful if swallowed. Harmful if absorbed through skin. Avoid contact with eyes, skin or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Remove and wash contaminated clothing before reuse.

HAZARDS TO DOMESTIC ANIMALS

For external use on dogs and puppies only. Do not use on puppies under 8 weeks of age. Consult a veterinarian before using this product on debilitated, aged, and medicated animals. Do not use on pregnant or nursing animals. Individual sensitivities, while rare, may occur after using any pesticide product. If signs of sensitivity occur bathe your pet with mild soap and rinse with large amounts of water. If signs continue, consult a veterinarian immediately. Certain medications can interact with pesticides. Consult a veterinarian before using on medicated animals.

DO NOT USE ON CATS. May be toxic or potentially fatal if applied to or ingested by cats. Accidental application to cats and/or grooming a recently treated dog may result in tremors and/or uncoordinated muscle movements. If this occurs, immediate veterinary care should be provided. Cats which actively groom or engage in close physical contact with recently treated dogs may be at risk of serious harmful effect.

PHYSICAL OR CHEMICAL HAZARDS

Flammable. Keep away from heat and open flame.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

Pesticide Storage: Store in a cool, dry place inaccessible to children and pets.

Pesticide Disposal: If partly filled: Call your local solid waste agency or 1-800-CLEANUP for disposal instructions. Never place unused product down any indoor or outdoor drain.

Container Disposal: If empty: Do not reuse this container. Place in trash or offer for recycling if available

EPA Reg. No. 773-95
EPA Est. No. 6175-LA-1

Net Contents: 1 applicator(s), each 0.07fl.oz. (2.0 mL)
Made in the USA

Intervet Inc d/b/a Merck Animal Health

Summit, NJ 07901 Copyright © 2011, Intervet Inc., a subsidiary of Merck & Co., Inc. All rights reserved. Rev. 11/11

(MAH logo)

DO NOT REMOVE APPLICATOR (TUBE) UNTIL READY TO USE

ACTIVYL Tick Plus for Dogs and Puppies

**For dogs 8 weeks of age and older and
over > 22 lb up to 44 lb ONLY**

Cat Warning Icon

Indoxacarb: ... 13.01%

Permethrin: ... 42.50%

KEEP OUT OF REACH OF CHILDREN

Caution

READ ENTIRE LABEL BEFORE USE

EPA Reg. No. 773-95

EPA Est. No. 6175-LA-1

Intervet Inc

Lot No: XXXXXX

0.03 fl.oz. (1.0mL)

**ACTIVYL Tick Plus for
Dogs and Puppies 8 weeks & older and
>22 lb and up to 44 lb ONLY**

Indoxacarb 13.01%


Permethrin 42.50%

EPA Reg. No. 773-95

Intervet Inc, EPA Est. No. 6175-LA-1

0.02 fl.oz. (0.51mL) Lot No:XXXXXx

**DO NOT USE
ON CATS**



Keep Out of Reach of Children
CAUTION
Read Entire Label Before Use