



LICENSED

PERIOD 2015-2017 LIC. NO.

9653.70



Department of Agriculture
STATE OF HAWAII

LICENSED

PERIOD 2012-2014 LIC. NO.



Department of Agriculture
STATE OF HAWAII

LICENSED

PERIOD 2009-2011 LIC. NO.

copy.

2006-2008

DEPT. OF AGRICULTURE
HAWAII
Period 2009-2011 No. 9653.70


LICENSED

0061-5052-01

ATROBAN® DELICE

POUR-ON INSECTICIDE

NON-SYSTEMIC POUR-ON INSECTICIDE FOR BEEF, LACTATING AND NON-LACTATING DAIRY CATTLE
CONTROLS LICE AND FLIES ON CATTLE — CONTROLS KEDS AND LICE ON SHEEP



Active Ingredient:

Permethrin: 3-(phenoxyphenyl)methyl (±)-cis, trans-3-(2,2-dichloroethyl)-2,2-dimethylcyclopropanecarboxylate*	1.0%
Other Ingredients**	99.0%
	100.0%

* cis/trans ratio: Min 35% (±) cis and max 65% (±) trans
** Contains petroleum distillates.

KEEP OUT OF REACH OF CHILDREN CAUTION



PRECAUTIONARY STATEMENTS, HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION — Avoid contact with eyes.

TREATMENT OF PRACTICAL TREATMENT

If in Eyes: Immediately flush eyes with plenty of water. Get medical attention if discomfort persists.
If Swallowed: Call a physician immediately. DO NOT INDUCE VOMITING UNLESS UNDER MEDICAL ATTENTION.
Hole to Physician: Solvent presents aspiration hazard. Gastric lavage is indicated if material was taken internally.

ENVIRONMENTAL HAZARDS

This pesticide is extremely toxic to fish. Use with care when applying to areas adjacent to any body of water. Do not add directly to water. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

PHYSICAL OR CHEMICAL HAZARDS

Do not use or store near heat or open flame.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.
Storage: Keep container sealed when not in use. Do not store near food or feed. Pesticide Disposal: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility. Container Disposal: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill or incineration, or if allowed by State and local authorities, by burning. If burned, stay out of smoke.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

READY TO USE — No dilution necessary.

Apply To	Target Insects	Application Instructions
Lactating and Non-Lactating Dairy Cattle and Beef Cattle and Calves	LICE Horn flies, Face flies and Aids in Control of, Horse flies, Stable flies, Mosquitoes, Black flies, and Ticks.	Pour along back and down face. Apply 1/2 fl oz (15 cc) per 100 lb body wt of animal, up to a maximum of 5 fl oz for any one animal.
Sheep	Sheep Keds LICE	Pour along back. Apply 1/4 fl oz (7.5 cc) per 50 lb body wt of animal, up to a maximum of 3 fl oz for any one animal.

For cattle and sheep, repeat treatment as needed, but not more than once every 2 weeks. For optimum lice control, two treatments at 14-day intervals are recommended.

Special Note: Atroban® DELICE Pour-On Insecticide is not effective in controlling cattle grubs.


NOTICE OF WARRANTY

SCHERING-PLOUGH ANIMAL HEALTH CORP. MAKES NO WARRANTY OF MERCHANTABILITY, FITNESS FOR ANY PARTICULAR PURPOSE, OR OTHERWISE EXPRESSED OR IMPLIED CONCERNING THIS PRODUCT OR ITS USES WHICH EXTEND BEYOND THE USE OF THE PRODUCT UNDER NORMAL CONDITIONS IN ACCORDANCE WITH THE STATEMENTS ON THIS LABEL.

EPA Reg. No. 773-66 EPA Est. No. 6175-LA-1
Copyright © 1997, Schering-Plough Animal Health Corp., Union, NJ 07083. All rights reserved. 20729422 Rev. 12/00


Schering-Plough Animal Health

Net Contents: 2.5 Gallons (9.5 L)



ATROBAN® DELICE

POUR-ON INSECTICIDE



**NON-SYSTEMIC POUR-ON INSECTICIDE FOR BEEF,
LACTATING AND NON-LACTATING DAIRY CATTLE
CONTROLS LICE AND FLIES ON CATTLE
CONTROLS KEDS AND LICE ON SHEEP**

Active Ingredient:

Permethrin: 3-(phenoxyphenyl)methyl (+)-cis, trans-3-(2,2-dichloroethyl)-2,2-dimethylcyclopropanecarboxylate* 1.0%
Other Ingredients** 99.0%
100.0%

*cis:trans ratio: Min 35% (+) cis and max 65% (+) trans

**Contains petroleum distillates.

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

**SEE BACK PANEL FOR ADDITIONAL
PRECAUTIONARY STATEMENTS**

EPA Reg. No. 773-66 EPA Est. No. 6175-LA-1

Copyright © 1997, Schering-Plough Animal Health Corp., Union, NJ 07083
All rights reserved. 20723025 Rev. 12/00

**Treats 46
550 lb Cattle**

 Schering-Plough Animal Health

**Net Contents:
One U.S. Gallon (3.785 L)**

ATROBAN® DELICE

POUR-ON INSECTICIDE

PRECAUTIONARY STATEMENTS, HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION – Avoid contact with eyes.

STATEMENT OF PRACTICAL TREATMENT

If in Eyes: Immediately flush eyes with plenty of water. Get medical attention if discomfort persists.

If Swallowed: Call a physician immediately. DO NOT INDUCE VOMITING UNLESS UNDER MEDICAL ATTENTION.

Note to Physician: Solvent presents aspiration hazard. Gastric lavage is indicated if material was taken internally.

ENVIRONMENTAL HAZARDS

This pesticide is extremely toxic to fish. Use with care when applying to areas adjacent to any body of water. Do not add directly to water. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

PHYSICAL OR CHEMICAL HAZARDS

Do not use or store near heat or open flame.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

Storage: Keep container sealed when not in use. Do not store near food or feed.

Outside Disposal: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

Container Disposal: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill or incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

READY TO USE – No dilution necessary.

Apply To	Target Insects	Application Instructions
Lactating and Non-Lactating Dairy Cattle and Beef Cattle and Calves	LICE Horn flies, Face flies and Aids in Control of: Horse flies, Stable flies, Mosquitoes, Black flies, and Ticks.	Pour along back and down face. Apply 1/2 fl oz (15 cc) per 100 lbs body wt of animal, up to a maximum of 5 fl oz for any one animal.
Sheep	Sheep Keds LICE	Pour along back. Apply 1/4 fl oz (7.5 cc) per 50 lb body wt of animal, up to a maximum of 3 fl oz for any one animal.

For cattle and sheep, repeat treatment as needed, but not more than once every 2 weeks. For optimum lice control, two treatments at 14-day intervals are recommended.

Special Note: Atroban® DELICE Pour-On Insecticide is not effective in controlling cattle grubs.

NOTICE OF WARRANTY

SCHERING-PLOUGH ANIMAL HEALTH CORP. MAKES NO WARRANTY OF MERCHANTABILITY, FITNESS FOR ANY PARTICULAR PURPOSE, OR OTHERWISE EXPRESSED OR IMPLIED CONCERNING THIS PRODUCT OR ITS USES WHICH EXTEND BEYOND THE USE OF THE PRODUCT UNDER NORMAL CONDITIONS IN ACCORDANCE WITH THE STATEMENTS ON THIS LABEL.

20722916 Rev. 8/00

