DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

DO NOT ALLOW CHILDREN TO APPLY PRODUCT. TO PREVENT HARM TO YOU AND YOUR DOG, READ ENTIRE LABEL BEFORE EACH USE. FOLLOW ALL DIRECTIONS AND PRECAUTIONARY STATEMENTS CAREFULLY.

Do not use on rabbits or animals other than dogs. Do not allow your dog to ingest this product. Do not use on puppies under 12 weeks of age. Use entire contents of tube vial on each dog. Do not split one tube between dogs. Do not use multiple tubes on one dog. Weigh your dog to be sure you are applying the right dose formulated for the weight of your dog. Separate the treated dog from all other dogs and cats for 24 hours after treatment has been applied.

Side Effects: Monitor your dog after application. The most common signs of ingestion are excessive salivation and foaming at the mouth. If these symptoms occur, immediately feed your dog and continue to monitor your dog for the next 24 hours. Some dogs may experience temporary startle effects when any product is applied. Dogs may experience some temporary irritation at the site of product application such as redness, scratching or other signs of discomfort. If signs of sensitivity occur, bathe your dog with a mild soap and rinse with large amounts of water. If signs persist or become more severe within a few days of application, consult a veterinarian immediately by calling 1-800-660-1842. If your dog has an unusual reaction to the initial application, consult a veterinarian before repeating application.

DO NOT USE ON CATS. May be toxic and POTENTIALLY FATAL if applied to or ingested by a cat. Keep cats away from treated dogs for 24 hours. Cats that actively groom or engage in close physical contact with treated dog may be at risk of serious harmful effects. Cats exhibiting signs of ingestion such as excessive salivation and foaming at the mouth should be taken to the veterinarian immediately.

STORAGE AND DISPOSAL

PESTICIDE STORAGE: Do not remove tube from the pack until ready to use. Store in a cool (below 25˚C) dry place inaccessible to children and pets. Do not refrigerate. Protect from direct sunlight.

CONTAINER DISPOSAL: If empty: Non-refillable container. Do not reuse or refill this container. Offer for recycling if available. **If partly filled:** Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

WARRANTY

Read the entire Directions for Use and Warranty before using this product. By using this product, user or buyer accepts the following warranty. The directions for use of this product are believed to be adequate and must be followed carefully. It is impossible to eliminate all risks associated with the use of this product. Unintended consequences may result because of unknown factors. All such risks shall be assumed by the user or buyer. Merial is committed to providing high quality products. To the extent consistent with applicable law, Merial makes no warranties, express or implied, of merchantability or of fitness for a particular purpose or otherwise, that extend beyond statements on this label. To the extent consistent with applicable law, Merial, the manufacturer, or the Seller shall not be liable for indirect, special, incidental or consequential damages. To the extent consistent with applicable law, the exclusive remedy of the user or buyer shall not exceed the purchase price paid.

FRONTLINE® TRITAK™ FOR DOGS

FOR USE ONLY ON DOGS 23-44lbs

3 DOSES

FRONTLINE® TRITAK™ FOR DOGS

FOR USE ONLY ON DOGS 23-44lbs

3 DOSES

3 DOSES

NET CONTENTS: 0.135 FL OZ (4.02 mL)
3–0.045 FL OZ (1.34mL) APPLICATORS

ACTIVE INGREDIENTS:

Fipronil	9.8%
Cyphenothrin	5.2%
(S)-methoprene	8.8%
OTHER INGREDIENTS:	76.2%
TOTAL	100.0%

KEEP OUT OF REACH OF CHILDREN

WARNING. SEE BACK LABEL PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS

DO NOT USE ON CATS

MERIAL — A SANOFI COMPANY

CONSULT VETERINARIAN BEFORE USE

Kills fleas, flea eggs and flea larvae

Kills ticks including those that may transmit Lyme disease, and chewing lice. Controls mites that may cause sarcoptic mange.

Starts killing fleas and ticks in just 5 minutes

FRONTLINE® TRITAK™ FOR DOGS

FOR USE ONLY ON DOGS 23-44lbs

3 DOSES

STATE OF HAWAII
Department of Agriculture
ACCEPTED
LICENSE NO. 9671.21

FRONTLINE® TRITAK™ FOR DOGS

- Triple Action: Effectively breaks the flea lifecycle
- Prevents all flea stages · Kills flea eggs and flea larvae
- Starts killing fleas and ticks in just five minutes
- Fast-Acting • Long-Lasting • Waterproof

SATISFACTION GUARANTEED — MERIAL

- Adult Fleas
- Flea Eggs
- Flea Larvae
- Ticks
- Chewing Lice

HOW TO OPEN

Remove product tubes from the package. Separate one tube from the others. Hold the tube with notched end pointing up and away from the face and body. Use scissors to cut off the narrow end at the notches along the line.

HOW TO APPLY

Invert tube over dog and use open end to part dog's hair. Squeeze tube firmly to apply all of the solution to the dog's skin, as directed below. Repeat application may be made if necessary, but do not apply more often than once every 30 days.

ONLY FOR DOGS WEIGHING BETWEEN 23-44 LBS.

Apply one tube (1.34 ml) (0.045 fl. oz.) as a spot on to the dog's back between the shoulder blades.

FREQUENCY OF APPLICATION

Use FRONTLINE® TRITAK™ For Dogs monthly for complete control of fleas, ticks and chewing lice. Studies show that FRONTLINE® TRITAK™ For Dogs kills fleas (starts killing in as little as 5 minutes) for up to three months. If your pet is at high risk for flea reinfestation or in a highly infested environment, apply monthly. Apply monthly to control ticks and chewing lice. FRONTLINE® TRITAK™ For Dogs remains effective even after bathing, water immersion or exposure to sunlight. Allow treated area to dry thoroughly. Do not reapply for 30 days.

Please call our experts at **1-800-660-1842**, Monday-Friday, 8am-8pm EST with any questions or comments about flea, tick, chewing lice or sarcoptic mite control or FRONTLINE® TRITAK™ For Dogs.

Distributed by: Merial Limited • 3239 Satellite Blvd., Duluth, GA 30096-4640, USA • *MADE IN USA*

®FRONTLINE is a registered trademark, and ™TRITAK is a trademark, of Merial.
©2013 Merial. All rights reserved.

EPA Reg. No. 2517-142-65331 EPA Est. No. 2517-NE-02 AHRC P004301-1 72336

FRONTLINE® TRITAK™ FOR DOGS

FRONTLINE® TRITAK™ For Dogs contains fipronil, cyphenothrin, and the insect growth regulator (IGR) (S)-methoprene to effectively kill fleas (starts killing in 5 minutes), ticks (starts killing in 5 minutes), and kills chewing lice. The active ingredients in FRONTLINE® TRITAK™ For Dogs feature a sustained release formula developed for fast-acting and long-lasting control of fleas, ticks, and chewing lice for up to 1 month on dogs.

READ ENTIRE LABEL BEFORE EACH USE. USE ONLY ON DOGS AND PUPPIES OVER 12 WEEKS OF AGE.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS. WARNING. Causes substantial but temporary eye injury. Do not get in eyes or on clothing. Harmful if swallowed. Harmful if absorbed through skin. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals. Wash hands thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet. Remove and wash contaminated clothing before reuse.

HAZARDS TO DOMESTIC ANIMALS. For external use on dogs only. Individual sensitivities, while rare, may occur after using any pesticide product on dogs. Certain medications can interact with pesticides. Do not treat your dog with more than one pesticide product at a time. Over dosing your dog can result in serious illness and even death. Consult a veterinarian before using on medicated, debilitated, aged, pregnant or nursing dogs. Consult a veterinarian before using on dogs with known organ dysfunction. If your dog is exhibiting signs of and/or is being treated for skin dermatitis, talk to your veterinarian before applying any topical flea and tick control product.

FIRST AID		
If In Eyes	• Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for further treatment advice.	**HOT LINE NUMBER** Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For more information on this pesticide product (including health concerns, medical emergencies or pesticide incidents), you may call 1-800-660-1842.
If On Skin Or Clothing	• Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.	
If Swallowed	• Immediately call a poison control center or doctor. • Do not induce vomiting unless told to do so by a poison control center or doctor. • Have person sip a glass of water if able to do so. • Do not give anything by mouth to an unconscious person.	**NOTE TO PHYSICIAN OR VETERINARIAN** Treat patient symptomatically

FRONTLINE® TRITAK™ FOR DOGS
Fipronil 9.8% w/w, Cyphenothrin 5.2% w/w
(S)-methoprene 8.8% w/w 1.34 mL (0.045 fl oz)
FOR DOGS 23-44 LBS
P004291

FRONT

KEEP OUT OF REACH OF CHILDREN
WARNING: Read directions and precautions
before using. Use scissors to open.
EPA REG. No. 2517-142-65331
P004283

BACK