

3M™ C. diff Solution Tablets is a Hospital Use Disinfectant. 3M™ C. diff Solution Tablets is for use in: Hospitals, nursing homes, medical and dental offices and clinics, operating rooms, isolation wards, and medical research facilities. 3M™ C. diff Solution Tablets is a disinfectant that disinfects pre-cleaned, hard, non-porous, inanimate surfaces. This cleaning process may be accomplished with any cleaner solution including 3M™ C. diff Solution Tablets. 3M™ C. diff Solution Tablets provides effective cleaning strength that will not dull high gloss floor finishes with repeated use. Dissolves in 2 minutes.

**STORAGE AND DISPOSAL**

Do not contaminate water, food, or feed by storage and disposal. **PESTICIDE STORAGE:** Store in a cool, dry, well-ventilated area at a temperature below 40°C/104°F. Avoid moisture getting into container. **PESTICIDE DISPOSAL:** Pesticide may be acutely hazardous. Wastes resulting from the use of this product must be disposed of on-site, or at an approved waste disposal facility.

**CONTAINER HANDLING:** Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available or puncture and dispose of in a sanitary landfill.

Sold and exported by:  
**3M Commercial Solutions Division**  
 St. Paul, MN 55144-1000  
 Made in Ireland  
 www.3m.com/facility  
 3M is a trademark of 3M.  
 Used under license in Canada.  
 © 3M 2016. All rights reserved.  
 84-8718-5590-3



# C. diff Solution Tablets

**ACTIVE INGREDIENT:**  
 Sodium dichloro-s-triazinetriene..... 48.21%\*  
**OTHER INGREDIENTS:** ..... 51.79%  
**TOTAL** ..... 100.00%  
 \*Equivalent to 31.10% active chlorine by tablet weight.  
 Refer to dilution chart for Available Chlorine concentrations.



**KEEP OUT OF REACH OF CHILDREN**  
**DANGER**

**PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS**  
**DANGER: Corrosive. Causes irreversible eye damage. Harmful if swallowed, inhaled, or absorbed through skin.** Do not get in eyes, on skin, or clothing. Avoid breathing dust. Wear chemical-resistant gloves and safety glasses or face shield when making up solution. Wash thoroughly with soap and water after handling, and before eating, drinking, chewing gum, using tobacco, or using the toilet. Remove and wash contaminated clothing before reuse.


**FIRST AID: IF IN EYES:** Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice. **IF SWALLOWED:** Call a poison control center, or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor. Do not give anything by mouth to an unconscious person. **IF INHALED:** Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control center or doctor for further treatment advice. **IF ON SKIN OR CLOTHING:** Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice. Have the product container or label with you when calling a poison control center or doctor, or going for treatment.

**NOTE TO PHYSICIAN:** Probable mucosal damage may contraindicate the use of gastric lavage.  
**IN CASE OF EMERGENCY, CALL: 1-800-364-3577 (1-800-3M-HELPS).**  
**REFER TO SAFETY DATA SHEET (SDS: 35-7889-7) FOR MORE INFORMATION.**  
**PHYSICAL OR CHEMICAL HAZARDS: STRONG OXIDIZING AGENT:** Use only clean dry utensils. Mix only in water. Contamination with moisture, dirt, organic matter or other chemicals or any other foreign matter may start a chemical reaction with generation of heat, liberation of hazardous gases and possible generation of fire and explosion. Avoid any contact with flaming or burning material such as a lighted cigarette. Do not use this product in any chlorinating device which has been used with any inorganic or un-stabilized chlorinated compounds (e.g., calcium hypochlorite). Such use may cause fire or explosion.

**DIRECTIONS FOR USE:**  
 It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Read the entire label and use strictly in accordance with precautionary statements and directions.

**HEALTHCARE DISINFECTION**  
 3M™ C. diff Solution Tablets is a Hospital Use Disinfectant. General healthcare disinfectant (538 ppm available chlorine disinfectant solution) is effective against standard gram positive and gram negative bacteria (*Staphylococcus aureus*, *Pseudomonas aeruginosa* and *Salmonella enterica*) and cold and flu viruses (Respiratory Syncytial Virus, Influenza H1N1). This product is effective as a healthcare disinfectant for micro-organisms and blood borne viruses when used at level of 1076 ppm available chlorine disinfectant solution. It is effective against HIV, Hepatitis B, Hepatitis A, *Enterococcus faecalis* Vancomycin Resistant, *Trichophyton mentagrophytes*, *Klebsiella pneumoniae*, *Staphylococcus epidermidis*, *Escherichia coli* O157:H7, *Staphylococcus aureus* - MRSA & GRSA, *Poliovirus type 1*, *Herpes simplex virus type 1*, *Canine parvovirus*, *Newcastle Disease Virus*, *Pseudorabies*, Feline Calicivirus, Norovirus, Canine Distemper Virus, Infectious Canine Hepatitis, Teschen/Taifu disease, Avian Influenza, *Porcine parvovirus*, Runtling & Sturling Virus (*Ictosynovitis*), *Actinobacillus pleuropneumoniae*, *Bordetella bronchiseptica* (*rhinitis*), *Brachyspira* (*Tropoema/Serpulina*) *hyodysenteriae* (swine dysentery), Gumboro disease, *Streptococcus uberis*, *Streptococcus dysgalactiae*, and *Clostridium perfringens* USDA with a 10 minute contact time. For Transmissible gastroenteritis (TGE), Swine Vesicular disease, African Swine Fever, Hog Cholera/Classical Swine Fever, and Avipox (Fowl Pox). A 30 minute contact time is required. Re-apply product as necessary to ensure surface remains wet.  
 This product is also effective as a healthcare disinfectant for critical areas of potentially contaminated with *Clostridium difficile* spores when used at a level of 4306 ppm available chlorine disinfectant solution. A 4 minute contact time is required.

**NET CONTENTS**  
 140 TABLETS  
 3.3 grams per tablet  
 NET 1.02 lbs (462 g)  
**EPA Reg. No.** 71847-6-10350  
**EPA Est. No.** 71847-IRL-001



This product is also effective as a healthcare disinfectant for critical areas potentially contaminated with *Mycobacterium bovis* (Tb) when used at a level of 5382 ppm available chlorine disinfectant solution. A 4 minute contact time is required.

**Notice to User:** This product is not to be used as a terminal sterilant/high level disinfectant on any surface or instrument that (1) is introduced directly into the human body, either into or in contact with the blood stream or normally sterile areas of the body, or (2) contacts intact mucous membranes, but which does not ordinarily penetrate the blood barrier or otherwise enter normally sterile areas of the body. This product may be used to pre-clean or decontaminate critical or semi-critical medical devices prior to sterilization or high level disinfection.

**GENERAL DISINFECTION DIRECTIONS:** Prepare a 538 ppm solution; (refer to Dilution Chart). Add 1 tablet to a half gallon of water. Apply to pre-cleaned surface with mop, cloth, sponge, brush, wipe, foaming equipment, or coarse trigger sprayer. Allow surface to remain wet for 10 minutes. Allow to air dry. Prepare a fresh solution weekly when using open containers (buckets) or if solution becomes diluted. All treated equipment that will contact food, feed, or drinking water must be rinsed with potable water before reuse.

**DISINFECTION FOR SURFACES CONTAMINATED WITH HIV:** Prepare a 1076 ppm solution (refer to Dilution Chart). Keep in a properly labeled container to protect against contamination. Unused solution must be discarded.

**DISINFECTION/VIROCIDAL DIRECTIONS:** Prepare a 1076 ppm solution (refer to Dilution Chart). Apply use solution to pre-cleaned, hard, non-porous, inanimate surfaces with brush, spray device, sponge, cloth, or mop to wet all surfaces thoroughly. Allow to remain wet for 10 minutes, then remove product by wiping with brush, sponge, or cloth. For sprayer applications use a coarse spray device. Spray 6-8 inches from surface, and rub with brush, sponge, wipe, or cloth. Do not breathe spray mist. Before using this product, food products, and packaging materials must be removed from the room or carefully protected. Prepare fresh solution weekly when using closed containers (spray bottles). Prepare a fresh solution daily when using open containers (buckets) or if solution becomes diluted. All treated equipment that will contact food, feed, or drinking water must be rinsed with potable water before reuse.

**Kills HIV-1, Hepatitis A, and Hepatitis B viruses on PRE-CLEANED ENVIRONMENTAL SURFACES/OBJECTS PREVIOUSLY SOILED WITH BLOOD/BODY FLUIDS:** In health care setting or other settings in which there is an expected likelihood of soiling of inanimate surfaces /objects with blood or body fluids, and in which the surfaces /objects likely to be soiled with blood or body fluids can be associated with the potential for transmission of Human Immunodeficiency Virus Type 1 (HIV-1) (associated with AIDS), Hepatitis A, and Hepatitis B virus.

**SPECIAL INSTRUCTIONS FOR CLEANING AND DECONTAMINATION AGAINST HIV-1 OF SURFACES/OBJECTS SOILED WITH BLOOD/BODY FLUIDS: PERSONAL PROTECTION:** Specific barrier protection items to be used when handling items soiled with blood or body fluids are disposable latex gloves, gowns, masks, and eye coverings. **CLEANING PROCEDURE:** Blood and other body fluids must be thoroughly cleaned from surfaces and objects before application of 3M™ C. diff Solution Tablets. This cleaning process may be accomplished with any cleaning solution including 3M™ C. diff Solution Tablets. **DISPOSAL OF INFECTIOUS MATERIALS:** Blood and other body fluids should be autoclaved and disposed of according to federal, state, and local regulations for infectious waste disposal. **CONTACT TIME:** Leave surfaces wet for 10 minutes.

**DISINFECTION FOR SURFACES CONTAMINATED WITH CLOSTRIDIUM DIFFICILE IN 4 MINUTES**  
**Special Label Instructions for Cleaning Prior to Disinfection against Clostridium difficile spores: Personal Protection:** Wear appropriate barrier protection such as gloves, gowns, masks or eye covering. **Cleaning Procedure:** Fecal matter/waste must be thoroughly cleaned from surfaces/objects before disinfection by application with a clean cloth, mop, and/or sponge saturated with the disinfectant product. Cleaning is to include vigorous wiping

and/or scrubbing, until all visible soil is removed. Special attention is needed for high-touch surfaces. Surfaces in patient rooms are to be cleaned in an appropriate manner, such as from right to left or left to right, on horizontal surfaces, and top to bottom, on vertical surfaces, to minimize spreading of the spores. Restrooms are to be cleaned last. Do not reuse soiled cloths.

**Infectious Materials Disposal:** Materials used in the cleaning process that may contain feces/wastes are to be disposed of immediately in accordance with local regulations for infectious materials disposal. Prepare a 4306 ppm solution; (refer to Dilution Chart). Add 4 tablets to 1 quart of water. Apply to pre-cleaned surface with mop, cloth, sponge, brush, wipe, foaming equipment, or coarse trigger sprayer. Allow surface to remain wet for 4 minutes. Allow to air dry. Prepare a fresh solution weekly when using closed containers (spray bottles). Prepare a fresh solution daily when using open containers (buckets) or if solution becomes diluted. All treated equipment that will contact food, feed, or drinking water must be rinsed with potable water before reuse. See package insert for additional directions for use.

**GENERAL DISINFECTION WITHOUT PRECLEAN**  
 3M™ C. diff Solution Tablets is a healthcare disinfectant when used at a level of 4,306 ppm available chlorine disinfectant solution. It is effective against Norovirus with a 1-minute contact time. It is effective against *Salmonella enterica*, *Staphylococcus aureus*, *Pseudomonas aeruginosa*, *carbapenem resistant Klebsiella pneumoniae*, *Acinetobacter baumannii* with a 4 minute contact time.

**GENERAL DISINFECTION WITHOUT PRECLEAN DIRECTIONS:** Prepare 4.306 ppm solution (refer to Dilution Chart). Add 4 tablets to 1 quart of water. Apply to surface with mop, cloth, sponge, brush, wipe, foaming equipment, or coarse trigger sprayer. Allow surface to remain wet for 4 minutes. Prepare a fresh solution weekly when using closed containers (spray bottles). Prepare a fresh solution daily when using open containers (buckets) or if solution becomes diluted. All treated equipment that will contact food, feed, or drinking water must be rinsed with potable water before reuse.

**DISINFECTION FOR SURFACES CONTAMINATED WITH MYCOBACTERIUM BOVIS (Tb) IN 4 MINUTES**

**Special Label Instructions for Cleaning Prior to Disinfection against Mycobacterium bovis:** This product when used as directed below is effective against *Mycobacterium bovis*. This product can be used on hard nonporous surfaces in commercial institutional hospital and premises including kitchens, bathrooms, sick rooms, laundry rooms, eating establishments, pet kennels, and veterinary premises. To disinfect hard nonporous surfaces first clean surface by removing gross filth (loose dirt debris, food material, etc). Prepare a 5,382 ppm available chlorine solution. Thoroughly wet surface with the solution and allow it to remain in contact with the surface for 4 minutes. Allow to air dry. Prepare fresh solution weekly when using in closed containers (spray bottles). Prepare fresh solution daily when using open containers (buckets) or if solution becomes diluted. All treated equipment that will contact food, feed, or drinking water must be rinsed with potable water before reuse.

DILUTION CHART		
3.3 Gram Tablet Solution ppm (mg/L) Available Chlorine	Tablets	Quarts of water
538	1	2
1,076	1	1
2,153	2	1
4,306	4	1
5,382	5	1



**C. diff Solution Tablets**

- Broad spectrum disinfectant
- Kills *Clostridium difficile* spores in 4 minutes
- Effective against Hepatitis A and Hepatitis B
- Easy, Convenient, Accurate Measurement
- Surface Friendly
- Smellier better than bleach
- Stable in solution (as compared to bleach)
- Can be applied spray, mop, wipe, cloth, sponge, brush, foaming equipment, or coarse trigger sprayer.
- 3M™ C. diff Solution Tablets is effective against the following micro-organisms on pre-cleaned, hard, nonporous, inanimate surfaces: *Salmonella enterica*, *Staphylococcus aureus*, *Pseudomonas aeruginosa*, *Klebsiella pneumoniae*, *Staphylococcus epidermidis*, *Escherichia coli* O157:H7, *Staphylococcus aureus*-methicillin-resistant (MRSA) & glycopeptide-resistant (GRSA), carbapenem resistant *Klebsiella pneumoniae*, *Acinetobacter baumannii*, Poliovirus Type1, Herpes Simplex Virus type 1, Hepatitis A Virus, Hepatitis B Virus, Human Immunodeficiency Virus Type 1 (associated with AIDS or AIDS Virus), Influenza virus H1N1, Respiratory Syncytial Virus, Canine Parvovirus, Newcastle Disease Virus, Pseudorabies, Canine Distemper Virus, Feline Calicivirus, Norovirus, *Enterococcus faecalis* Vancomycin Resistant, *Trichophyton mentagrophytes*, *Mycobacterium bovis* (TB), and *Clostridium difficile*.
- 3M™ C. diff Solution Tablets is designed to provide effective cleaning, and disinfection in areas where it is of prime importance in controlling the hazard of cross contamination on treated pre-cleaned, hard, nonporous, inanimate surfaces.

6.38

 <b>Requester:</b> Colleen Vossen <b>Creator:</b> deZinnia_19366 <b>File Name:</b> 34871855903.ai <b>Structure #:</b> SS-71909 <b>Date:</b> 05/26/16  <b>Scale:</b> 1 Inch	<b>Printed Colors – Front:</b> 	<b>This artwork has been created as requested by 3M. 3M is responsible for the artwork AS APPROVED and assumes full responsibility for its correctness.</b>
	<b>Printed Colors – Back:</b>	
<b>Match Colors:</b>		