



STATE OF HAWAII  
Department of Agriculture

**ACCEPTED**

LICENSE NO.

**9786.143**

# PropiMax<sup>®</sup>

## EC

### Fungicide

For control of certain diseases in almond, berries, carrot, celery, cereal grains, cilantro (coriander) leaves, corn, cranberry, garden beet, leaf petioles, nonbearing citrus fruits, onions, parsley leaves (fresh and dried), peanut, pecan, peppermint, pineapple, rice, sorghum, spearmint, stone fruits, sugarcane, tree nuts, turfgrass grown for seed, turfgrass and ornamentals, wild rice, and nonbearing fruit and nut trees in nurseries and landscape settings.

Group	3	FUNGICIDE
-------	---	-----------

Active Ingredient:

propiconazole: 1-[[2-(2,4-dichlorophenyl)-4-propyl-1,3-dioxolan-2-yl]methyl]-1H-1,2,4-triazole ..... 41.8%

Other Ingredients ..... 58.2%  
Total ..... 100.0%

Contains petroleum distillates.

Contains 3.6 lb active ingredient per gallon.

## Keep Out of Reach of Children WARNING AVISO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

### Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. Refer to the label booklet under "Agricultural Use Requirements" in the Directions for Use section for information about this standard.

For additional Precautionary Statements, First Aid, Storage and Disposal and other use information see inside this label.

**Notice:** Read the entire label before using. Use only according to label directions. **Before using this product, read Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies at end of label booklet. If terms are unacceptable, return at once unopened.**

In case of emergency endangering health or the environment involving this product, call 1-800-992-5994.

Agricultural Chemical: Do not ship or store with food, feeds, drugs or clothing.

EPA Reg. No. 62719-346

EPA Est. 34704-MS-2  
900-018842 / 00303908

®Trademark of Dow AgroSciences LLC

Produced for  
Dow AgroSciences LLC  
9330 Zionsville Road  
Indianapolis, IN 46268

Net Contents **1 gal**



Table of Contents	Page
First Aid .....	1
Precautionary Statements.....	1
Hazards to Humans and Domestic Animals .....	1
Personal Protective Equipment (PPE).....	1
User Safety Requirements.....	2
User Safety Recommendations.....	2
Engineering Controls.....	2
Environmental Hazards .....	2
Physical or Chemical Hazards.....	2
Directions for Use .....	2
Agricultural Use Requirements.....	2
Storage and Disposal .....	2
Crops .....	3
Product Information .....	3
Integrated Pest Management .....	3
Fungicide Resistance Management .....	3
Spray Equipment.....	3
Mixing Directions.....	4
Application Directions .....	4
Rotational Crops .....	7
Uses.....	7
Almond .....	7
Bulb Onion (Subgroup 3-07A) <sup>1</sup> and Green Onion (Subgroup 3-07B) <sup>2</sup> .....	8
Bushberries (Subgroup 13-07B) <sup>1</sup> , Caneberries (Subgroup 13-07A) <sup>2</sup> , and Low Growing Berries (Subgroup 13-07G) (Except Cranberry) <sup>3</sup> .....	9
Carrot .....	11
Cereal Grains (Crop Group 15) .....	12
Corn (Field, Sweet, Pop, and Seed) .....	14
Cranberry .....	15
Garden Beet.....	15
Leaf Petioles (Subgroup 4B).....	16
Parsley Leaves (Fresh and Dried) and Cilantro (Coriander) Leaves .....	16
Peanut.....	17
Pecan .....	19
Peppermint and Spearmint .....	20
Pineapple .....	20
Rice .....	21
Sorghum.....	23
Stone Fruits (Crop Group 12) .....	24
Sugarcane .....	26
Tree Nuts (Crop Group 14) .....	27
Turfgrass Grown for Seed.....	28
Wheat.....	29
Terms and Conditions of Use.....	30
Warranty Disclaimer.....	30
Inherent Risks of Use.....	30
Limitation of Remedies .....	30

## First Aid

**If in eyes:** Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

**If on skin or clothing:** Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

**If swallowed:** Immediately call a poison control center or doctor. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give **any** liquid to the person. Do not give anything by mouth to an unconscious person.

**If inhaled:** Remove person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. Call a poison control center or doctor for further treatment advice.

Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact 1-800-992-5994 for emergency medical treatment information.

**Note to Physician:** Contains petroleum distillate – vomiting may cause aspiration pneumonia.

## Precautionary Statements

### Hazards to Humans and Domestic Animals

# WARNING

**Causes Substantial But Temporary Eye Injury • Harmful If Swallowed, Inhaled, Or Absorbed Through Skin • Prolonged Or Frequently Repeated Skin Contact May Cause Allergic Reactions In Some Individuals**

**Do not get in eyes or on clothing. Avoid contact with skin. Avoid breathing spray mist.**

### Personal Protective Equipment (PPE)

Some materials that are chemical resistant to this product are listed below. If you want more options, follow the instructions for Category G on the EPA chemical resistance category selection chart.

**Applicators and other handlers must wear:**

- Long-sleeved shirt and long pants
- Chemical-resistant gloves, such as barrier laminate or viton
- Shoes plus socks
- Protective eyewear

**User Safety Requirements**

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

**User Safety Recommendations**

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

**Engineering Controls**

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 160.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

**Environmental Hazards**

This pesticide is toxic to fish. Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters or rinsate.

**Physical or Chemical Hazards**

Do not use, pour, spill or store near heat or open flame.

**Directions for Use**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Read all Directions for Use carefully before applying.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your state or tribe, consult the agency responsible for pesticide regulation.

**Agricultural Use Requirements**

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours. Exception: If the product is soil-injected or soil-incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is:

- Coveralls
- Chemical-resistant gloves, such as barrier laminate or viton
- Shoes plus socks
- Protective eyewear

**Storage and Disposal**

Do not contaminate water, food, or feed by storage or disposal.

**Pesticide Storage:** Store in original container in secured dry storage area. Prevent cross-contamination with other pesticides and fertilizers. For minor spills, leaks, etc., follow all precautions indicated on this label and clean up immediately.

Take special care to avoid contamination of equipment and facilities during cleanup and disposal of wastes.

## Storage and Disposal (Cont.)

**Pesticide Disposal:** Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.

**Container Handling:** Nonrefillable container. Do not reuse or refill this container.

Triple rinse or pressure rinse container (or equivalent) promptly after emptying. **Triple rinse** as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. **Pressure rinse** as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 psi for at least 30 seconds. Drain for 10 seconds after the flow begins to drip. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures allowed by state and local authorities.

## Crops

### Product Information

PropiMax<sup>®</sup> EC fungicide is a broad-spectrum fungicide for the control of certain diseases in listed crops.

**Important:** Do not use in greenhouses or as a tree injection.

**Failure to follow directions and precautions on this label may result in crop injury, poor disease control, or illegal residues.**

### Integrated Pest Management

Integrate this product into an overall disease and pest management (IPM) strategy whenever the use of a fungicide is required. Follow cultural practices known to reduce disease development. Consult

your local agricultural authorities for additional IPM strategies established for your area. PropiMax EC may be used in state agricultural extension advisory (disease forecasting) programs, the specified application timing based upon environmental factors favorable for disease development.

### Fungicide Resistance Management

PropiMax EC belongs to the sterol demethylation inhibitor (DMI) class of fungicides and is classified as Group 3 Fungicide by EPA. Since certain fungi can develop resistance to this class of products, use PropiMax EC as part of a resistance management strategy that includes alternation and/or tank mixing with another fungicide mode of action. After two consecutive applications of PropiMax EC, another propiconazole product, or another DMI, rotate to a product that is effective on the target pathogen and has a mode of action different from PropiMax EC. Apply the alternate products within the intervals specified on the label for PropiMax EC. Do not apply PropiMax EC at rates below those specified on the label. If tank mixing, use the full label rate of PropiMax EC with the full label rates of other products effective on the target pest. Consult your local or state agricultural authorities for resistance management strategies that are appropriate for your disease management program.

### Spray Equipment

Thorough coverage is necessary to provide good disease control.

To avoid spray drift, do not apply when conditions favor drift beyond the target area. Avoid spray overlap as crop injury may occur.

Air assisted or air blast sprayers move spray droplets into the canopy using a forced air stream. Set up the fan to deliver only enough air volume to penetrate the canopy and provide good coverage. Adjust deflectors or other aiming devices to direct spray only to the target area.

Equip sprayers with nozzles that provide accurate and uniform application. Be certain that nozzles are the same size and uniformly spaced across the boom. Calibrate the sprayer before use.

Use a pump with sufficient capacity to maintain 35 to 40 psi at nozzles and provide sufficient agitation in the tank to keep the mixture in suspension (this requires recirculation of 10% of tank volume per minute). Use a jet agitator or liquid sparger tube for agitation. Do not use air sparging.

Although PropiMax EC is an emulsifiable concentrate, a best practice is to use screens to protect the pump and to prevent nozzles from clogging. Screens placed on the suction side of the pump should be 16-mesh or coarser. Do not place a screen in the recirculation line. Use 50-mesh or coarser screens between the pump and boom and, where required, at the nozzles. Check nozzle manufacturer's directions.

For more information on spray equipment and calibration, consult sprayer manufacturers and state recommendations. For specific local directions and spray schedules, consult the current state agricultural experiment station recommendations.

## Mixing Directions

Application Rate Reference Table

Application Rate of PropiMax EC (fl oz/acre)	Active Ingredient Equivalent (lb ai/acre)	Acres per Gallon of PropiMax EC
2	0.056	64
4	0.1125	32
6	0.169	21.3
8	0.225	16
10	0.28	12.8
12	0.34	10.7
16	0.45	8
20	0.56	6.4
24	0.67	5.3
30	0.84	4.3
32	0.9	4

Prepare no more spray mixture than is required for the immediate application. Thoroughly clean spray equipment before using this product. Agitate the spray solution before and during application. Rinse spray tank thoroughly with clean water after each day's use and dispose of pesticide rinsate by application to an already treated area.

**PropiMax EC - Alone:** Add 1/2 to 2/3 of the required amount of water to the spray or mixing tank. With the agitator running, add PropiMax EC to the tank. Continue agitation while adding the remainder of the water. Begin application of the spray solution after

PropiMax EC has completely dispersed into the mix water. Maintain agitation until all of the mixture has been sprayed.

**PropiMax EC - Tank Mix:** PropiMax EC is usually compatible with all tank mix partners listed on this label. To determine the physical compatibility of PropiMax EC with other products, use a jar test. Using a quart jar, add the proportionate amounts of the products to 1 quart of water. Add wettable powders and water dispersible granular products first, then liquid flowables and emulsifiable concentrates last. After thoroughly mixing, let stand for at least 5 minutes. If the combination remains mixed or can be readily remixed, it is physically compatible. Once compatibility has been proven, use the same procedure for adding required ingredients to the spray tank. Add 1/2 to 2/3 of the required amount of water to the spray or mixing tank. With the agitator running, add the tank mix partner into the tank. Allow the material to completely dissolve and disperse into the mix water. Continue agitation while adding the remainder of the water and PropiMax EC to the spray tank. Allow PropiMax EC to completely disperse. Spray the mixture with the agitator running.

Do not apply this product in a tank mix with a dodine fungicide or crop injury may occur.

If using PropiMax EC in a tank mix, observe all directions for use, crops/sites, use rates, dilution ratios, precautions, and limitations required by the label of the tank mix product. Do not exceed any label dosage rate and the most restrictive label precautions and limitations must be followed. Do not tank mix this product with any product that prohibits such mixing. Tank mixes or other applications of products referenced on this label are permitted only in those states in which the referenced products are registered.

## Application Directions

PropiMax EC is most effective when applied and allowed to dry before a rainfall. Avoid applying PropiMax EC under conditions when uniform coverage cannot be obtained or when excessive spray drift may occur. Do not apply in a manner that results in exposure to humans or animals.

### Ground Application

For tree crops, apply PropiMax EC in a minimum of 50 gallons of water per acre (gpa) unless otherwise specified. For all other crops, apply PropiMax EC in a minimum of 10 gpa unless otherwise specified.

### **Aerial Application**

For tree crops, apply PropiMax EC in a minimum of 10 gpa unless otherwise specified. For all other crops, apply PropiMax EC in a minimum of 2 gpa unless otherwise specified.

### **Chemigation Application**

This product may be applied through properly equipped chemigation systems for disease control in the labeled crops. Refer to crop specific use directions for application rates, timing and frequency of application. Do not apply PropiMax EC by chemigation to other labeled crops except as specified in Dow AgroSciences supplemental labeling.

**Directions for Sprinkler Chemigation:** Apply this product only through center pivot, solid set, hand move, or moving wheel irrigation systems. Do not apply this product through any other type of irrigation system. Use only with drive systems that provide uniform water distribution.

When applying this product by chemigation, do not exceed labeled rates or apply more frequently than specified for conventional application methods. PropiMax EC, alone or in combination with other pesticides that are registered for application through irrigation systems, may be applied through irrigation systems. For chemigation application to labeled crops, apply in 0.1 to 0.25 inches of water unless otherwise specified. Chemigation with excessive water may lead to a decrease in efficacy.

**Note:** Do not inject PropiMax EC at full strength or deterioration of valves and seals may occur. Use a dilution ratio of at least 10 parts water to 1 part PropiMax EC. PropiMax EC is corrosive to many seal materials. Leather seals are best. EPDM or silicone rubber seals can be used but should be replaced once a year. Do not use viton, Buna-N, neoprene, or PVC seals.

**Chemigation Equipment Preparation:** The following use directions are to be followed when this product is applied through sprinkler irrigation systems. Thoroughly clean the chemigation system and tank of any fertilizer or chemical residues, and dispose of the residues according to state and federal laws. Flush the injection system with soap or a cleaning agent and water. Determine the amount of this product needed to cover the desired area. Mix according to instructions in the Mixing Directions section. Continually agitate the mixture during mixing and application.

**Center Pivot Equipment Calibration:** In order to calibrate the irrigation system and injector to apply

the mixture containing this product, determine the following: 1) Determine size of area to be treated; 2) Determine the time required to apply 1/8 to 1/2 inch of water over the area to be treated when the system and injection equipment are operated at normal pressures as specified by the equipment manufacturer. When applying PropiMax EC through irrigation equipment, use the lowest obtainable water volume while maintaining uniform distribution. Run the system at 80 to 95% of the manufacturer's rated capacity.; 3) Using only water, determine the injection pump output when operated at normal line pressure; 4) Determine the amount of PropiMax EC required to treat the area covered by the irrigation system; 5) Add the required amount of PropiMax EC and sufficient water to meet the injection time requirements of the solution tank. Maintain constant solution tank agitation during the injection period. Operate system at normal pressures specified by the manufacturer of the injection equipment and used for the time interval established during calibration. Inject this product at the end of an irrigation cycle or as a separate application to maximize foliar absorption and retention. Stop injection equipment after treatment is completed. Continue to operate the system until the solution with this product has cleared the last sprinkler head. Do not use end guns when applying PropiMax EC through center pivot systems because of non-uniform application.

### **Solid Set, Hand Move, and Moving Wheel**

**Equipment Calibration:** In order to calibrate the irrigation system and injector to apply the mixture containing this product, determine the following: 1) Determine the acreage covered by the sprinkler; 2) Fill the injector solution tank with water and adjust flow rate to use the contents over a 20- to 30-minute interval. When applying PropiMax EC through irrigation equipment, use the lowest obtainable water volume while maintaining uniform distribution.; 3) Determine the amount of PropiMax EC required to treat the area covered by the irrigation system; 4) Add the required amount of PropiMax EC into the same quantity of water used to calibrate the injection equipment. Maintain constant solution tank agitation during the injection period. Operate the system at normal pressures specified by the manufacturer of the injection equipment and used for the time interval established during the calibration. Inject PropiMax EC at the end of the irrigation cycle or as a separate application to maximize foliar fungicide retention. Stop injection equipment after treatment is completed. Continue to operate the system until the solution of PropiMax EC has cleared the last sprinkler head.

**Chemigation Operation:** Start the water pump and irrigation system, and let the system achieve the desired pressure and speed before starting the injector. Make sure the system is fully charged with water before starting injection of PropiMax EC. Time the injection to last at least as long as it takes to bring the system to full pressure. Check for leaks and uniformity and make repairs before any chemigation takes place. Start the injection system and calibrate according to manufacturer's specifications. This procedure is necessary to deliver the desired rate per acre in a uniform manner. When the application is finished, allow the entire irrigation and injection system to be thoroughly flushed clean before stopping the system.

**Chemigation Equipment Requirements:**

- The system must contain an air gap, an approved backflow prevention device, a functional check valve, vacuum relief valve (including inspection port), and low-pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from back flow. Refer to the American Society of Agricultural Engineer's Engineering Practice 409 for more information or state specific regulations.
  - The pesticide injection pipeline must contain a functional, automatic, quick-closing check-valve to prevent the flow of fluid back toward the injection pump.
  - The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
  - The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.
  - Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.
  - To ensure uniform mixing of the fungicide in the water line, inject the mixture in the center of the pipe diameter or just ahead of an elbow or tee in the irrigation line so that the turbulence created at those points will assist in mixing. The injection point must be located after all backflow prevention devices on the water line.
- Ensure the tank holding the fungicide mixture is free of rust, fertilizer, sediment, and foreign material, and equipped with an in-line strainer situated between the tank and the injector point.

**Chemigation Precautions:**

- Crop injury, lack of effectiveness, or illegal pesticide residues in the crop can result from non-uniform distribution of treated water.
- If you have questions about calibration, contact State Extension Service specialists, equipment manufacturers, or other experts.
- Public water system means a system for the provision to the public of piped water for human consumption if such system has at least 15 service connections or regularly serves an average of at least 25 individuals daily at least 60 days out of the year.
- A person knowledgeable of the chemigation system and responsible for its operation, or under the supervision of the responsible person, shall operate the system and make necessary adjustments and continuously monitor the injection.

**Chemigation Restrictions:**

- Chemigation systems connected to public water systems must contain a functional, reduced-pressure zone (RPZ), back flow preventer or the functional equivalent in the water supply line upstream from the point of pesticide introduction. As an option to the RPZ, discharge the water from the public water system into a reservoir tank prior to pesticide introduction. There shall be a complete physical break (air gap) between the flow outlet end of the fill pipe and the top or overflow rim of the reservoir tank of at least twice the inside diameter of the fill pipe.
- The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection.
- The pesticide injection pipeline must contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
- The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops, or in cases where there is no water pump, when the water pressure decreases to the point where pesticide distribution is adversely affected.

- Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.
- Do not connect an irrigation system used for pesticide application (including greenhouse systems) to a public water system unless the pesticide label-prescribed safety devices for public water systems are in place with current certification. Specific local regulations may apply and must be followed.
- Do not apply when wind speed favors drift beyond the area intended for treatment. End guns must be turned off during the application if they irrigate nontarget areas.
- Do not allow irrigation water to collect or run off and pose a hazard to livestock, wells, or adjoining crops.
- Do not enter treated area during the reentry interval specified in the Agricultural Use Requirements section of this label unless required PPE is worn.
- Do not apply through sprinkler systems that deliver a low coefficient of uniformity such as certain water drive units.

### Rotational Crops

To avoid possible illegal residues, do not plant any other crop intended for food, grazing, or any component of animal feed or bedding within 105 days of an application of PropiMax EC to the preceding crop unless the second crop appears on this label. Alfalfa can be planted 75 days after the last application of PropiMax EC if the total application of propiconazole has not exceeded 0.22 lb active ingredient per acre during the previous year.

## Uses

### Almond

Apply PropiMax EC by either ground or aerial equipment in a minimum of 15 gpa.

Target Disease		Rate (fl oz/acre)	Application Timing	Use Restrictions
Common Name	Scientific Name			
brown rot blossom blight	<i>Monilinia laxa</i> , <i>M. fructicola</i>	4 – 8	Apply at 5 to 10% bloom and 50 to 100% bloom. Under severe disease conditions, use the highest rate. The minimum re-treatment interval is 7 days.	<ul style="list-style-type: none"> <li>• Do not apply more than a total of 32 fl oz of PropiMax EC (0.9 lb active ingredient) per acre per season.</li> <li>• <b>Preharvest Interval:</b> Do not apply within 60 days of harvest.</li> <li>• Do not graze livestock in treated areas or cut treated cover crop for feed.</li> </ul>
anthracnose	<i>Collectotrichum acutatum</i>	8	Apply beginning at bud break on a 7- to 14-day interval.	

## Bulb Onion (Subgroup 3-07A)<sup>1</sup> and Green Onion (Subgroup 3-07B)<sup>2</sup>

<sup>1</sup>Bulb onion (subgroup 3-07A) including bulb daylily, bulb fritillaria, bulb garlic, bulb lily, bulb onion, bulb shallot, Chinese bulb onion, great-headed bulb garlic, pearl onion, potato bulb onion, serpent bulb garlic, cultivars, varieties, and/or hybrids of these

<sup>2</sup>Green onion (subgroup 3-07B) including beltville bunching onion, Chinese fresh leaf chive, elegans hosta, fresh leaf chive, fresh leaf shallot, fresh onion, green onion, leaf fritillaria, kurrat, lady's leek, leek, macrostem onion, tree top onion, Welsh tops onion, wild leek, cultivars, varieties, and/or hybrids of these

Apply PropiMax EC by either ground equipment in a minimum of 15 gpa or aerial equipment in a minimum of 5 gpa.

Target Disease		Rate (fl oz/acre)	Application Timing	Use Restrictions
Common Name	Scientific Name			
botrytis leaf blight (suppression only) purple blotch	<i>Botrytis squamosa</i>	4 - 8	Begin applications when conditions favor disease development and reapply on a 7- to 10-day interval. Use the higher rate and shorter interval when disease conditions are severe.	<ul style="list-style-type: none"> <li>Do not apply more than a total of 16 fl oz of PropiMax EC (0.45 lb active ingredient) per acre per season.</li> <li><b>Preharvest Interval:</b> Bulb onion-Do not apply within 14 days of harvest. There is no preharvest interval for green onion types.</li> </ul>
	<i>Alternaria pom</i>	2 - 4	Apply with another fungicide registered for control of target diseases. Begin applications when conditions favor disease development and reapply on a 7-day interval or according to the tank mix partner label. Use the higher rate when disease conditions are severe. To achieve optimum disease control, a wetting agent or spreader-sticker may be used.	

## Bushberries (Subgroup 13-07B)<sup>1</sup>, Caneberries (Subgroup 13-07A)<sup>2</sup>, and Low Growing Berries (Subgroup 13-07G) (Except Cranberry)<sup>3</sup>

<sup>1</sup>Bushberries (subgroup 13-07B) including aronia berry, black currant, buffalo currant, Chilean guava, edible honeysuckle, elderberry, European barberry, gooseberry, highbush blueberry, highbush cranberry, huckleberry, jostaberry, juneberry, lingonberry, lowbush blueberry, native currant, red currant, salal, sea buckthorn, cultivars, varieties, and/or hybrids of these

<sup>2</sup>Caneberries (subgroup 13-07A) including bingleberry, blackberry, black raspberry, boysenberry, dewberry, loganberry, lowberry, marionberry, olallieberry, red raspberry, wild raspberry, youngberry, cultivars, varieties and/or hybrids of these

<sup>3</sup>Low growing berries (subgroup 13-07G) except cranberry) including bearberry, bilberry, lowbush blueberry, cloudberry, lingonberry, muntries, partridgeberry, strawberry, cultivars, varieties, and/or hybrids of these

### Bushberries (Subgroup 13-07B) and Caneberries (Subgroup 13-07A)

Apply PropiMax EC by either ground equipment in a minimum of 15 gpa or aerial equipment in a minimum of 5 gpa.

Target Disease		Rate (fl oz/acre)	Application Timing	Use Restrictions
Common Name	Scientific Name			
leaf spot, stem canker rust	<i>Septoria albopuncata</i>  <i>Pucciniastrum vaccinii</i>	6	Apply when conditions favor disease development. Reapply on a 4-week interval.	<ul style="list-style-type: none"> <li>Do not apply more than a total of 30 fl oz of PropiMax EC (0.84 lb active ingredient) per acre per season.</li> <li><b>Preharvest Interval:</b> Do not apply within 30 days of harvest.</li> </ul>
leaf spot	<i>Septoria</i> spp.		Make the first application any time prior to bloom and again after petal fall. If needed, reapply just after harvest.	
leaf and cane spot	<i>Septoria rubi</i>		Apply as a delayed dormant spray after training in the spring. Reapply in late spring, at bud break, and once flowering has begun.	
mummyberry disease	<i>Monilinia vaccinicorymbosi</i>		Make first application beginning at green tip and repeat in 7 to 10 days. If conditions are favorable for disease development, reapply at pink bud and every 7 to 10 days through petal fall.	

Target Disease (Cont.)		Rate (fl oz/acre)	Application Timing	Use Restrictions
Common Name	Scientific Name			
powdery mildew	<i>Microsphaera vaccinii</i>	6	Apply at 5 to 10% bloom. Reapply at full bloom and on a 14-day interval while conditions favor disease development.	<ul style="list-style-type: none"> <li>Do not apply more than a total of 30 fl oz of PropiMax EC (0.84 lb active ingredient) per acre per season.</li> <li><b>Preharvest Interval:</b> Do not apply within 30 days of harvest.</li> </ul>

#### Low Growing Berries (Subgroup 13-07G) (Except Cranberry)

Apply PropiMax EC by either ground equipment in a minimum of 20 gpa or aerial equipment in a minimum of 15 gpa.

Target Disease		Rate (fl oz/acre)	Application Timing	Use Restrictions
Common Name	Scientific Name			
anthracnose leaf rust leaf spot powdery mildew	<i>Colletotrichum acutatum</i> <i>Phragmidium potentillae</i> <i>Cercospora fragariae</i> <i>Sphaerotheca maculeris</i>	4	Apply when disease levels are no more than 5%. Apply up to 4 times on a 7-day interval. Do not make more than 2 consecutive applications of this product before rotating to another fungicide with a different mode of action.	<ul style="list-style-type: none"> <li>Do not apply more than a total of 16 fl oz of PropiMax EC (0.45 lb active ingredient) per acre per season.</li> <li><b>Preharvest Interval:</b> There is no preharvest interval</li> </ul>

## Carrot

Apply PropiMax EC by either ground equipment in a minimum of 15 gpa or aerial equipment in a minimum of 5 gpa.

Target Disease		Rate (fl oz/acre)	Application Timing	Use Restrictions
Common Name	Scientific Name			
<i>Alternaria dauci</i> (suppression only) leaf blight powdery mildew	<i>Cercospora carotae</i> <i>Erysiphe polygoni</i>	4	Apply when conditions favor disease development. Reapply on a 7- to 10-day interval using the shorter interval when disease conditions are severe. If desired, a spreader-sticker may be used.	<ul style="list-style-type: none"> <li>Do not apply more than a total of 16 fl oz of PropiMax EC (0.45 lb active ingredient) per acre per season.</li> <li><b>Preharvest Interval:</b> Do not apply within 14 days of harvest.</li> </ul>
		2	Apply the product with 0.75 lb active ingredient of chlorothalonil per acre. Begin applications when conditions favor disease development. Reapply on a 7- to 10-day interval.	

## Cereal Grains (Crop Group 15)<sup>1</sup>

<sup>1</sup>Cereal grains (crop group 15) including barley, buckwheat, corn, pearl millet, proso millet, oats, popcorn, rice, rye, sorghum, teosinte, triticale, wheat, wild rice and other commodities included in the cereal grains crop grouping. **For corn, popcorn, rice, sorghum, wheat, and wild rice, see specific use directions as designated on this label.**

Apply PropiMax EC by either ground, aerial, or chemigation equipment.

Target Disease		Rate (fl oz/acre)	Application Timing	Use Restrictions
Common Name	Scientific Name			
<b>early season suppression of:</b> glume blotch  leaf blight powdery mildew  tan spot	<i>Stagonospora nordorum</i> <i>Septoria tritici</i> <i>Blumeria</i> spp., <i>Erysiphe</i> spp. <i>Pyrenophora tritici-repentis</i>	2 – 4	Apply in the spring. Reapply through full head emergence for season long control. The minimum re-treatment interval is 14 days.	<ul style="list-style-type: none"> <li>• Do not apply more than a total of 8 fl oz of PropiMax EC (0.22 lb active ingredient) per acre per season.</li> <li>• Do not apply more than a total of 4 fl oz of PropiMax EC (0.11 lb active ingredient) per acre per season if forage or hay will be harvested.</li> <li>• Do not apply after Feekes 10.5 in wheat.</li> <li>• <b>Preharvest Interval:</b>  <b>All cereals-</b>Do not apply within 30 days of harvest for forage or hay. <b>Cereals other than wheat-</b>Do not apply within 45 days of harvest for grain and straw.</li> </ul>

Target Disease (Cont.)		Rate (fl oz/acre)	Application Timing	Use Restrictions
Common Name	Scientific Name			
<b>control of:</b> barley scald  barley stripe  glume blotch  helminthosporium leaf blight leaf blight net blotch powdery mildew  rust spot blotch tan spot	<i>Rhynchosporium secalis</i> <i>Pyrenophora graminea</i> <i>Stagonospora nodorum</i> <i>Drechslera tritici-repentis</i> <i>Septoria tritici</i> <i>Pyrenophora teres</i> <i>Blumeria</i> spp., <i>Erysiphe</i> spp. <i>Puccinia</i> spp. <i>Bipolaris sorokiniana</i> <i>Pyrenophora tritici-repentis</i>	4	Protecting the flag leaf is important for maximizing yield. Highest yields are normally obtained when the product is applied when the flag leaf is 50% to fully emerged.  The minimum re-treatment interval is 14 days. Using an oil base adjuvant may improve the spray coverage and canopy penetration.  In wheat, the product can be applied through full head emergence (Feekes growth stage 10.5). Do not apply after this stage to avoid possible illegal residues.	Do not apply more than a total of 8 fl oz of PropiMax EC (0.22 lb active ingredient) per acre per season. <ul style="list-style-type: none"> <li>Do not apply more than a total of 4 fl oz of PropiMax EC (0.11 lb active ingredient) per acre per season if forage or hay will be harvested.</li> <li>Do not apply after Feekes 10.5 in wheat.</li> <li><b>Preharvest Interval:</b> <b>All cereals-</b>Do not apply within 30 days of harvest for forage or hay. <b>Cereals other than wheat-</b>Do not apply within 45 days of harvest for grain and straw.</li> </ul>
foot rot	<i>Pseudocercospora</i> spp.		Apply with half rates of fungicides such as thiophanate-methyl. Apply at tillering but before elongation has occurred.	
fusarium head blight suppression			Apply at approximately 50% flowering. Adding a penetrating type of adjuvant may increase fusarium head blight suppression.	

## Corn (Field, Sweet, Pop, and Seed)

Apply PropiMax EC by either ground, aerial, or chemigation equipment.

Target Disease		Rate (fl oz/acre)	Application Timing	Use Restrictions
Common Name	Scientific Name			
helminthosporium leaf blights  northern corn leaf blight northern corn leaf spot southern corn leaf blight	<i>H. maydis</i> , <i>H. turcicum</i> , <i>H. carbonum</i> <i>Setosphaeria turcica</i>  <i>Cochliobolus carbonum</i> <i>Cochliobolus heterostrophus</i>	2 – 4	Apply when disease first appears and reapply on a 7- to 14-day schedule. Use the low rate when disease pressure is low and the high rate under heavy disease pressure or if conditions favor disease development.	<b>Field corn, popcorn, seed corn</b> <ul style="list-style-type: none"> <li>Do not apply more than a total of 16 fl oz of PropiMax EC (0.45 lb active ingredient) per acre per season.</li> <li>Do not apply more than a total of 8 fl oz of PropiMax EC (0.22 lb active ingredient) per acre per season on field corn harvested for forage.</li> </ul>
eye spot gray leaf spot  rusts	<i>Aureobasidium zeae</i> <i>Cercospora zeaemaydis</i> <i>Puccinia</i> spp.	4	Apply when disease first appears. If conditions favor disease development, reapply on a 7- to 14-day schedule. For best disease control, early applications at initial disease onset perform better.	<ul style="list-style-type: none"> <li><b>Preharvest Interval:</b> Do not apply within 30 days of harvest for forage, grain and stover.</li> <li><b>Sweet corn</b></li> <li><b>Preharvest Interval:</b> Do not apply within 14 days of harvest for ears or forage.</li> </ul>

## Cranberry

(For use only in Oregon, Washington and Wisconsin)

Apply PropiMax EC by either ground or aerial equipment in a minimum of 20 gpa.

Target Disease		Rate (fl oz/acre)	Application Timing	Use Restrictions
Common Name	Scientific Name			
cottonball	<i>Monilinia oxycocci</i>	4 – 6	Make the first application at leaf bud break and repeat in 14 days. Make the third application at early bloom and repeat in 14 days. Under severe disease pressure, use the higher rate for control.	<ul style="list-style-type: none"><li>• Do not apply more than a total of 24 fl oz of PropiMax EC (0.67 lb active ingredient) per acre per season.</li><li>• <b>Preharvest Interval:</b> Do not apply within 45 days of harvest.</li></ul>

## Garden Beet

Apply PropiMax EC by either ground, aerial, or chemigation equipment.

Target Disease		Rate (fl oz/acre)	Application Timing	Use Restrictions
Common Name	Scientific Name			
leaf spot powdery mildew	<i>Cercospora beticola</i> <i>Erysiphe polygoni</i>	3 - 4	Apply when disease first appears and reapply on a 14-day interval. Do not make more than 2 consecutive applications of this product before rotating to another fungicide with a different mode of action. Immediately switch to a fungicide that has a different mode of action if disease levels continue to increase.	<ul style="list-style-type: none"><li>• Do not apply more than a total of 16 fl oz of PropiMax EC (0.45 lb active ingredient) per acre per season.</li><li>• <b>Preharvest Interval:</b> Do not apply within 14 days of harvest.</li></ul>

## Leaf Petioles (Subgroup 4B)<sup>1</sup>

<sup>1</sup>Leaf petioles (subgroup 4B) including cardoon, celery, Chinese celery, celtuce, Florence fennel, rhubarb, Swiss chard

Apply PropiMax EC by either ground equipment in a minimum of 10 gpa or aerial equipment in a minimum of 5 gpa.

Target Disease		Rate (fl oz/acre)	Application Timing	Use Restrictions
Common Name	Scientific Name			
early blight late blight leaf spot	<i>Cercospora apii</i> <i>Septoria apicola</i> <i>Cercospora</i> spp. <i>Septoria</i> spp.	4	Apply on a 7-day schedule. If desired, a spreader-sticker may be used.	<ul style="list-style-type: none"> <li>Do not apply more than a total of 16 fl oz of PropiMax EC (0.45 lb active ingredient) per acre per season.</li> <li><b>Preharvest Interval:</b> Do not apply within 14 days of harvest.</li> </ul>

## Parsley Leaves (Fresh and Dried) and Cilantro (Coriander) Leaves

Apply PropiMax EC by either ground, aerial, or chemigation equipment.

Target Disease		Rate (fl oz/acre)	Application Timing	Use Restrictions
Common Name	Scientific Name			
leaf spot  powdery mildew	<i>Alternaria</i> spp. <i>Cercospora</i> spp. <i>Erysiphe</i> spp.	3 - 4	Apply when disease first appears and reapply on a 14-day interval. Do not make more than 2 consecutive applications of this product before rotating to another fungicide with a different mode of action.  Immediately switch to a fungicide that has a different mode of action if disease levels continue to increase.	<ul style="list-style-type: none"> <li>Do not apply more than a total of 16 fl oz of PropiMax EC (0.45 lb active ingredient) per acre per season.</li> <li><b>Preharvest Interval:</b> Do not apply within 14 days of harvest.</li> </ul>

## Peanut

Apply PropiMax EC by either ground, aerial, or chemigation equipment.

Target Disease		Rate (fl oz/acre)	Application Timing	Use Restrictions
Common Name	Scientific Name			
early leaf spot	<i>Cercospora arachidicola</i>	2.5 – 4	Apply beginning 35 to 40 days after planting or at first appearance of disease and reapply on a 10- to 14-day schedule. Under heavy disease pressure, use the higher rate.  This product may be used in State Agricultural Extension advisory (disease forecasting) programs which specify application timing based upon environmental factors favorable for disease development.	<ul style="list-style-type: none"> <li>• Do not apply more than a total of 16 fl oz of PropiMax EC (0.45 lb active ingredient) per acre per season.</li> <li>• <b>Preharvest Interval:</b> Do not apply within 14 days of harvest with the low use rate of 4 fl oz per acre. Do not apply within 21 days of harvest with the high use rate of 8 fl oz per acre.</li> <li>• Do not feed hay from treated fields to livestock if the high rate of 8 fl oz per acre was used.</li> </ul>
late leaf spot rust web blotch	<i>Cercosporidium personatum</i> <i>Puccinia arachidis</i> <i>Phoma arachidicola</i>	4		

Target Disease (Cont.)		Rate (fl oz/acre)	Application Timing	Use Restrictions
Common Name	Scientific Name			
southern stem rot	<i>Sclerotium rolfsii</i>	4	<p>Apply to the crown and pegging zones of the plant by chemigation or directed ground application using one of the following schedules:</p> <p><b>A.</b> Apply 4 fl oz per acre. Begin applications 45 days after planting or at first appearance of disease. Repeat on a 14-day interval.</p> <p><b>B.</b> Apply 8 fl oz per acre. Make the first application at pegging (approximately 60 days after planting) or at first appearance of disease. Make the second application 3 to 4 weeks later.</p> <p>When applying in irrigation water, use a minimum of 0.25 to 0.5 inches of water per acre. Use enough water so the fungicide penetrates the peanut canopy and reaches the crown of the plant where the disease is most active.</p> <p>When applying via irrigation or as a directed ground application, additional methods should be employed for leaf spot control.</p>	<ul style="list-style-type: none"> <li>• Do not apply more than a total of 16 fl oz of PropiMax EC (0.45 lb active ingredient) per acre per season.</li> <li>• <b>Preharvest Interval:</b> Do not apply within 14 days of harvest with the low use rate of 4 fl oz per acre. Do not apply within 21 days of harvest with the high use rate of 8 fl oz per acre.</li> <li>• Do not feed hay from treated fields to livestock if the high rate of 8 fl oz per acre was used.</li> </ul>

## Pecan

Apply PropiMax EC by either ground or aerial equipment in a minimum of 20 gpa.

The active ingredient in this product may have effects on federally listed threatened and endangered species or critical habitat in some counties. When using this product, you must follow the measures contained in the County Bulletin for the county in which you are applying the pesticide. To determine whether your county has a bulletin consult [www.epa.gov/espp/bulletins.htm](http://www.epa.gov/espp/bulletins.htm). Bulletins may also be available from local pesticide dealers, extension offices, or state pesticide agencies.

Target Disease		Rate (fl oz/acre)	Application Timing	Use Restrictions
Common Name	Scientific Name			
pecan scab	<i>Cladosporium caryigenum</i>	4 – 8	Apply on a 14-day schedule during bud break and pre-pollination sprays. Apply 6 to 8 fl oz per acre during nut formation and cover sprays. Use higher rates when disease pressure is heavier.	<ul style="list-style-type: none"> <li>Do not apply more than a total of 32 fl oz of PropiMax EC (0.9 lb active ingredient) per acre per season.</li> <li>Do not apply after shuck split.</li> <li>Do not graze livestock in treated areas or cut treated cover crop for feed.</li> </ul>
downy spot liver spot powdery mildew vein spot zonate leaf spot	<i>Mycosphaerella caryigena</i> <i>Gnomonia caryae</i> pv <i>pecanae</i> <i>Microsphaera penicillata</i> <i>Gnomonia nerviseda</i> <i>Cristulariella moricola</i>			

## Peppermint and Spearmint

Apply PropiMax EC by ground equipment in a minimum of 20 gpa.

Target Disease		Rate (fl oz/acre)	Application Timing	Use Restrictions
Common Name	Scientific Name			
rust	<i>Puccinia menthae</i>	4	Begin applications when the plants are 2 to 4 inches high or when conditions favor disease development. Reapply within 14 days of the first application.	<ul style="list-style-type: none"> <li>Do not apply more than a total of 8 fl oz of PropiMax EC (0.22 lb active ingredient) per acre per season.</li> <li><b>Preharvest Interval:</b> Do not apply within 30 days of harvest.</li> </ul>

## Pineapple

(For use only in Hawaii)

Target Disease		Rate (fl oz/100 gal water)	Application Timing	Use Restrictions
Common Name	Scientific Name			
butt rot	<i>Ceratocystis paradoxa</i>	0.75 (1:17,000)	<p>Treatments can be made in either a cold or hot water dip.</p> <p><b>Cold water dip:</b> Immerse crowns to give thorough wetting, remove, and allow to drain.</p> <p><b>Hot water dip:</b> Maintain water temperature at 125°F (52°C). Soak crown for 20 to 30 minutes, remove, and allow to drain.</p>	<ul style="list-style-type: none"> <li>Do not use treated crowns for food or feed.</li> <li>Do not graze while plant is growing.</li> <li>Do not graze tops until fruit is harvested.</li> <li>Dispose of used dip solution according to local, state, and federal regulations.</li> </ul>

## Rice

Apply PropiMax EC by aerial equipment only.

The active ingredient in this product may have effects on federally listed threatened and endangered species or critical habitat in some counties. When using this product, you must follow the measures contained in the County Bulletin for the county in which you are applying the pesticide. To determine whether your county has a bulletin consult [www.epa.gov/esp/bulletins.htm](http://www.epa.gov/esp/bulletins.htm). Bulletins may also be available from local pesticide dealers, extension offices, or state pesticide agencies.

Target Disease		Rate (fl oz/acre)	Application Timing	Use Restrictions
Common Name	Scientific Name			
aggregate sheath spot black sheath rot  brown leaf spot  false smut (suppression only) kernel smut leaf smut narrow brown leaf spot sheath blight sheath spot stem rot (suppression only)	<i>Rhizoctonia oryzae-sativa</i> <i>Gaeumannomyces graminis</i> <i>Helminthosporium oryzae</i> <i>Ustilagoideae virens</i>  <i>Tilletia barclayana</i> <i>Entyloma oryzae</i> <i>Cercospora oryzae</i>  <i>Rhizoctoria solani</i> <i>Rhizoctonia oryzae</i> <i>Sclerotium oryzae</i>	6 – 10	Timing of application depends upon disease severity, disease complex and rice variety/ growth stage. Consult local extension experts for economic thresholds established for various rice varieties and diseases.  Apply at first internode elongation (up to 2-inch panicle) using one of the following schedules: <b>A.</b> Apply 6 fl oz per acre and repeat at swollen boot. Make the second application 14 days later but before the boot splits and head emerges. Sheath blight is best controlled when the first application is applied at disease appearance in the field. Apply the first application when 5% or fewer of the tillers are infected. <b>B.</b> Apply 10 fl oz per acre. Use this rate if greater than 10% of the tillers are infected with sheath blight. If disease reappears, use another fungicide for the second application.  Tank mix option: Apply 6 fl oz of the product in a tank mix with registered fungicides for control of rice diseases.	<ul style="list-style-type: none"> <li>Do not apply more than a total of 12 fl oz of PropiMax EC (0.34 lb active ingredient) per acre per season.</li> <li><b>Preharvest Interval:</b> Do not apply within 35 days of harvest.</li> <li>Do not apply to stubble or ratoon crop rice.</li> <li>Do not use in rice fields where commercial farming of crayfish will be practiced.</li> <li>Do not drain water from treated rice fields into ponds used for commercial fish farming.</li> <li>Do not use water drained from treated fields to irrigate other crops.</li> <li>Do not release flood water within 7 days of application.</li> </ul>

Target Disease (Cont)		Rate (fl oz/acre)	Application Timing	Use Restrictions
Common Name	Scientific Name			
<b>Wild Rice (For use only in Minnesota)</b>				
brown spot helminthosporium leaf blight	<i>Bipolaris</i> spp. <i>Helminthosporium</i> spp.	6 – 8	Apply 6 fl oz per acre at both booting and heading, or make a single application of 8 fl oz per acre at booting. The minimum re-treatment interval is 10 days.	<ul style="list-style-type: none"> <li>• Do not apply more than a total of 12 fl oz of PropiMax EC (0.34 lb active ingredient) per acre per season.</li> <li>• <b>Preharvest Interval:</b> Do not apply within 35 days of harvest.</li> <li>• Do not apply to stubble or ratoon crop rice.</li> <li>• Do not use in rice fields where commercial farming of crayfish will be practiced.</li> <li>• Do not drain water from treated rice fields into ponds used for commercial fish farming.</li> <li>• Do not use water drained from treated fields to irrigate other crops.</li> <li>• Do not release flood water within 7 days of application.</li> </ul>

## Sorghum

Apply PropiMax EC by either ground equipment in a minimum of 15 gpa or aerial equipment in a minimum of 10 gpa.

Target Disease		Rate (fl oz/acre)	Application Timing	Use Restrictions
Common Name	Scientific Name			
ergot	<i>Claviceps sorghi</i>	3 – 4	Make the first application at or just prior to flowering. Reapply on a 5- to 7-day interval. Make up to 4 applications.	<ul style="list-style-type: none"> <li>• Do not apply more than a total of 16 fl oz of PropiMax EC (0.45 lb active ingredient) per acre per season.</li> <li>• Do not apply more than a total of 8 fl oz of PropiMax EC (0.22 lb active ingredient) per acre per season on sorghum harvested for forage.</li> <li>• <b>Preharvest Interval:</b> Do not apply within 30 days of harvest for forage and within 21 days of harvest for grain and stover.</li> <li>• Do not graze livestock or cut for green chop or silage within 30 days of application.</li> </ul>

## Stone Fruits (Crop Group 12)<sup>1</sup>

<sup>1</sup>Stone fruits (crop group 12) including apricot, chickasaw plum, damson plum, Japanese plum, nectarine, peach, plum, plumcot, prune, sweet cherry, tart cherry, and cultivars and/or hybrids of these included in the stone fruits crop grouping

Apply PropiMax EC by either ground or aerial equipment in a minimum of 15 gpa. Diseases on stone fruits are most effectively controlled by ground applications.

Target Disease		Rate (fl oz/acre)	Application Timing	Use Restrictions
Common Name	Scientific Name			
brown rot blossom blight	<i>Monilinia</i> spp.	4	Apply at early bloom stage. If disease pressure is low, a second application may be made as needed through petal fall. Under conditions of high disease pressure and/or very susceptible varieties, make a second application from 75 to 100% bloom and a third application at petal fall.	<ul style="list-style-type: none"> <li>Do not apply more than a total of 20 fl oz of PropiMax EC (0.56 lb active ingredient) per acre per season.</li> <li><b>Preharvest Interval:</b> 0 days</li> <li>Applications of PropiMax EC during bloom to stanley plums have occasionally caused fruit to be less oval in shape and smaller in size at harvest. To avoid this, do not apply to stanley plums earlier than 21 days before harvest.</li> </ul>
fruit brown rot	<i>Monilinia</i> spp.		Apply a maximum of two sprays as needed during the preharvest period up to the day of harvest. If high inoculum and severe disease conditions persist, apply another fungicide after applying PropiMax EC twice.	

Target Disease (Cont.)		Rate (fl oz/acre)	Application Timing	Use Restrictions
Common Name	Scientific Name			
cherry leaf spot powdery mildew rust	<i>Blumeriella jaapii</i> <i>Podosphaera</i> spp. <i>Tranzschelia discolor</i>	4	See brown rot blossom blight. Make up to two additional applications on a 10- to 14-day interval from the end of petal fall to harvest.	<ul style="list-style-type: none"> <li>• Do not apply more than a total of 20 fl oz of PropiMax EC (0.56 lb active ingredient) per acre per season.</li> <li>• <b>Preharvest Interval:</b> 0 days</li> <li>• Applications of PropiMax EC during bloom to stanley plums have occasionally caused fruit to be less oval in shape and smaller in size at harvest. To avoid this, do not apply to stanley plums earlier than 21 days before harvest.</li> </ul>

## Sugarcane

Target Disease		Rate (fl oz/100 gal water)	Application Timing	Use Restrictions
Common Name	Scientific Name			
pineapple disease	<i>Ceratocystic paradoxa</i>	0.75 (1:17,000)	<p>Apply to cut seed pieces using one of the following methods:</p> <p><b>Cold water dip:</b> Immerse seed pieces to thoroughly wet, remove, and allow to drain.</p> <p><b>Hot water dip:</b> Maintain water temperature at 125°F (52°C). Soak seed pieces for 20 to 30 minutes, remove, and allow to drain.</p> <p><b>Conveyor belt treatment:</b> Treat seed pieces with a solution of PropiMax EC and water using in-line directed spray sufficient to wet cut ends.</p>	<ul style="list-style-type: none"> <li>• Do not use treated seed pieces for food or feed.</li> <li>• Dispose of used dip solution according to local, state, and federal regulations.</li> </ul>

## Tree Nuts (Crop Group 14)<sup>1</sup>

<sup>1</sup>Tree nuts (crop group 14) including almond, beechnut, Brazil nut, butternut, cashew, chestnut, chinaquapin, hickory, macadamia, pecan, walnut and other commodities included in the tree nuts crop grouping. For almond and pecan, see specific use directions elsewhere on the label.

Apply PropiMax EC by either ground or aerial equipment in a minimum of 15 gpa. Diseases on tree nuts are most effectively controlled by ground applications.

Target Disease	Rate (fl oz/acre)	Application Timing	Use Restrictions
foliar diseases	4 – 8	Apply at first sign of disease and reapply on a 7- to 14-day interval.	<ul style="list-style-type: none"><li>• Do not apply more than a total of 32 fl oz of PropiMax EC (0.9 lb active ingredient) per acre per season.</li><li>• <b>Preharvest Interval:</b> Do not apply within 60 days of harvest.</li><li>• Do not graze livestock in treated areas or cut treated cover crop for feed.</li></ul>

## Turfgrass Grown for Seed

(For use only in Idaho, Minnesota, Nebraska, Oregon, and Washington)

Apply PropiMax EC by either ground or aerial equipment in a minimum of 10 gpa. PropiMax EC may also be applied by chemigation.

Target Disease		Rate (fl oz/acre)	Application Timing	Use Restrictions
Common Name	Scientific Name			
ergot stem disease powdery mildew rusts selenophoma stem eyespot	<i>Erysiphe graminis</i> <i>Puccinia</i> spp. <i>Selenophoma</i> spp.	4 – 8	Apply when powdery mildew infections, <i>Selenophoma</i> infections, and/or rust pustules are noticeable and increasing in number in late spring or early summer. Apply on a 14- to 21-day interval.  To maximize control of severe rust pressure, apply 8 fl oz per acre and reapply at 14-day intervals until the seed is mature. Make the last application at least 20 days before seed matures.  For bluegrass, it is important to begin applications early in the growing season.	<ul style="list-style-type: none"> <li>Do not apply more than a total of 32 fl oz of PropiMax EC (0.9 lb active ingredient) per acre per season.</li> <li>Do not feed hay cut within 20 days of the last application.</li> <li>Do not graze treated areas within 140 days of the last application.</li> </ul>

## Wheat

Apply PropiMax EC by either ground, aerial, or chemigation equipment.

Target Disease		Rate (fl oz/acre)	Application Timing	Use Restrictions
Common Name	Scientific Name			
<b>early season suppression of:</b> glume blotch leaf blight powdery mildew tan spot	<i>Stagonospora nordorum</i> <i>Septoria tritici</i> <i>Blumeria</i> spp., <i>Erysiphe</i> spp. <i>Pyrenophora tritici-repentis</i>	2 – 4	Apply in the spring. Reapply up to Feekes growth stage 10.5 for season long control. The minimum re-treatment interval is 14 days.	<ul style="list-style-type: none"> <li>Do not apply more than a total of 8 fl oz of PropiMax EC (0.22 lb active ingredient) per acre per season.</li> <li>Do not apply more than a total of 4 fl oz of PropiMax EC (0.11 lb active ingredient) per acre per season if forage or hay will be harvested.</li> <li>Do not apply after Feekes 10.5 in wheat.</li> <li><b>Preharvest Interval:</b> Do not apply within 30 days of harvest for forage or hay.</li> </ul>
<b>control of:</b> glume blotch helminthosporium leaf blight leaf blight net blotch powdery mildew rust spot blotch tan spot	<i>Stagonospora nordorum</i> <i>Drechslera tritici-repentis</i> <i>Septoria tritici</i> <i>Pyrenophora teres</i> <i>Blumeria</i> spp., <i>Erysiphe</i> spp. <i>Puccinia</i> spp. <i>Bipolaris sorokiniana</i> <i>Pyrenophora tritici-repentis</i>	4	Protecting the flag leaf is important for maximizing yield. Highest yields are normally obtained when the product is applied when the flag leaf is 50% to fully emerged. The minimum re-treatment interval is 14 days. Using an oil base adjuvant may improve the spray coverage and canopy penetration. In wheat, the product can be applied through full head emergence (Feekes growth stage 10.5). Do not apply after this stage to avoid possible illegal residues.	
foot rot	<i>Pseudocercospora</i> spp.		Apply with half rates of fungicides such as thiophanate-methyl. Apply at tillering but before elongation has occurred.	
fusarium head blight suppression			Apply at approximately 50% flowering. Adding a penetrating type of adjuvant may increase fusarium head blight suppression.	

---

## Terms and Conditions of Use

---

If terms of the following Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies are not acceptable, return unopened package at once to the seller for a full refund of purchase price paid. Otherwise, use by the buyer or any other user constitutes acceptance of the terms under Warranty Disclaimer, Inherent Risks of Use and Limitations of Remedies.

---

## Warranty Disclaimer

---

Dow AgroSciences warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions, subject to the inherent risks set forth below. Dow AgroSciences MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.

---

## Inherent Risks of Use

---

It is impossible to eliminate all risks associated with use of this product. Plant injury, lack of performance, or other unintended consequences may result because of such factors as use of the product contrary to label instructions (including conditions noted on the label, such as unfavorable temperature, soil conditions, etc.), abnormal conditions (such as excessive rainfall, drought, tornadoes, hurricanes), presence of other materials,

the manner of application, or other factors, all of which are beyond the control of Dow AgroSciences or the seller. To the extent permitted by law, all such risks shall be assumed by buyer.

---

## Limitation of Remedies

---

To the extent permitted by law:

- a) The exclusive remedy for losses or damages resulting from this product (including claims based on contract, negligence, strict liability, or other legal theories), shall be limited to, at Dow AgroSciences' election, one of the following:
  - i. Refund of purchase price paid by buyer or user for product bought, or
  - ii. Replacement of amount of product used.
- b) Dow AgroSciences shall not be liable for losses or damages resulting from handling or use of this product unless Dow AgroSciences is promptly notified of such loss or damage in writing. In no case shall Dow AgroSciences be liable for consequential or incidental damages or losses.
- c) The terms of the Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies cannot be varied by any written or verbal statements or agreements. No employee or sales agent of Dow AgroSciences or the seller is authorized to vary or exceed the terms of the Warranty Disclaimer or this Limitation of Remedies in any manner.

©Trademark of Dow AgroSciences LLC  
EPA accepted 01/05/12



# PropiMax<sup>®</sup> EC

## Fungicide

For control of certain diseases in almond, berries, carrot, celery, cereal grains, cilantro (coriander) leaves, corn, cranberry, garden beet, leaf petioles, nonbearing citrus fruits, onions, parsley leaves (fresh and dried), peanut, pecan, peppermint, pineapple, rice, sorghum, spearmint, stone fruits, sugarcane, tree nuts, turfgrass grown for seed, turfgrass and ornamentals, wild rice, and nonbearing fruit and nut trees in nurseries and landscape settings.

Group	3	FUNGICIDE
-------	---	-----------

Active Ingredient:

propiconazole: 1-[[2-(2,4-dichlorophenyl)-4-propyl-1,3-dioxolan-2-yl]methyl]-1*H*-1,2,4-triazole ..... 41.8%

Other Ingredients ..... 58.2%  
 Total ..... 100.0%

Contains petroleum distillates.  
 Contains 3.6 lb active ingredient per gallon.

## Keep Out of Reach of Children

# WARNING AVISO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

**Agricultural Use Requirements**  
 Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. Refer to the label booklet under "Agricultural Use Requirements" in the Directions for Use section for information about this standard.

For additional Precautionary Statements, First Aid, Storage and Disposal and other use information see inside this label.

**Notice:** Read the entire label before using. Use only according to label directions. **Before using this product, read Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies at end of label booklet. If terms are unacceptable, return at once unopened.**

In case of emergency endangering health or the environment involving this product, call 1-800-992-5994.

Agricultural Chemical: Do not ship or store with food, feeds, drugs or clothing.

EPA Reg. No. 62719-346 900-018842 / 00303908

®Trademark of Dow AgroSciences LLC

**Produced for**  
**Dow AgroSciences LLC**  
 9330 Zionsville Road  
 Indianapolis, IN 46268

Net Contents **1 gal**



(01) 0 06 62974 34116 1

PEEL FILM HERE