Dow AgroSciences LLC
9330 Zionsville Road
Indianapolis, IN 46268-1054

Building 308/2E
800-892-6740
Contact Person: Tracey Wiley

Department of Agriculture
STATE OF HAWAII
LICENSED
PERIOD 2014-2016 LIC. NO.

Department of Agriculture
STATE OF HAWAII
LICENSED
PERIOD 2011-2013 LIC. NO.

9786.205

Dow AgroSciences

PRODUCT DATA SHEET

Kerb* WSP

62719-397

EPA Accepted: 7/18/99, Notif 8/30/99 and Notif 2/9/00

Comments:

Label Code: 192324A

Department of Agriculture
STATE OF HAWAII
LICENSED
PERIOD 2008-2010 LIC. NO.

LICENSED
HAWAII
DEPT. OF AGRICULTURE
Period 2002-2004 No. 9786.205

Kerb WSP is a new registration with the Dow AgroSciences logo as a result of our company acquiring Rohm and Haas Agricultural Products. This label is identical in compound to the last Rohm and Haas label (EPA Reg. No. 707-159) re-registered for the year 2002 in your state. Dow AgroSciences will handle and start the discontinuation process of 707-159 when the next renewal comes due in your state.

Note: The pdf pre-printed page numbers at the top, sides and immediate bottom of the label pages might not appear consistent due to eliminating empty printoff pages set aside for "notes", extra pdf overpage run-offs, etc. When applicable, please use the page numbers at the very bottom

* Trademark of Dow AgroSciences LLC

JUL 24 2002
Pesticides

RESTRICTED-USE PESTICIDE

Restricted Use Pesticide

Because pronamide has produced tumors in laboratory animals, this product is for retail sale to and use only by Certified Applicators or persons under their direct supervision, and only for those uses covered by the Certified Applicator's certification.

KERB® WSP

TURF & ORNAMENTAL HERBICIDE

ACTIVE INGREDIENT

Pronamide

3,5-dichloro-N-(1,1-dimethyl-2-propynyl) benzamide 51.0%

Inert Ingredients 49.0%

Total 100.0%

Notice: Before using this product, read the entire Precautionary Statements, Conditions of Sale and Warranty, Directions for Use, Use Restrictions and Storage and Disposal Instructions. If the Conditions of Sale and Warranty are not acceptable, return the product unopened within thirty days of purchase to the place of purchase.

KEEP OUT OF REACH OF CHILDREN

CAUTION

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Call a physician or Poison Control Center. Drink one or two glasses of water and induce vomiting by touching back of throat with finger. If person is unconscious, do not give anything by mouth and do not induce vomiting.

IF INHALED: Remove victim to fresh air. If not breathing, give artificial respiration, preferably mouth-to-mouth. Get medical attention.

IF ON SKIN: Wash with plenty of soap and water. Get medical attention if irritation persists.

IF IN EYES: Flush eyes with plenty of water. Call a physician if irritation persists.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

Causes moderate eye irritation. Avoid contact with eyes or clothing. Wash thoroughly with soap and water after handling.

ENVIRONMENTAL HAZARDS

Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. Refer to the label booklet under "Agricultural Use Requirements" in the Directions for Use section for information about this standard.

Refer to label booklet for Directions for Use including Storage and Disposal.

EPA Reg. No. 62719-397

EPA Est. 65387-AR-01
192324A

Dow AgroSciences
® Trademark of Dow AgroSciences LLC
Dow AgroSciences LLC
Indianapolis, IN 46268 U.S.A.
Emergency: 1-800-992-5994

Net Weight 3 lb
(3 x 1 lb. Water-Soluble Pouch)

00192324 8384-B7

RESTRICTED USE PESTICIDE

Because pronamide has produced tumors in laboratory animals, this product is for retail sale to and use only by Certified Applicators or persons under their direct supervision, and only for those uses covered by the Certified Applicator's certification.

KERD® WSP

TURF & ORNAMENTAL HERBICIDE

USE DIRECTIONS FOR:

- **Grasses (Ornamental)**
- **Woody Ornamentals, Nursery Stock of Ornamentals, Christmas Trees**

NOTICE: Before using this product, read the entire Precautionary Statements, Conditions of Sale and Warranty, Directions for Use, Use Restrictions and Storage and Disposal Instructions. If the Conditions of Sale and Warranty are not acceptable, return the product unopened within thirty days of purchase to the place of purchase.

KEEP OUT OF REACH OF CHILDREN

CAUTION

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Call a physician or Poison Control Center. Drink one or two glasses of water and induce vomiting by touching back of throat with finger. If person is unconscious, do not give anything by mouth and do not induce vomiting.
IF INHALED: Remove victim to fresh air. If not breathing, give artificial respiration, preferably mouth-to-mouth. Get medical attention.
IF ON SKIN: Wash with plenty of soap and water. Get medical attention if irritation persists.
IF IN EYES: Flush eyes with plenty of water. Call a physician if irritation persists.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

Causes moderate eye irritation. Avoid contact with eyes or clothing. Wash thoroughly with soap and water after handling.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Applicators and other handlers must wear:

- Coveralls over short-sleeved shirt and short pants • Waterproof gloves
- Chemical-resistant footwear plus socks
- Chemical-resistant headgear for overhead exposure
- Chemical-resistant apron when cleaning equipment, mixing, or loading

Clothing that has been drenched or heavily contaminated should be disposed of in accordance with State or local regulations.

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standards (WPS) for agricultural pesticides [40 CFR 170.240(d) (4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ACTIVE INGREDIENT

Pronamide

3,5-dichloro-N-(1,1-dimethyl-2-propynyl)-benzamide 51.0%
INERT INGREDIENTS 49.0%
TOTAL 100.0%

EPA Reg. No. 62719-397

Dow AgroSciences

® Trademark of Dow AgroSciences LLC
Dow AgroSciences LLC • Indianapolis, IN 46268 U.S.A.
Emergency: 1-800-992-5994

ENVIRONMENTAL HAZARDS

Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application.

For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of **24 hours.**

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil or water, is:

- Coveralls over short-sleeved shirt and short pants
- Waterproof gloves • Chemical-resistant footwear plus socks
- Chemical-resistant headgear for overhead exposure

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are not within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries or greenhouses.

- Keep unprotected persons out of treated area until sprays have dried.

CONDITIONS OF SALE AND WARRANTY

Dow AgroSciences warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label only when used in accordance with label directions under normal conditions of use. **Dow AgroSciences MAKES NO OTHER EXPRESS OR IMPLIED WARRANTIES EITHER OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR USE.** Handling, storage and use of the product by Buyer or User are beyond the control of Dow AgroSciences and Seller. Risks such as crop injury, ineffectiveness or other unintended consequences resulting from, but not limited to, weather or soil conditions, presence of other materials, disease, pests, drift to other crops or property or failure to follow label directions will be assumed by the Buyer or User. **IN NO CASE WILL Dow AgroSciences OR SELLER BE HELD LIABLE FOR CONSEQUENTIAL, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE HANDLING, STORAGE OR USE OF THIS PRODUCT.**

USE RESTRICTIONS

Hand-spray applications of pronamide may be made only to ornamentals and nursery stock of ornamentals.

CHEMIGATION

Do not apply this product through any type of irrigation system.

GENERAL USE INFORMATION

KERB WSP selective herbicide is formulated as a wettable powder containing 50% active ingredient packaged in a one-pound water-soluble pouch. It is effective for the control of a wide range of grasses and certain broadleaf weeds. The product is a soil-active herbicide with uptake by sensitive weeds occurring through the roots. Before using this herbicide for a specific crop use, study the following general use information which provides important instructions for the safe and effective application of the product.

WEED SPECTRUM

KERB WSP herbicide is recommended for both preemergence and early postemergence control of winter annual and perennial grasses and chickweed, and for preemergence control only of certain other broadleaf weeds and certain other grasses listed.

WEEDS CONTROLLED BOTH PREEMERGENCE AND EARLY POSTEMERGENCE

BARLEY, FOXTAIL	*Hordeum jubatum*
BARLEY, VOLUNTEER	*Hordeum vulgare*
BLUEGRASS, ANNUAL	*Poa annua*
BLUEGRASS, KENTUCKY	*Poa pratensis*
BROME, DOWNY (CHEATGRASS)	*Bromus tectorum*
CHICKWEED, COMMON	*Stellaria media*
CHICKWEED, MOUSE-EAR	*Cerastium vulgatum*
OAT, VOLUNTEER	*Avena sativa*
ORCHARDGRASS	*Dactylis glomerata*
QUACKGRASS	*Agropyron repens*
RYE, VOLUNTEER	*Secale cereale*
RYEGRASS, ITALIAN	*Lolium multiflorum*
RYEGRASS, PERENNIAL	*Lolium perenne*
WHEAT, VOLUNTEER	*Triticum aestivum*

WEEDS CONTROLLED ONLY PREEMERGENCE

BARNYARDGRASS	*Echinochloa crus-galli*
CANARYGRASS	*Phalaris canariensis*
CARPETWEED	*Mollugo verticillata*
CRABGRASS, LARGE	*Digitaria sanguinalis*
DODDER, FIELD	*Cuscuta campestris*
FOXTAIL, YELLOW	*Setaria lutescens*
GOOSEFOOT, NETTLELEAF	*Chenopodium murale*
GOOSEGRASS	*Eleusine indica*
HENBIT	*Lamium amplexicaule*
KNOTWEED, PROSTRATE	*Polygonum aviculare*
LAMBSQUARTERS, COMMON	*Chenopodium album*
LOVEGRASS	*Eragrostis diffusa*
MALLOW, LITTLE (CHEESEWEED)	*Malva parviflora*
MORNINGGLORY, ANNUAL	*Ipomoea purpurea*
MUSTARD, WILD	*Brassica kaber*
NETTLE, BURNING	*Urtica urens*
NIGHTSHADE, BLACK	*Solanum nigrum*
NIGHTSHADE, HAIRY	*Solanum sarachoides*
PANICUM, FALL	*Panicum dichotomiflorum*
PURSLANE, COMMON	*Portulaca oleracea*
RADISH, WILD	*Raphanus sativus*
ROCKET, LONDON	*Sisymbrium irio*
SHEPHERDSPURSE	*Capsella bursa-pastoris*
SMARTWEED, PALE	*Polygonum lapathifolium*
SORREL, RED (from seed)	*Rumex acetosella*
TOMATO, VOLUNTEER	*Solanum esculentum*

NOTE: The weed species controlled by KERB WSP herbicide are dependent on the rate used, specific crop culture involved, and the associated conditions of temperature, soil type and moisture availability. Refer to specific crop use directions for weed species controlled.

DOSAGE

The rate of KERB WSP herbicide required will vary depending on the crop culture involved and weed species to be controlled. See specific crop use directions for all dosage recommendations. All dosage recommendations listed in this label are in terms of pounds KERB WSP herbicide per broadcast acre. For banded application, the amount of KERB WSP herbicide used per acre should be reduced according to the following formula:

$$\frac{\text{Band Width (in inches)}}{\text{Row Width (in inches)}} \times \text{Rate per Broadcast Acre} = \text{Amount Needed per Acre for Banded Application}$$

TIMING AND APPLICATION

Unless specific directions are given under the crop to be treated, KERB WSP herbicide should be applied in the fall or early winter, when temperatures do not exceed 55°F, BUT PRIOR TO SOIL FREEZE-UP. Best weed control results when KERB WSP herbicide is applied preemergence to the weeds and when application is followed by rainfall or irrigation to move the product into the root zone of the germinating weeds.

KERB WSP herbicide should be thoroughly mixed in clean water at the recommended concentration and applied uniformly as a spray. For ground application, use a conventional low pressure herbicide sprayer equipped with flat fan nozzles spaced and calibrated to uniformly deliver 20 to 50 gallons of spray per acre. For aerial applications apply in a coarse droplet spray at 5 to 10 gallons per acre. Accurately calibrate spray equipment prior to each use.

COMPATIBILITY

KERB WSP herbicide is compatible with most commonly used agricultural pesticides, crop oil concentrate and adjuvants. When preparing tank mixes, user should consult spray compatibility charts or State Cooperative Extension Service Specialists prior to actual use.

NOTE: KERB WSP herbicide is compatible with boron and crop oil concentrate; however, the water-soluble pouches must be completely dissolved before adding spray oils or products containing boron to spray mixtures.

EFFECT OF SOIL TYPE, MOISTURE AND TEMPERATURE

KERB WSP herbicide is most active in coarse to medium-textured soils of low organic matter and is relatively inactive in peat or muck soils or mineral soils high in organic matter content at rates recommended in this label. Herbicidal activity is best in soils containing less than 4 percent organic matter. Use in soils with higher organic matter may result in inconsistent or incomplete weed control.

The herbicidal activity of KERB WSP is mainly through root absorption in sensitive weed species. Rain, melting snow or irrigation is ESSENTIAL following treatment to move KERB WSP herbicide into the root zone of germinating weeds.

Under field conditions, KERB WSP herbicide will remain relatively stable with little loss of herbicidal activity when soil temperatures are less than 55°F. As soil temperatures increase, degradation of the active ingredient takes place. KERB WSP herbicide may degrade rather quickly if left exposed on the soil surface in warm weather. If KERB WSP herbicide is applied when air temperatures exceed 85°F, it should be soil incorporated to a shallow depth (top two to three inches) or watered into the soil as soon as possible.

CULTURAL CONSIDERATIONS

For best results apply KERB WSP herbicide to a trash-free soil surface. Clean cultivation before application is preferable, but not necessary. To obtain optimum weed control in areas not clean cultivated, the area to be treated should be free of surface liter (dead or decaying crop and weed debris, mowing clippings, etc.). Trash-free areas create ideal conditions for rapid movement of KERB WSP into the weed root zone following rain or irrigation.

ROTATION CROP PLANTING INFORMATION

Follow the directions given below when rotation crops will be planted to areas previously treated with KERB WSP herbicide:

NOTE: "Crops not listed in Crop Groups A, B, C, D (below) must not be planted back sooner than 365 days after the last KERB WSP herbicide application regardless of the rate of KERB WSP used."

Further, rotational crops are restricted to artichokes, apples, apricots, cherries, nectarines, peaches, pears, plums, prunes or grapes in fields previously planted with these crops and treated with more than 4.0 pounds of KERB WSP (2.0 lb. ai), per planted acre[1].

In fields planted with blackberries, boysenberries or raspberries and treated with more than 4.0 pounds of KERB WSP per planted acre, rotational crops are restricted to these crops and artichokes, apples, apricots, cherries, nectarines, peaches, pears, plums, prunes or grapes.

Whether KERB WSP herbicide is bed-topped, banded or broadcast, the beds should be knocked down and the field cross-disced before rotation crops other than artichokes, lettuce, endive, radicchio or escarole are planted.

Where the KERB WSP herbicide treatment is to be followed by a rotation crop within 180 days of application, bed-topped or banded applications are suggested.

Where rotation crops are to follow within 365 days of the KERB WSP herbicide treatment, the following rotational plan applies:

Amount of KERB WSP Applied/Planted Acre[1]	Waiting Period in Days Before Planting the Crops Indicated[2] Group A	Group B	Group C	Group D
1.0 lb.	0	90	90	180
2.0 lbs.	0	90	120	180
3.0 lbs.	0	120	180	180
4.0 lbs.	0	150	210	180

[1] Planted Acre refers to Acreage planted. KERB WSP at 4.0 pounds of product per treated acre on two 5-inch treated bands (40-inch centers) is the equivalent of one pound KERB WSP herbicide per planted acre. KERB WSP at 4.0 pounds of product as a broadcast treatment (total acreage sprayed) is the equivalent of 4.0 pounds of KERB WSP herbicide per planted acre.

[2] Crop Group A – Artichokes
Crop Group B – Beans, cotton, carrots, celery
Crop Group C – Broccoli, cabbage, cauliflower, cucurbits, spinach, sugarbeets, onion, tomatoes
Crop Group D – Lettuce, endive, escarole, radicchio

GRASSES (ORNAMENTAL)

GROWN FOR TURF OR SEED

GENERAL INFORMATION

KERB WSP herbicide is a selective herbicide recommended for the preemergence and postemergence control of annual bluegrass (*Poa annua*) from warm season grasses and the removal of perennial rye grass (*Lolium perenne*) from warm season during spring transition. Warm season grasses include ornamental bermudagrass (*Cynodon dactylon*), Zoysiagrass, St. Augustinegrass and Centipedegrass or bermudagrass grown for seed.

ANNUAL BLUEGRASS (*Poa Annua*) CONTROL

KERB WSP herbicide will control annual bluegrass from pregermination and seedling stages through tillering, heading and seed formation. KERB WSP herbicide acts slowly on seedling to mature annual bluegrass. Following application of KERB WSP annual bluegrass may first become dark green and then gradually turn yellow and die over a 3 to 5 week period

For effective control of annual bluegrass in turf or seed crops, moisture is necessary to move KERB WSP herbicide in the weed root zone. Refer to the Moisture Requirements section of this label for details.

DOSAGE AND TIMING

For annual bluegrass control KERB WSP herbicide is recommended at the rate of 1.0 to 3.0 pounds of product (0.5 to 1.5 pounds active ingredient) per acre broadcast application. The dosage rate required is dependent on the growth stage of annual bluegrass at time of application. Follow the dosage rate and timing recommendations given below:

Annual Bluegrass Growth Stage	KERB WSP Lbs. Per Acre** Broadcast Application
Preemergence or early postemergence	1.0 - 2.0*
Postemergence – early tillering to heading	1.5 - 2.0
Postemergence – seed forming stages	2.0 - 3.0

*Use the higher rate when longer preemergence residual control is desired.
**One acre equals 43,560 sq. ft.

REMOVAL OF PERENNIAL RYE GRASS FROM WARM SEASON GRASSES

KERB WSP herbicide will remove postemergent perennial rye grass from warm season grasses during the spring to control the transition from cool season overseed to warm season grasses. KERB WSP herbicide works slowly to control mature perennial rye grass. After an application of KERB WSP herbicide, perennial rye grass will gradually die over a 4 to 6 week period. The length of this transition is dependent upon environmental factors such as temperature, rainfall and mowing height of the turf.

DOSAGE AND TIMING

For removal of perennial rye grass from warm season grasses, KERB WSP herbicide is recommended at a rate of 1.0 to 2.0 lbs. of product (0.5 - 1 lb. active ingredient) per broadcast acre. It is best to apply KERB WSP herbicide to warm season grasses at 50% greenup. Application of KERB WSP herbicide to dormant warm season grasses can slow greenup.

APPLICATION

Mix the recommended amount of KERB WSP herbicide in clean water and apply uniformly with a low pressure ground sprayer in 20 to 50 gallons of water per acre or 0.5 to 1 gallon of water per 1000 sq. ft. The sprayer should be equipped with flat fan nozzles, spaced to provide uniform distribution without skips or excessive overlapping of spray patterns.

NOTE: Avoid spraying on fairways or approaches that may drain onto bentgrass greens or to areas overseeded with sensitive cool season grasses.

Do not make an application of a wetting agent for the purpose of frost protection or soil penetration to greens or tees 14 days prior to or after a KERB WSP application as injury may result.

MOISTURE REQUIREMENTS

KERB WSP herbicide acts mainly through root absorption in sensitive weed species. If no rainfall occurs within a day or so of the application, a light overhead irrigation should be made to move the chemical into the weed root zone. Avoid heavy irrigations of more than 1 inch to reduce the possibility of excess washing or leaching of the chemical from the area of application.

KERB WSP DEACTIVATION FOR OVERSEEDING

Where it is desirable to reseed sooner than 90 days following the application of KERB WSP herbicide, an application of an activated charcoal, such as Gro-Safe®, is needed. Apply the activated charcoal at the rate of 10 pounds per 1000 square feet. Allow at least 14 days between the KERB WSP application and the charcoal application for control of emerged annual bluegrass. Reseed no sooner than seven days following charcoal application.

GRASSES (ORNAMENTAL)
TURF OR SEED CROP

– SPECIFIC USE RESTRICTIONS

- This product may be used on seeded, sodded, or sprigged ornamental turf or turf grown for seed that is well established. Use of this product on turf that has been weakened by weather-, pest-, disease- chemical-, or mechanical-related stress may increase the chances of turf injury.
- This product should only be applied to turf lawns that are composed of the following turfgrass species:
  Bermudagrass (*Cynodon dactylon*)
  Centipedegrass (*Eremochloa ophiuroides*)
  St. Augustinegrass (*Stenotaphrum secundatum*)
  Zoysiagrass (*Zoysia japonica*)
- Avoid spraying on hill sides, fairways, or approaches that may drain onto bentgrass greens or to areas overseeded with sensitive cool season grasses. Do not make an application of a wetting agent for the purpose of frost protection or soil penetration to greens or tees 14 days prior to or after a KERB application as injury may result.
- Do not apply KERB WSP herbicide to areas that are to be overseeded with susceptible cool season grasses within 90 days of treatment unless deactivation is planned.
- Do not apply KERB WSP to dicondra, perennial bluegrass, annual and perennial ryegrasses, fescues and bentgrasses.
- Do not graze treated areas and do not feed clippings to livestock.
- Do not apply more than one application of KERB WSP herbicide per season and do not apply KERB WSP herbicide at rates higher than 3.0 pounds product (1.5 pounds active ingredient) per acre.

WOODY ORNAMENTALS • NURSERY STOCK OF ORNAMENTALS • CHRISTMAS TREES

GENERAL INFORMATION

KERB WSP herbicide is a selective herbicide recommended for fall applications to established woody ornamentals, nursery stock of ornamentals and Christmas trees for the control of winter annual and perennial grasses and certain broadleaf weeds.

CROP TOLERANCE

At recommended rates of KERB WSP herbicide the following trees and shrubs are tolerant to topical applications made in the fall:

ARBORVITAE	FIRETHORN	MOUNTAIN ASH
ASH	FLOWERING CHERRY	MOUNTAIN LAUREL
AZALEA	FLOWERING CRABAPPLE	OAK
BARBERRY	FLOWERING QUINCE	OHIO BUCKEYE
BASSWOOD	FORSYTHIA	PINE
BEECH	GINKO	POPLAR
BIRCH	HAWTHORN	PRIVET
BOXWOOD	HEMLOCK	RHODODENDRON
BRADFORD PEAR	HOLLY	SPIREA
CEDAR	HONEY LOCUST	SPRUCE
COTONEASTER	JUNIPER	SWEETGUM
DOGWOOD	LILAC	SYCAMORE
DOUGLAS FIR	LINDEN	TULIPTREE
EASTERN REDBUD	LONDON PLANE	VIBURNUM
ELM	MAGNOLIA	WALNUT
EUONYMUS	MAPLE	WILLOW
FIR	MOCK ORANGE	YEW

KERB WSP herbicide should only be used on established trees and woody ornamentals. KERB WSP herbicide is not recommended for use on seedling trees or shrubs less than one year old or to fall transplanted stock transplanted less than one year or to spring transplanted stock transplanted less than six months.

WEED CONTROL

KERB WSP herbicide is effective in fall applications at the rate of 2.0 to 4.0 pounds of product (1.0 to 2.0 pounds active ingredient) per broadcast acre for the preemergence and postemergence control of susceptible winter annual and perennial grasses and chickweed and for preemergence control only of other broadleaf weeds listed on this label. Refer to chart in Dosage and Timing section below for specific weeds controlled.

DOSAGE AND TIMING

KERB WSP herbicide is recommended in a single, fall application, either directed or topically applied, to woody ornamentals, nursery stock of ornamentals or Christmas trees at the rate of 2.0 to 4.0 pounds of product (1.0 to 2.0 pounds active ingredient) per broadcast acre. Application of KERB WSP herbicide should be made in the fall prior to leaf drop and soil freeze-up. For control of winter annual or perennial grasses or chickweed, applications can be made either preemergence or postemergence to the weeds. For control of other labeled broadleaf weeds, preemergence applications must be used to achieve control.

The dosage rate required will depend on the weed species present in the area to be treated. Follow the weed control recommendations given in the chart below:

Weeds Controlled	Pounds KERB WSP Per Acre Broadcast Application
BARLEY, FOXTAIL BLUEGRASS, ANNUAL BROME, DOWNY (CHEATGRASS) CHICKWEED GRAIN, VOLUNTEER RYEGRASS, ITALIAN SORREL, RED (from seed)	2.0
MUSTARD, WILD ROCKET, LONDON SHEPHERDSPURSE	3.0
BLUEGRASS, KENTUCKY ORCHARDGRASS QUACKGRASS RYEGRASS, PERENNIAL	4.0

APPLICATION

Mix the recommended amount of KERB WSP herbicide in clean water and apply uniformly in 20 to 50 gallons per acre. Use a low pressure ground sprayer equipped with flat fan nozzles spaced to provide uniform distribution.

Dosage recommendations listed on this label are for surface broadcast application. For banded treatments down the row, the amount of KERB WSP used per acre should be reduced according to the following formula:

$$\frac{\text{Band Width (in inches)}}{\text{Row Width (in inches)}} \times \text{Rate per Broadcast Acre} = \text{Amount Needed per Acre for Banded Application}$$

KERB WSP herbicide should not be soil incorporated.

NOTE: Most ornamental turf grass species and ground covers are sensitive to KERB WSP. Care should be exercised to avoid contact of KERB WSP herbicide with these plants from either direct application, spray drift or from applications to areas that may drain onto established ornamental turf and ground cover.

SOIL AND MOISTURE REQUIREMENTS

KERB WSP herbicide is most active in coarse to medium-textured soils of low organic matter and is relatively inactive in peat or muck soils or mineral soils high in organic matter content at rates recommended in this label. Herbicidal activity is best in soils containing less than 4 percent organic matter. Use in soils of higher organic matter content may result in inconsistent or incomplete weed control.

KERB WSP herbicide acts mainly through root absorption in sensitive weed species. Dependable rainfall or overhead irrigation is essential following application for effective weed control.

WOODY ORNAMENTALS/NURSERY STOCK OF ORNAMENTALS/CHRISTMAS TREES – SPECIFIC USE RESTRICTIONS

- Apply KERB WSP herbicide in the fall prior to soil freeze-up.
- Do not soil incorporate KERB WSP herbicide.
- Do not harvest plants for food or feed for at least one year after treatment.
- Do not make more than one application of KERB WSP herbicide per year and do not apply KERB WSP herbicide at rates higher than 4.0 pounds of product (2.0 lbs. ai) per acre.

STORAGE AND DISPOSAL

PROHIBITIONS: Do not contaminate water, food or feed by storage or disposal. Open dumping is prohibited.

STORAGE: Store in a cool, dry place but not below 32°F (0°C). Do not remove package from container except for immediate use.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Completely empty bag into application equipment. Then dispose of in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

STEPS TO BE TAKEN IN CASE MATERIAL IS RELEASED OR SPILLED: Sweep up and place in containers for recovery or disposal. Keep dusting to a minimum. Wear MSHA/NIOSH respirator for dusts encountered. Flush area with water. Keep spill out of all sewers and open bodies of water. Protective Clothing – See Precautionary Statements.

ATTENTION: This product contains propyzamide (pronamide) a chemical known to the State of California to cause cancer.

GLEAN is a registered trademark of the E. I. duPont de Nemours & Company, Inc.
Gro-Safe is a registered trademark of American Norit Company, Inc.