

Group	3	FUNGICIDE
-------	---	-----------



Enable[®]

2F



STATE OF HAWAII
Department of Agriculture

ACCEPTED

LICENSE NO.
9786.208

PEEL FILM HERE

Fungicide

Group	3	FUNGICIDE
-------	---	-----------

Active Ingredient

fenbuconazole: a-[2-(4-chlorophenyl)ethyl]-a-phenyl-1H-1,2,4-triazole-1-propanenitrile...	23.5%
Other Ingredients.....	76.5%
Total.....	100.0%

Contains 2 lb of active ingredient per gallon

Keep Out of Reach of Children CAUTION

First Aid

If inhaled: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. Call a poison control center or doctor for further treatment advice.

Hot Line Number: Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-992-5994 day or night, for emergency treatment information.

Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. Refer to the label booklet under "Agricultural Use Requirements" in the Directions for Use section for information about this standard.

For additional Precautionary Statements, First Aid, Storage and Disposal and other use information see inside this label.

Notice: Read the entire label. Use only according to label directions. **Before using this product, read Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies at end of label booklet. If terms are unacceptable, return at once unopened.**

In case of emergency endangering health or the environment involving this product, call 1-800-992-5994.

Agricultural Chemical: Do not ship or store with food, feeds, drugs or clothing.

EPA Reg. No. 62719-416 900-021470 / 00284587

®Trademark of Dow AgroSciences LLC

Produced for
Dow AgroSciences LLC
9330 Zionsville Road
Indianapolis, IN 46268

Net Contents **1 gal**

Group	3	FUNGICIDE
-------	---	-----------



Enable[®]

2F

Fungicide

Group	3	FUNGICIDE
-------	---	-----------

Active Ingredient

fenbuconazole: a-[2-(4-chlorophenyl)ethyl]-a-phenyl-1H-1,2,4-triazole-1-propanenitrile.....	23.5%
Other Ingredients.....	76.5%
Total	100.0%

Contains 2 lb of active ingredient per gallon

**Keep Out of Reach
of Children**
CAUTION

First Aid

If inhaled: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. Call a poison control center or doctor for further treatment advice.

Hot Line Number: Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-992-5994 day or night, for emergency treatment information.

Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. Refer to the label booklet under "Agricultural Use Requirements" in the Directions for Use section for information about this standard.

For additional Precautionary Statements, First Aid, Storage and Disposal and other use information see inside this label.

Notice: Read the entire label. Use only according to label directions. **Before using this product, read Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies at end of label booklet. If terms are unacceptable, return at once unopened.**

In case of emergency endangering health or the environment involving this product, call 1-800-992-5994.

Agricultural Chemical: Do not ship or store with food, feeds, drugs or clothing.

EPA Reg. No. 62719-416

EPA Est. 5905-GA-01
900-021470 / 00284587

®Trademark of Dow AgroSciences LLC

Produced for
Dow AgroSciences LLC
9330 Zionsville Road
Indianapolis, IN 46268

Net Contents **1 gal**

Table of Contents	Page
Precautionary Statements.....	1
Hazards to Humans and Domestic Animals.....	1
Personal Protective Equipment (PPE).....	1
Engineering Controls.....	1
User Safety Recommendations.....	1
Environmental Hazards.....	1
Directions for Use.....	1
Agricultural Use Requirements.....	2
Storage and Disposal.....	2
General Product Information.....	2
Resistance Management.....	2
Use Rate Determination.....	2
Mixing Directions.....	3
Application Directions.....	3
Orchard Use.....	3
Aerial Spray Drift Management.....	3
Compatibility.....	3
Plantback Intervals.....	3
Uses.....	4
Banana.....	4
Citrus Fruits(Crop Group 10).....	4
Pecan.....	5
Peppers.....	5
Sugarbeets.....	6
Terms and Conditions of Use.....	6
Warranty Disclaimer.....	6
Inherent Risks of Use.....	7
Limitation of Remedies.....	7

Precautionary Statements

Hazards to Humans and Domestic Animals

CAUTION

Harmful if Inhaled

Avoid breathing spray mist. Remove and wash contaminated clothing before reuse.

Personal Protective Equipment (PPE)

Some materials that are chemical-resistant to this product are listed below. If you want more options follow the instructions for Category A on an EPA chemical resistant category selection sheet.

WPS Uses: Applicators and other handlers who handle this pesticide for any use covered by the Worker Protection Standard (40 CFR Part 170):

- Long-sleeved shirt and long pants
- Shoes plus socks

- Chemical-resistant gloves made of any waterproof material, such as polyvinyl chloride, nitrile rubber, or butyl rubber.
- Chemical-resistant headgear for overhead exposure

User Safety Requirements

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

Engineering Controls

When handlers use closed systems, enclosed cabs or aircraft in a manner that meet the requirements listed in the Worker Protection Standards (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

User Safety Recommendations

Users should:

- Wash hands before eating, drinking, changing gum, using tobacco or using the toilet.
- Remove clothing/PPE immediately if pesticides get inside. Then wash and put on clean clothing.
- Remove PPE immediately after handling product. Wash outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

Environmental Hazards

This product is toxic to fish. Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high-water mark. Drift or runoff from treated areas may be hazardous to aquatic organisms in adjacent aquatic sites. Do not contaminate water when disposing of equipment washwaters or rinsate.

Directions for Use

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Read all Directions for Use carefully before applying.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your state or tribe, consult the agency responsible for pesticide regulation.

Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil or water, is:

- Coveralls
- Shoes plus socks
- Chemical-resistant gloves made of any waterproof material, such as polyvinyl chloride, nitrile rubber, or butyl rubber.

Storage and Disposal

Do not contaminate water, food or feed by storage or disposal.

Pesticide Storage: Store in a cool dry well-ventilated area, but not below 32°F (0°C).

Pesticide Disposal: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

Container Handling: Nonrefillable container. Do not reuse or refill this container.

Triple rinse or pressure rinse container (or equivalent) promptly after emptying. **Triple rinse** as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. **Pressure rinse** as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application

Storage and Disposal (Cont.)

equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 psi for at least 30 seconds. Drain for 10 seconds after the flow begins to drip. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures allowed by state and local authorities.

Product Information

Enable® 2F fungicide is a systemic fungicide with preventative and curative properties used for the control of specific diseases listed on this label. For optimal disease control with Enable 2F, apply sprays on a preventative spray program.

Enable 2F is considered not hazardous to bees, including those which are actively foraging in the area during application, when used as directed.

A non-polymer containing spray adjuvant approved for use with registered pesticide products may be added to spray solutions according to manufacturer's use instructions to achieve optimum disease control.

Do not apply Enable 2F in greenhouses.

Resistance Management

Enable 2F belongs to the sterol demethylation inhibitor (DMI) class of fungicides or target site of action Group 3 fungicide. Since certain fungi can develop resistance to this class of products, use Enable 2F as part of a resistance management strategy that includes alternation and/or mixing with fungicides of different target site of action. Consult your local or state agricultural authorities for resistance management strategies that are appropriate for your disease management program.

Use Rate Determination

Carefully read, understand and follow label use rates and restrictions. Under low disease conditions, use minimum application rates, while maximum application rates and shortened spray schedules are for severe or threatening disease conditions. For proper application, determine the number of acres to be treated, the label use rate and the gallonage to be applied per acre. Prepare only the amount of spray solution required to treat the measured acreage. Carefully calibrate spray equipment prior to use.

Mixing Directions

Thoroughly shake container before using contents. Triple-rinse container used to measure Enable 2F and add rinsate to spray tank. Add a registered spray adjuvant after Enable 2F is fully dispersed in the spray solution. **Note:** Reduced product efficacy may occur if water containing suspended soil particles, such as water from ponds, streams or unlined ditches, is used.

Application Directions

Do not make ground or aerial applications within 75 feet of streams, rivers, ponds, lakes or reservoirs.

Ground Application

Thorough coverage sprays generally result in optimum disease control. To achieve good coverage use proper spray pressure, gallonage per acre, nozzles, nozzle spacing and tractor speed. Consult spray nozzle and accessory catalogues for specific information on proper equipment calibration. For tree fruits and nuts, apply the same amount of Enable 2F per acre in either dilute or concentrate sprays.

Aerial Application

Apply in a minimum of 5 gallons of water per acre on annual crops and 10 gallons of water per acre on perennial tree fruits and nuts. Do not apply Enable 2F under conditions when uniform coverage cannot be obtained or when excessive spray drift may occur. Do not apply directly to humans or animals. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Flagmen and loaders should wash thoroughly before eating, drinking, or smoking and at the end of each day's spray operation.

Orchard Use

Apply Enable 2F to orchards using a dilute or concentrate spray.

Dilute Spray: Dilute or thorough coverage applications are based upon the amount of spray solution necessary to thoroughly wet the trees until spray run-off. To determine the number of gallons of dilute spray required per acre, contact your state agricultural experiment station, certified pest control advisor, or extension specialist for assistance.

Concentrate Spray: Concentrate spray applications assume all of the tree is uniformly covered with spray solution but not to the point of run-off. A lower spray volume is used to deliver the same application rate per acre as used for a dilute spray. Use a minimum of 50 gallons of spray solution per acre for concentrate sprays with Enable 2F.

Aerial Spray Drift Management

Avoiding spray drift at the application site is the responsibility of the applicator. The interaction of many equipment and weather-related factors determine the potential for spray drift. The applicator is responsible for considering all these factors when making decisions. The following drift management requirements must be followed to avoid off-target movement from aerial applications to agricultural field crops.

- The distance of the outer most nozzles on the boom must not exceed 3/4 the length of the wingspan or rotor.
- Nozzles must always point backward parallel with the air stream and never be pointed downwards more than 45 degrees.

Where states have more stringent regulations, they must be observed.

Compatibility

Enable 2F is compatible with most commonly used agriculture fungicides, insecticides, growth regulators, micronutrients and spray adjuvants. For any registered adjuvant, follow the manufacturer's use instructions to achieve proper spray concentrations. When preparing tank mixes, consult spray compatibility charts or State Cooperative Extension Service Specialists prior to actual use.

Use of Adjuvants: Surfactants may improve spray coverage, especially for waxy leaf surfaces. For best results do not use penetrants. Use only adjuvants that are registered for agricultural use and are known to be safe for the crop. Follow directions on the manufacturer's label. When an adjuvant is to be used with this product, Dow AgroSciences recommends the use of a Chemical Producers and Distributors Association certified adjuvant. For nonionic surfactants, a spray mix concentration of 0.25% v/v or less generally is sufficient. For organosilicones, a spray mix concentration of 0.05% v/v or less generally is sufficient. Conduct a compatibility test (jar test) using the relative proportions of the tank mix ingredients before mixing ingredients in the spray tank.

Plantback Intervals

Plantback intervals (PBI) for crops without primary label uses (such as rotational crops), where the last application of the product expended is <0.188 lb active per acre, the PBI must be 35 days. Where the last application of the product expended is 0.188 to 1 lb active per acre, the PBI must be 210 days.

Uses

Banana

Disease	Enable 2F fl oz/acre	Use Directions	Restrictions
sigatoka (<i>Mycosphaerella</i> spp.)	6 (0.09 lb active)	Apply when leaves first appear and repeat every 14 to 21 days or as required. Best control of diseases is achieved on a protectant application schedule and when an emulsifiable spray oil is added to the spray solution. Apply aerially in a minimum of 10 gallons of water per acre or by ground in a minimum of 20 gallons of water per acre.	<ul style="list-style-type: none"> Do not make more than 8 applications of Enable 2F (0.75 lb active) per acre per year. Preharvest Interval: Up to day of harvest. Do not graze livestock in treated areas. Do not cut cover crops for livestock food. Do not apply this product by chemigation to banana.

Citrus Fruits (Crop Group 10)¹

¹Citrus fruits (crop group 10) including calamondin, citrus citron, chironja, citrus hybrids, grapefruit, kumquat, lemon, lime, orange (sour, sweet, mandarin), pummelo, satsuma mandarin, tangerine, tangelo, tangor

Apply aerially in a minimum of 10 gallons of water per acre or by ground in a minimum of 20 gallons of water per acre. Obtain optimum disease control by applying Enable 2F with a nonionic surfactant or spray oil at the manufacturer's specified rate and making thorough coverage applications.

Disease	Enable 2F fl oz/acre	Use Directions	Restrictions
greasy spot (<i>Mycosphaerella citri</i>)	8 (0.12 lb active)	Make the first application at the normal timing for greasy spot in your area. Make a second application for control of greasy spot of foliage and rind blotch of fruit after the first summer flush. Application for control of greasy spot may also occur at other times.	<ul style="list-style-type: none"> Do not make more than 3 applications or apply more than a total of 24 fl oz of Enable 2F (0.38 lb active) per acre per year. Preharvest Interval: Up to day of harvest. Minimum Re-Treatment Interval: 21 days. Do not allow livestock to graze in treated areas. Do not apply Enable 2F with polymer based spray adjuvants. Do not apply this product by chemigation to citrus.
scab (<i>Elsinoe fawcettii</i>)		Make the first application between feather flush and 25% expansion of spring leaf flush. Make a second application at petal fall. Follow 3 to 4 weeks later with another fungicide for control of late scab and early melanose.	
sooty mold (<i>Capnodium</i> spp.)		Applications made for control of greasy spot or scab will also control sooty mold.	

Pecan

Disease	Enable 2F fl oz/acre	Use Directions	Restrictions
downy spot <i>(Mycosphaerella caryigena)</i> gnomonium leafspot <i>(Gnomonia dispersa)</i> leaf blotch <i>(Mycosphaerella dendroides)</i> powdery mildew <i>(Microspora penicillata)</i> scab <i>(Cladosporium caryigenum)</i> vein spot <i>(Gnomonia nerviseda)</i>	8 (0.125 lb active)	Begin applications at bud break and continue at 10- to 14-day intervals through pollination. Apply at 14- to 21-day intervals through the cover sprays. Apply aerially in a minimum of 10 gallons of water per acre or by ground in a minimum of 20 gallons of water per acre.	<ul style="list-style-type: none"> Do not make more than 6 applications or apply more than a total of 48 fl oz of Enable 2F (0.75 lb active) per acre per season. Preharvest Interval: Do not apply after shuck split or within 28 days of harvest. Do not graze livestock in treated areas or feed cover crops grown in treated areas to livestock. Do not apply after shuck split. Do not apply this product by chemigation to pecan.

Peppers¹

¹Peppers including bell pepper, chili pepper, cooking pepper, pimento, and sweet pepper

Disease	Enable 2F fl oz/acre	Use Directions	Restrictions
cercospora leaf spot <i>(Cercospora capsici)</i> powdery mildew <i>(Leveillula taurica)</i>	6 - 12 (0.094 – 0.188 lb active)	Begin applications when disease is first observed. Reapply 10 to 14 days later or as necessary for the level of disease pressure present. Apply uniformly in a spray volume that assures thorough coverage of the crop. A minimum spray volume of 30 gallons per acre is necessary. Disease control may be reduced at low spray volumes or if spray coverage is inadequate. For optimal disease control, apply as a ground application. Aerial application is permitted; however, disease control may be reduced when product is applied by air. For optimum performance, add a nonionic surfactant to the solution.	<ul style="list-style-type: none"> Do not apply more than 12 fl oz of Enable 2F per acre per application or more than a total of 48 fl oz of Enable 2F (0.75 lb active) per acre per crop. Do not make more than 2 consecutive applications of Enable 2F. Alternate within 14 days with a product that is effective on the target pathogen and has a different mode of action than Enable 2F. Preharvest Interval: Do not apply within 7 days of harvest. Minimum Re-Treatment Interval: 10 days Chemigation: Do not apply this product through any type of irrigation system.

Sugarbeets

Disease	Enable 2F fl oz/acre	Use Directions	Restrictions
cercospora leaf spot (<i>Cercospora beticola</i>) powdery mildew (<i>Erysiphe</i> spp.)	8 (0.125 lb active)	Apply Enable 2F at intervals of 14 to 21 days beginning when conditions are favorable for disease development. For resistance management, alternate applications of Enable 2F with a non-triazole fungicide labeled for use on sugarbeets. For optimum disease control, an adjuvant, such as crop oil concentrate that is labeled for crop use, may be added to the spray mixture at the manufacturer's specified rate. Add an adjuvant to the spray solution last. For best spray coverage of foliage, apply aerially in a minimum of 5 to 7 gallons of water per acre or by ground in a minimum of 20 gallons of water per acre.	<ul style="list-style-type: none">Do not make more than 2 applications or apply more than a total of than 16 fl oz of Enable 2F (0.25 lb active) per acre per year.Preharvest Interval: Do not apply within 14 days of harvest.Do not apply this product by chemigation to sugarbeets.

Terms and Conditions of Use

If terms of the following Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies are not acceptable, return unopened package at once to the seller for a full refund of purchase price paid. Otherwise, use by the buyer or any other user constitutes acceptance of the terms under Warranty Disclaimer, Inherent Risks of Use and Limitations of Remedies.

Warranty Disclaimer

Dow AgroSciences warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions, subject to the inherent risks set forth below. Dow AgroSciences MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.

Inherent Risks of Use

It is impossible to eliminate all risks associated with use of this product. Plant injury, lack of performance, or other unintended consequences may result because of such factors as use of the product contrary to label instructions (including conditions noted on the label, such as unfavorable temperature, soil conditions, etc.), abnormal conditions (such as excessive rainfall, drought, tornadoes, hurricanes), presence of other materials, the manner of application, or other factors, all of which are beyond the control of Dow AgroSciences or the seller. All such risks shall be assumed by buyer.

Limitation of Remedies

To the extent permitted by law, the exclusive remedy for losses or damages resulting from this product (including claims based on contract, negligence, strict liability, or other legal theories) shall be limited to, at Dow AgroSciences' election, one of the following:

1. Refund of purchase price paid by buyer or user for product bought, or
2. Replacement of amount of product used

To the extent permitted by law, Dow AgroSciences shall not be liable for losses or damages resulting from handling or use of this product unless Dow AgroSciences is promptly notified of such loss or damage in writing. In no case shall Dow AgroSciences be liable for consequential or incidental damages or losses.

To the extent permitted by law, the terms of the Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies cannot be varied by any written or verbal statements or agreements. No employee or sales agent of Dow AgroSciences or the seller is authorized to vary or exceed the terms of the Warranty Disclaimer or this Limitation of Remedies in any manner.

®Trademark of Dow AgroSciences LLC
EPA accepted 07/11/13

51 1/4"

1 7/16"

14 15/16"

9 15/16"

15 1/16"


9 7/8"

Do Not Cut To Open
Contents In Plastic



4 7/8"

00281846



Dow AgroSciences



Fungicide 4 x 1 gal



Dow AgroSciences



Fungicide 4 x 1 gal

00281846

4G/Z27.5/S/13
USA/AA2110



Dow AgroSciences



Fungicide 4 x 1 gal

13 5/16"

Dow AgroSciences

Enable 2F

Fungicide

Group 3 FUNGICIDE

Active Ingredient		
fenbuconazole: a-[2-(4-chlorophenyl)ethyl]-a-phenyl-1H-1,2,4-triazole-1-propanenitrile.....	23.5%	
Other Ingredients.....	76.5%	
Total.....	100.0%	

Contains 2 lb of active ingredient per gallon

Keep Out of Reach of Children

CAUTION

First Aid

If inhaled: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. Call a poison control center or doctor for further treatment advice.

Hot Line Number: Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-992-5994 day or night, for emergency treatment information.

Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. Refer to the label booklet under "Agricultural Use Requirements" in the Directions for Use section for information about this standard.

Refer to inside of label booklet for additional precautionary information including Directions for Use.

Notice: Read the entire label. Use only according to label directions. Before using this product, read Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies at end of label booklet. If terms are unacceptable, return at once unopened.

In case of emergency endangering health or the environment involving this product, call 1-800-992-5994.

Agricultural Chemical: Do not ship or store with food, feeds, drugs or clothing.

EPA Reg. No. 62719-416 EPA Est. 5905-GA-01

*Trademark of Dow AgroSciences LLC

Produced for
Dow AgroSciences LLC
9330 Zionsville Road
Indianapolis, IN 46268

4 x 1 gal

900-018199 / 00284586
BOX MFG. DATE:

4 7/8"

23 1/16"
CORR.

MASTER 900-006156
4 x 1 gal, 4 x 4 L, 4 x 4.45 L Mottled White
Dimensions: 14.75" L x 9.625" W x 12.625" H