



Department of Agriculture
STATE OF HAWAII

LICENSED

PERIOD **2014-2016** LIC. NO.

9786.230

PEEL FILM HERE 



Department of Agriculture
STATE OF HAWAII



Dow AgroSciences

Fore[®] 80WP Rainshield[®]

SPECIALTY FUNGICIDE

Product in Water-Soluble Packets

Group	M3	FUNGICIDE
-------	----	-----------

Active Ingredients

Mancozeb: A coordination product of zinc ion
and manganese ethylene bisdithiocarbamate..... 80.0%

In which the ingredients are:

Manganese⁺⁺ 16.00%

Zinc⁺⁺ 2.00%

Ethylene bisdithiocarbamate
ion (C₄H₆N₂S₄) 62.00%

Other Ingredients..... 20.0%

Total 100.0%

Keep Out of Reach of Children

CAUTION

Refer to inside of label booklet for additional precautionary
information including Directions for Use.

LICENSED

PERIOD **2014-2016** LIC. NO.

9786.230

In case of emergency endangering health or the environment
involving this product, call 1-800-992-5994.

Agricultural Chemical: Do not ship or store with food, feeds,
drugs or clothing.

EPA Reg. No. 62719-388

900-022165 / 00187717

MADE IN COLOMBIA

®Trademark of The Dow Chemical Company ("Dow") or an
affiliated company of Dow

Produced for
Dow AgroSciences LLC
9330 Zionsville Road
Indianapolis, IN 46268

Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. Refer to the label booklet under "Agricultural Use Requirements" in the Directions for Use section for information about this standard

Notice: Read the entire label. Use only according to label directions. **Before using this product, read Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies at end of label booklet. If terms are unacceptable, return at once unopened.**

First Aid

If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

If on skin or clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-992-5994 day or night, for emergency treatment information.

Precautionary Statements

Hazards to Humans and Domestic Animals

CAUTION

Harmful If Absorbed Through The Skin • Causes Moderate Eye Irritation • Avoid contact with skin, eyes or clothing • Prolonged Or Frequently Repeated Skin Contact May Cause Allergic Reactions In Some Individuals

Personal Protective Equipment (PPE)

Some materials that are chemical-resistant to this product are butyl rubber and neoprene rubber. If you want more options, follow the instructions for category "A" on an EPA chemical resistance category selection chart.

Mixers, loaders, applicators and all other handlers must wear:

- Long-sleeved shirt
- Long pants
- Shoes and socks
- Chemical resistant gloves made of any waterproof material (except for pilots)

In addition, all handlers (except pilots, groundboom applicators, and airblast applicators) must wear:

- A NIOSH-approved respirator with a dust/mist filter with MSHA/NIOSH approval number prefix TC-21C or any R, P or HE filter.
- Use an R, P or HE filter when mixing Fore 80WP Rainshield with an oil-based adjuvant.

See engineering controls for additional requirements.

User Safety Requirements

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry. Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them.

Engineering Controls

Water-soluble packets when used correctly qualify as a closed mixing/loading system under the Worker Protection Standard for Agricultural Pesticides [40 CFR 170.240(d)(4)]. Mixers and loaders using water-soluble packets must:

- Wear the personal protective equipment required in the PPE section of this labeling for mixers/loaders
- Be provided, and have immediately available, and wear in an emergency, such as a broken package, spill, or equipment breakdown: a NIOSH-approved respirator with a dust/mist filter with MSHA/NIOSH approval number prefix TC-21C or any R, P or HE filter.

User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside, then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

Environmental Hazards

This pesticide is toxic to aquatic organisms. Drift and runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when cleaning equipment or disposing of equipment washwaters or rinsate.

Directions for Use

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Read all Directions for Use carefully before applying.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your state or tribe, consult the agency responsible for pesticide regulation.

Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 24 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil or water, is:

- Coveralls
- Shoes and socks
- Chemical-resistant gloves made of any waterproof material

Non-Agricultural Use Requirements

The requirements in this box apply to uses of this product that are not within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries or greenhouses.

Applications on golf courses or around municipal or industrial (office park) buildings and to ornamentals not grown for resale are not within the scope of the Worker Protection Standard.

- Keep unprotected persons out of treated area until sprays have dried.

Storage and Disposal

Do not contaminate water, food, or feed by storage and disposal.

Storage: Keep away from fire and sparks. Do not allow this product to become wet or overheated in storage: decomposition, impaired activity, or fire may result. Keep container closed when not in use. Decomposition produces a foul odor; if observed, check for hot containers and immediately remove to open areas for disposal.

Pesticide Disposal: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

Container Handling: Nonrefillable container. Do not reuse or refill this container. Completely empty bag into application equipment. Offer for recycling if available or dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

Product Information

Fore[®] 80WP Rainshield[®] specialty fungicide is a unique and different broad-spectrum protectant fungicide that, when used according to label directions, provides high levels of disease control.

Optimum disease control is achieved when the fungicide is applied in a regularly scheduled preventative spray program. The addition of an spreader-sticker will improve fungicide performance by providing a more uniform spray deposit, increased foliar redistribution, and improved fungicide retention during periods of wet weather.

Chemigation: Do not apply this product through any type of irrigation system.

Aerial application by fixed or rotary wing aircraft on turf and ornamentals is prohibited.

Spray Drift Management

A variety of factors including weather conditions (e.g., wind direction, wind speed, temperature, relative humidity) and method of application (e.g., ground, airblast) can influence pesticide drift. The applicator must evaluate all factors and make appropriate adjustments when applying this product.

Wind Speed

Do not apply at wind speeds greater than 15 mph.

Temperature Inversions

If applying at wind speeds less than 3 mph, the applicator must determine if a) conditions of temperature inversion exist, or b) stable atmospheric conditions exist at or below nozzle height. Do not make applications into areas of temperature inversions or stable atmospheric conditions.

Other State and Local Requirements

Applicators must follow all state and local pesticide drift requirements regarding application of mancozeb. Where states have more stringent regulations, they must be observed.

Equipment

All ground application equipment must be properly maintained and calibrated using appropriate carriers or surrogates.

Do not apply with a nozzle height greater than 4 feet above the crop canopy.

Use Rate Determination

Carefully read, understand, and follow label use rates and restrictions. Under low disease conditions, minimum label rates per application can be used while maximum label rates and the minimum interval may be used for severe or threatening disease conditions.

For proper application to turfgrass, determine the square footage to be treated, divide the footage by 1000, multiply by the required fungicide use rate per 1000 sq ft, and then determine the amount of water required to provide adequate coverage. When treating ornamentals, determine the required fungicide use rate and the spray gallonage required to provide a thorough coverage. Prepare only the amount of spray solution required to treat the desired area.

Handling of Water Soluble Packets

The enclosed packets of Fore 80WP Rainshield are water soluble. Do not remove the water soluble packets from overwrap packaging except to add directly to the spray tank. Do not allow packets to become wet prior to adding to the spray tank. Do not handle the packets with wet hands or wet gloves. Do not open water soluble packets. Do not split packets. Partial use of packets is not allowed. Always reseal overwrap bag containing unopened water soluble packets to protect the remaining unused packets.

Mixing Directions

Be sure the sprayer is clean and not contaminated with other materials. When using an agitated spray tank, fill the tank 1/2 to 3/4 full with clean water and start agitation. Be certain that the agitation system is working properly and creates a rolling rippling on the liquid surface. Turn off the agitator and add the required number of unopened soluble packets of Fore 80WP Rainshield as determined by the required dosage to the tank - being careful not to drop on top of each other. Wait 5 to 10 minutes to allow the packets to thoroughly dissolve before turning on the agitation. Continue filling tank with the remainder of the water. Always add Fore 80WP Rainshield first and insure that it is in solution prior to adding any additional materials to the tank.

Compatibility

Fore 80WP Rainshield is compatible with boron and spray oils; however, the water soluble packets must be completely dissolved before adding spray oils or products containing boron to spray mixtures.

Fore 80WP Rainshield is compatible with most commonly used agricultural fungicides, insecticides and growth regulators. When preparing tank mixes, consult spray compatibility charts or State Cooperative Extension Service Specialists prior to actual use. Do not tank mix with any product which contains a prohibition on tank mixing. When two pesticides are tank mixed, the more restrictive label conditions apply.

The physical spray compatibility of specific tank mixes can be checked by preparing a sample suspension in a glass container and observing the suspension for any adverse effects, which might interfere with proper spray application.

Spray Adjuvants

The addition of a surfactant to sprays containing Fore 80WP Rainshield will improve initial spray deposits, fungicide redistribution and weatherability. Place Fore 80WP Rainshield into suspension prior to adding an adjuvant to the spray mixture. Read

and carefully observe the precautionary statements and all other information appearing on both product labels prior to spray preparation. When an adjuvant is to be used with this product, Dow AgroSciences recommends the use of a Chemical Producers and Distributors Association certified adjuvant.

Application Directions

Fore 80WP Rainshield is a protectant fungicide. Optimum disease control is achieved with thorough coverage sprays applied on a regular schedule.

Disease Scouting

Fore 80WP Rainshield is a broad-spectrum protectant fungicide. If not applied on a routine protectant spray schedule, frequently observe turfgrass and ornamental plants for disease signs or symptoms. Apply fungicide at the required label use rate and spray schedule, at the first sign of disease, report of disease in the area, or during environmental conditions favorable for disease development.

Turfgrass

For use on golf courses, and industrial or municipal turf areas. Not for use on sod farms, athletic fields and residential turf.

Restrictions:

Golf Courses:

- For cool season turfgrasses; greens, tees and aprons - limit to a maximum of five applications per year at a maximum application rate of 21.75 lb of product per acre (17.4 lb ai per acre) per application.
- For cool season turfgrasses; fairways - limit to a maximum of four applications per year at a maximum application rate of 21.75 lb of product per acre (17.4 lb ai per acre) per application.
- For warm season turfgrasses; fairways - limit to a maximum of three applications per year at a maximum application rate of 21.75 lb of product per acre (17.4 lb ai per acre) per application.

- For warm season turfgrasses; greens, tees and aprons - limit to a maximum of four applications per year at a maximum application rate of 21.75 lb of product per acre (17.4 lb ai per acre) per application.
- Require a minimum of a 10-day interval between applications.

All Other Turfgrass:

- Limit to a maximum of four applications per year and a maximum application rate of 21.75 lb of product per acre (17.4 lb ai per acre) per application
- Require a minimum of a 10-day interval between applications

Each 24 oz water soluble packet treats 6000 sq ft at the 4 oz per 1000 sq ft rate, 4000 sq ft at the 6 oz per 1000 sq ft rate, and 3000 sq ft at the 8 oz per 1000 sq ft rate. Note that the 8 oz per 1000 sq ft rate is the same as the maximum rate of 21.75 lb per acre.

Start application when grass greens up in spring or when disease first appears and repeat at 10- to 14-day intervals or until disease threat is passed. When conditions are especially favorable for disease development, apply maximum fungicide use rate on a 10-day spray schedule. Apply in sufficient water to provide adequate coverage.

Turfgrass Tolerance

Maintain treated turfgrass in a vigorous growing condition. Turfgrass under stress will not respond as well to fungicide treatments as well-maintained turfgrass. Turfgrass tolerance to this product has been found to be acceptable; however, this product and tank mixtures with other products have not been tested on all varieties of every turfgrass species or under all possible growing conditions. If user is unfamiliar with the performance of Fore 80WP Rainshield or tank mixtures, under user growing conditions, treat a limited area of turfgrass prior to initiating large-scale applications.

Crop	Diseases Controlled	Fore 80WP Rainshield (oz/1000 sq ft)	Remarks (Also refer to directions for use)	Restrictions
assorted grasses	helminthosporium melting-out rust (leaf, stem, stripe)	4		<ul style="list-style-type: none"> • Do not graze treated areas. • Do not use on turfgrasses intended for grazing, such as range or pasture grasses. • Do not feed grass clippings to livestock. • Do not use on grasses grown for seed.
	copper spot fusarium blight red thread slime mold	4 - 8		
	algae	6		
	dollar spot	6 - 8		
	rhizoctonia brown patch	4	Apply on a 10-day spray schedule.	
	pythium blight	8	Apply at 10-day intervals	
	rapid blight (a putative chytridiomycete fungus)	8	Apply at a 14-day interval if conditions are favorable for disease development.	
	fusarium snow mold	6 - 8	Apply at 2- to 6-week intervals during winter.	
	gray leaf spot	8	Apply at 14-day intervals if conditions are favorable for disease development.	

Ornamentals

Restrictions:

- Cut flowers and greenhouse grown ornamentals: Limit to 20 applications per year.
- Do not use any part of treated plants, including fruit, nuts, bark, or sap, for food or feed purposes.
- Do not use on pachysandra.
- Do not apply more than a total of 8.4 lb of Fore 80WP Rainshield per acre per year.
- Not for use in residential greenhouses.

Neither the manufacturer nor the seller has determined the effects of using Fore 80WP Rainshield on ornamentals not specified on this label. Prior to any large-scale applications on such plants, the user should determine the effects of Fore 80WP Rainshield by testing a small section of the type of plants treated. User assumes all risk arising out of application to unlabeled plants. The Conditions of Sale and Warranty apply to all uses.

For outdoor or greenhouse use, apply the equivalent of 1 1/2 lb of Fore 80WP Rainshield per 100 gallons of dilute spray (1 1/2 lb of Fore 80WP Rainshield per acre). The addition of a surfactant to spray solutions will improve performance.

Begin spraying when plants are well leafed out or at first sign of disease in a full coverage spray at 7- to 10-day intervals throughout season or follow State Extension Service recommended retreatment intervals for disease control on the following ornamental plants:

Crop	Diseases Controlled	Crop Specific Use Directions
African violet	botrytis blight	
ageratum	botrytis blight rust	
almond (ornamental)	leaf spot	
alyssum		
anthurium	anthracnose, spadix rot	
apple (ornamental)	fabraea leaf spot rust scab	
arborvitae	cercospora blight	
areca palm	leaf spot	
ash, mountain	entomosporium leaf spot guignardia leaf blotch	
ash, white	anthracnose cylindrosporium leaf spot	
aster	leaf spot	
aster, perennial	puccinia rusts	

aucuba, japonica	alternaria leaf spot anthracnose	
azalea	cyindrocladium rot petal blight phytophthora twig and bud blight	Apply in a full coverage spray, 2 to 3 times a week, while flowers are opening. Direct spray into flowers and thoroughly spray ground under bushes.
begonia	botrytis blight	
bougainvillea	leaf spot	
boxwood		
buffaloberry	cyindrosporium leaf spot	
camellias	petal blight	Refer to azalea.
carnation	rust septoria leaf spot	
cedar, red (juniper)	cercospora blight phomopsis blight	
chrysanthemum	ascochyta blight botrytis petal spot rust	Apply twice weekly during blooming period.
cockscomb (<i>Celosia</i>)	alternaria leaf spot	

Crop (Cont.)	Diseases Controlled	Crop Specific Use Directions
conifers (Christmas trees)	lophodermium needle cast pine gall rust scirrhia brown spot	Begin application in spring or early summer before infection occurs. Repeat after heavy rains and at two- week intervals as long as needed.
cordyline	cercospora leaf spot	
crabapple (ornamental)	cedar-apple rust scab sphaeropsis leaf spot	
cypress, Arizona (<i>Cupressus</i> sp.)	cercospora blight monochaetia canker	
dahlia	botrytis blight	
delphinium		
dieffenbachia	leptosphaeria brown spot	
dogwood, flowering	anthracnose elsinoe leaf spot septoria leaf spot	Apply when buds begin to open, when bracts have fallen, 4 weeks later and again in late summer after flower buds for next season have formed.
dracaena	fusarium leaf spot	
elm	black leaf spot	

euonymus	anthracnose	
fatsia		
fern	rhizoctonia blight	
figus	cercospora leaf spot	
fig	cyllindrocladium leaf spot	
firethorn (<i>pyracantha</i>)	fusicladium scab	
fir, Douglas	Swiss needle cast	
fir, fraser		
fuchsia	botrytis blight rust	
geranium	rust	
gladiolus	botrytis blossom blight curvularia leaf spot	Make regular weekly applications starting before diseases appear and increase to 2 or 3 applications per week during periods of heavy disease and during rainy weather. On flower spikes, reduce spray concentration to 3/4 lb per 100 gallons.
gloxinia	botrytis blight	
gypsophila		

Crop (Cont.)	Diseases Controlled	Crop Specific Use Directions
hawthorn	cedar-apple rust fabraea leaf spot frogeye leaf spot hawthorn rust scab	
hickory	gnomonium leaf spot	
holly	purple spot	
hollyhock	anthracnose cercospora leaf spot puccinia rust	
honeysuckle	herpobasidium blight	
horsechestnut, buckeye	alternaria leaf spot guignardia leaf blotch	
hydrangea	botrytis blight cercospora leaf spot	
impatiens	botrytis blight	
iris	didymella leaf spot mycosphaerella leaf spot mystrosporium ink spot	(formerly didymella)

juniper	phomopsis blight	
larkspur	rust	
laurel, mountain	cercospora leaf spot petal blight	Refer to azalea.
ligustrum	cercospora leaf spot	
lily	botrytis blight	
magnolia	gloeosporium leaf spot	
maple	alternaria leaf spot phyllosticta leaf spot	
marigold	botrytis blossom blight	Do not use on French dwarf double or Signet type marigold seedlings.
narcissus	botrytis blight (fire) smoulder	
oak	actinopelte leaf spot taphrina leaf blister	
orchid (<i>Dendrobium</i>)	botrytis blossom blight	
oxalis	rust	

Crop (Cont.)	Diseases Controlled	Crop Specific Use Directions
pansy	anthracnose	
pears (ornamental)	fabraea leaf spot rust scab	
peony	botrytis blossom blight phytophthora blight	Apply in early spring and early fall, drenching soil around plants as well as the foliage. Promptly destroy all infected plant parts.
peperomia	cercospora leaf spot	
petunia	botrytis blight	
philodendron	dactylaria leaf spot phytophthora leaf spot	
phlox	leaf spot	
photinia	entomosporium leaf spot	
pine, Australia	cyclaneusma needle cast	
pine, Scotch	cyclaneusma needle cast gall rust	
pittosporum	alternaria leaf spot	
pleomele	fusarium leaf spot	
poinsettia	sphaceloma scab	

poplar	rust	
primrose	botrytis blight	
protea		
quince (ornamental)	fabraea leaf spot rust scab	
rhododendron	cercospora leaf spot discosia leaf spot petal blight	Refer to azalea.
rose	black spot cercospora leaf spot rust	
rosemary	rhizoctonia aerial blight	
schefflera	alternaria blight	
Scotts pine	needle cast	
skunkbush, sumac	cyindrosporium leaf spot	
snapdragon	rust	

Crop (Cont.)	Diseases Controlled	Crop Specific Use Directions
spathiphyllum	myrothecium leaf spot	
statice	cercospora frogeye	
strawflower	rust	
syngonium	cephalosporium leaf spot	
thorn apple	rust	
tulip	botrytis blight (fire)	
venus, flytrap	anthracnose	
viburnum	downy mildew ramularia leaf spot	
walnut	anthracnose	
zinnia	alternaria leaf blight	

ATTENTION: This product contains mancozeb and ETU, chemicals known to the State of California to cause cancer. ETU is also known to the State of California to cause birth defects or other reproductive harm.

Terms and Conditions of Use

If terms of the following Warranty Disclaimer, Inherent Risks of Use and Limitation of Remedies are not acceptable, return unopened package at once to the seller for a full refund of purchase price paid. To the extent permitted by law, otherwise, use by the buyer or any other user constitutes acceptance of the terms under Warranty Disclaimer, Inherent Risks of Use and Limitation of Remedies.

Warranty Disclaimer

Dow AgroSciences warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions, subject to the inherent risks set forth below. TO THE EXTENT PERMITTED BY LAW, Dow AgroSciences MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.

Inherent Risks of Use

It is impossible to eliminate all risks associated with use of this product. Crop injury, lack of performance, or other unintended consequences may result because of such factors as use of the product contrary to label instructions (including conditions noted on the label, such as unfavorable temperatures, soil conditions, etc.), abnormal conditions (such as excessive rainfall, drought, tornadoes, hurricanes), presence of other materials, the manner of application, or other factors, all of which are beyond the control of Dow AgroSciences or the seller. To the extent permitted by law, all such risks shall be assumed by buyer.

Limitation of Remedies

To the extent permitted by law, the exclusive remedy for losses or damages resulting from this product (including claims based on contract, negligence, strict liability, or other legal theories), shall be limited to, at Dow AgroSciences' election, one of the following:

1. Refund of purchase price paid by buyer or user for product bought, or
2. Replacement of amount of product used.

To the extent permitted by law, Dow AgroSciences shall not be liable for losses or damages resulting from handling or use of this product unless Dow AgroSciences is promptly notified of such loss or damage in writing. To the extent permitted by law, in no case shall Dow AgroSciences be liable for consequential or incidental damages or losses.

The terms of the Warranty Disclaimer, Inherent Risks of Use and Limitation of Remedies cannot be varied by any written or verbal statements or agreements. No employee or sales agent of Dow AgroSciences or the seller is authorized to vary or exceed the terms of the Warranty Disclaimer or Limitation of Remedies in any manner.

®Trademark of The Dow Chemical Company ("Dow") or an affiliated company of Dow
EPA accepted 06/05/13

Fore[®] 80WP Rainshield[®]

SPECIALTY FUNGICIDE

Active Ingredients

mancozeb: A coordination product of zinc ion
and manganese ethylene bisdithiocarbamate..... 80.0%

In which the ingredients are:

Manganese⁺⁺ 16.00%

Zinc⁺⁺ 2.00%

Ethylene bisdithiocarbamate
ion (C₄H₆N₂S₄)⁻⁻ 62.00%

Inert Ingredients 20.0%

Total 100.0%

Keep Out of Reach of Children

CAUTION

Refer to label booklet for Directions for Use including
Storage and Disposal.

EPA Reg. No. 62719-388

300-001955

[®]Trademark of The Dow Chemical Company ("Dow") or an affiliated
company of Dow

NET WEIGHT 1.5 LB

PLACE
BOOKLET
HERE

First Aid
If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.
If on skin or clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-992-5994, day or night, for emergency treatment information.

Precautionary Statements and Domestic Animals

CAUTION
Harmful if Absorbed Through The Skin • Causes Moderate Eye Irritation • Avoid contact with skin, eyes or clothing • Prolonged Or Frequently Repeated Skin Contact May Cause Allergic Reactions in Some Individuals

Personal Protective Equipment (PPE)

Some materials that are chemical-resistant to this product and butyl rubber and neoprene rubber. If you want more options, follow the instructions for category "A," on an EPA chemical resistance category selection chart.

Mixers, loaders, applicators and all other handlers must wear:

- Long-sleeved shirt
- Long pants
- Shoes and socks
- Chemical resistant gloves made of any waterproof material (except for pilots)

In addition, all handlers (except pilots, groundboom applicators, and airblast applicators) must wear:

- A NIOSH-approved respirator with a dust/mist filter with MSHA/NIOSH approval number prefix TC-21C or any R, P or HE filter.
- Use an R, P or HE filter when mixing Fore 80WP Rainshield with an oil-based adjuvant.

See engineering controls for additional requirements.

User Safety Requirements

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry. Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them.

Engineering Controls

Water-soluble packets when used correctly qualify as a closed mixing/loading system under the Worker Protection Standard for Agricultural Pesticides [40 CFR 170.170-240(d)(4)]. Mixers and loaders using water-soluble packets must:

- Wear the personal protective equipment required in the PPE section of this labeling for mixers/loaders
- Be provided, and have immediately available, and wear in an emergency, such as a broken package, spill, or equipment breakdown; a NIOSH-approved respirator with a dust/mist filter with MSHA/NIOSH approval number prefix TC-21C or any R, P or HE filter.

User Safety Recommendations

- Users should:
- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
 - Remove clothing/PPE immediately if pesticide gets inside, then wash thoroughly and put on clean clothing.
 - Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

Environmental Hazards

This pesticide is toxic to aquatic organisms. Drift and runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when cleaning equipment or disposing of equipment washwaters or rinsate.

Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. Refer to the label booklet under "Agricultural Use Requirements" in the Directions for Use section for information about this standard.

Storage and Disposal

Do not contaminate water, food, or feed by storage and disposal.

Storage: Keep away from fire and sparks. Do not allow this product to become wet or overheated in storage. decomposition, impaired activity, or fire may result. Keep container closed when not in use. Decomposition produces a foul odor; if observed, check for hot containers and immediately remove to open areas for disposal.

Pesticide Disposal: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

Container Handling: Nonrefillable container. Do not reuse or refill this container. Completely empty bag into application equipment. Offer for recycling if available or dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

Refer to label booklet for Directions for Use.



NET WEIGHT 4 X 1.5 LB WATER-SOLUBLE PACKETS

MADE IN COLOMBIA

®Trademark of The Dow Chemical Company ("Dow") or an affiliated company of Dow

Produced for
Dow AgroSciences LLC
9330 Zionsville Road
Indianapolis, IN 46268

EPA Reg. No. 62719-388 EPA Est. 67545-AZ-001
900-022164 / 00187718

Agricultural Chemical: Do not ship or store with food, feeds, drugs or clothing.

In case of emergency endangering health or the environment involving this product, call 1-800-992-5994.

once unopened.

Remedies at end of label booklet. If terms are unacceptable, return at once unopened.

Disclaimer, Inherent Risks of Use, and Limitation of Remedies at end of label booklet. If terms are unacceptable, return at once unopened.



Notice: Read the entire label. Use only according to label directions. Before using this product, read Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies at end of label booklet. If terms are unacceptable, return at once unopened.

SPECIALTY FUNGICIDE

80WP Rainshield®

Fore®



Dow AgroSciences

Keep Out of Reach of Children CAUTION

Total 100.0%

Other Ingredients:..... 20.0%

Ethylene bisdithiocarbamate ion (C₄H₈N₂S₂)..... 62.00%

Zinc⁺⁺ 2.00%

Manganese⁺⁺ 16.00%

In which the ingredients are:

zinc ion and manganese ethylene bisdithiocarbamate 80.0%

Active Ingredients

Mancozeb: A coordination product of zinc ion and manganese ethylene bisdithiocarbamate 80.0%

Product in Water-Soluble Packets

Group M3 FUNGICIDE

Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. Refer to the label booklet under "Agricultural Use Requirements" in the Directions for Use section for information about this standard.

Refer to inside of label booklet for additional precautionary information including Directions for Use.

14.5" Repeat

TOP OF BACK

.375"

.375"

.375"

.4375"

Zipper C/L 1.5" below top of pouch.

21" Finished Web Width

1"

1" NO COPY AREA
BOTTOM OF BAG

20-13/16

16-3/8

75-3/4

20-7/8

16-5/16

1-3/8

Do Not Cut To Open
Contents In Plastic



Fore[®] 80WP Rainshield[®]

SPECIALTY FUNGICIDE

Product in Water-Soluble Packets

Group	M3	FUNGICIDE
Active Ingredients		

Mancozeb: A coordination product of zinc ion and manganese ethylene bisdithiocarbamate.....	80.00%	
In which the ingredients are:		
Manganese ⁺⁺	16.00%	
Zinc ⁺⁺	2.00%	
Ethylene bisdithiocarbamate ion (C ₂ H ₄ N ₂ S ₂).....	62.00%	
Other Ingredients.....	20.00%	
Total.....	100.00%	

Keep Out of Reach of Children

CAUTION

Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. Refer to the label booklet under "Agricultural Use Requirements" in the Directions for Use section for information about this standard.

Refer to inside of label booklet for additional precautionary information including Directions for Use.

Notice: Read the entire label. Use only according to label directions. Before using this product, read Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies at end of label booklet. If terms are unacceptable, return at once unopened.

In case of emergency endangering health or the environment involving this product, call 1-800-992-5994.

Agricultural Chemical: Do not ship or store with food, feeds, drugs or clothing.

EPA Reg. No. 62719-388

EPA Est. 67545-AZ-001

MADE IN COLOMBIA

[®]Trademark of The Dow Chemical Company ("Dow") or an affiliated company of Dow

Produced for
Dow AgroSciences LLC
9330 Zionsville Road
Indianapolis, IN 46268

8 X 4 X 1.5 LB WATER-SOLUBLE PACKETS

00172874



Fore[®] 80WP Rainshield[®]

SPECIALTY FUNGICIDE



8 X 4 X 1.5 LB WATER-SOLUBLE PACKETS




Fore[®] 80WP Rainshield[®]

SPECIALTY FUNGICIDE



8 X 4 X 1.5 LB WATER-SOLUBLE PACKETS

00172874



4G/Z24.2/S/14
USA+/AA4992



Fore[®] 80WP Rainshield[®]

SPECIALTY FUNGICIDE



8 X 4 X 1.5 LB WATER-SOLUBLE PACKETS

900-022163 / 00187716
BOX MFG. DATE:



Dow AgroSciences

Fore[®]

80WP Rainshield[®]

SPECIALTY FUNGICIDE



Department of Agriculture
STATE OF HAWAII

LICENSED

PERIOD 2014-2016 LIC. NO.

9786.230



Group	M3	FUNGICIDE
-------	----	-----------

Active Ingredients

Mancozeb: A coordination product of zinc ion and manganese ethylene bisdithiocarbamate80.0%

In which the ingredients are:

Manganese⁺⁺ 16.00%
 Zinc⁺⁺ 2.00%
 Ethylene bisdithiocarbamate ion (C₄H₆N₂S₄) 62.00%
 Other Ingredients 20.0%
 Total 100.0%

Notice: Read the entire label. Use only according to label directions. **Before using this product, read Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies at end of label booklet. If terms are unacceptable, return at once unopened.**

In case of emergency endangering health or the environment involving this product, call 1-800-992-5994.

Agricultural Chemical: Do not ship or store with food, feeds, drugs or clothing.

EPA Reg. No. 62719-388



(01) 0 06 62974 47381 7

Keep Out of Reach of Children **CAUTION**

Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. Refer to the label booklet under "Agricultural Use Requirements" in the Directions for Use section for information about this standard.

Refer to back panel of bag for additional precautionary information including Personal Protective Equipment (PPE), User Safety Recommendations and Directions for Use including Storage and Disposal.

MADE IN COLOMBIA

® Trademark of The Dow Chemical Company ("Dow") or an affiliated company of Dow

**Produced for
Dow AgroSciences LLC
9330 Zionsville Road
Indianapolis, IN 46268**

NET WEIGHT 25 KG / 55 LB

**ENVIRONMENTALLY HAZARDOUS
SUBSTANCES, SOLID, N.O.S.
(MANCOZEB) UN3077**



**NOT REGULATED FOR
DOMESTIC SURFACE
TRANSPORTATION WITHIN
THE U.S.A.**



Table of Contents	Page
Precautionary Statements	1
Hazards to Humans and Domestic Animals	1
Personal Protective Equipment (PPE).....	1
User Safety	1
Environmental Hazards.....	1
Directions for Use.....	1
Agricultural Use Requirements	1
Non-Agricultural Use Requirements	1
Storage and Disposal	1
Product Information	2
Spray Drift Management.....	2
Use Rate Determination.....	2
Mixing Directions	2
Compatibility.....	3
Spray Adjuvants.....	3
Application Directions.....	3
Disease Scouting.....	3
Ornamentals	3
Terms and Conditions of Use	4
Warranty Disclaimer	4
Inherent Risks of Use	4
Limitation of Remedies.....	5

First Aid

If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

If on skin or clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-992-5994 day or night, for emergency treatment information.

Precautionary Statements

Hazards to Humans and Domestic Animals

CAUTION

Harmful If Absorbed Through The Skin • Causes Moderate Eye Irritation • Avoid contact with skin, eyes or clothing • Prolonged Or Frequently Repeated Skin Contact May Cause Allergic Reactions In Some Individuals

Personal Protective Equipment (PPE)

Some materials that are chemical-resistant to this product are butyl rubber and neoprene rubber. If you want more options, follow the instructions for category A on an EPA chemical resistance category selection chart.

Mixers, loaders, applicators and all other handlers must wear:

- Long-sleeved shirt
- Long pants
- Shoes and socks
- Chemical-resistant gloves made of any waterproof material (except for pilots)

In addition, all handlers (except pilots, groundboom applicators, and airblast applicators) must wear:

- A NIOSH-approved respirator with a dust/mist filter with MSHA/NIOSH approval number prefix TC-21C or any R, P or HE filter.
- Use an R, P or HE filter when mixing Fore 80WP Rainshield with an oil-based adjuvant.

See engineering controls for additional requirements

User Safety Requirements

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry. Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them.

Engineering Controls

When handlers use closed systems or enclosed cabs in a manner that meet the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d) (5)], the handler PPE requirements may be reduced or modified as specified in the WPS.

User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside, then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

Environmental Hazards

This pesticide is toxic to aquatic organisms. Drift and runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when cleaning equipment or disposing of equipment washwaters or rinsate.

Directions for Use

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your state or tribe, consult the agency responsible for pesticide regulation.

Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 24 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil or water, is:

- Coveralls
- Shoes and socks
- Chemical-resistant gloves made of any waterproof material

Non-Agricultural Use Requirements

The requirements in this box apply to uses of this product that are not within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries or greenhouses.

Applications on golf courses or around municipal or industrial (office park) buildings and to ornamentals not grown for resale are not within the scope of the Worker Protection Standard.

- Keep unprotected persons out of treated area until sprays have dried.

Storage and Disposal

Do not contaminate water, food, or feed by storage and disposal.

Storage: Keep away from fire and sparks. Do not allow this product to become wet or overheated in storage: decomposition, impaired activity, or fire may result. Keep container closed when not in use. Decomposition produces a foul odor; if observed, check for hot containers and immediately remove to open areas for disposal.

Pesticide Disposal: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

Container Handling: Nonrefillable container. Do not reuse or refill this container. Completely empty bag into application equipment. Offer for recycling if available or dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

Product Information

Read all Directions for Use carefully before applying.

Fore® 80WP Rainshield® is a unique and different specialty broad-spectrum protectant fungicide that, when used according to label directions, provides high levels of disease control.

Optimum disease control is achieved when the fungicide is applied in a regularly scheduled preventative spray program. The addition of an spreader-sticker will improve fungicide performance by providing a more uniform spray deposit, increased foliar redistribution, and improved fungicide retention during periods of wet weather.

Chemigation: Do not apply this product through any type of irrigation system.

Aerial application by fixed or rotary wing aircraft on ornamentals is prohibited.

Spray Drift Management

A variety of factors including weather conditions (e.g., wind direction, wind speed, temperature, relative humidity) and method of application (e.g., ground, airblast) can influence pesticide drift. The applicator must evaluate all factors and make appropriate adjustments when applying this product.

Wind Speed

Do not apply at wind speeds greater than 15 mph.

Temperature Inversions

If applying at wind speeds less than 3 mph, the applicator must determine if a) conditions of temperature inversion exist, or b) stable atmospheric conditions exist at or below nozzle height. Do not make applications into areas of temperature inversions or stable atmospheric conditions.

Other State and Local Requirements

Applicators must follow all state and local pesticide drift requirements regarding application of mancozeb. Where states have more stringent regulations, they must be observed.

Equipment

All ground application equipment must be properly maintained and calibrated using appropriate carriers or surrogates.

Do not apply with a nozzle height greater than 4 feet above the crop canopy.

Use Rate Determination

Carefully read, understand, and follow label use rates and restrictions. Under low disease conditions, minimum label rates per application can be used while maximum label rates and the minimum interval may be used for severe or threatening disease conditions.

When treating ornamentals, determine the required fungicide use rate and the spray gallonage required to provide a thorough coverage. Prepare only the amount of spray solution required to treat the desired area.

Mixing Directions

Be sure sprayer is clean and not contaminated with other materials prior to use. When using an agitated spray tank, fill the tank 1/2 to 3/4 full with clean water and start agitation. Be certain that the agitation system is working properly and creates a rolling rippling on the liquid surface. With the agitator running add the required amount of Fore 80WP Rainshield to the tank. Continue filling tank with the remainder of the water.

Slowly pour the appropriate amount of Fore 80WP Rainshield into a small container containing an equal volume of water while mixing. Mix until the Fore 80WP Rainshield is thoroughly wetted. Add additional water if necessary to make solution flowable. Add the contents of the slurry tank to a 1/2 filled sprayer, continue filling tank with remainder of water and mix well. Always add Fore 80WP Rainshield to spray mixture prior to adding any additional materials to the tank.

Compatibility

Fore 80WP Rainshield is compatible with most commonly used agricultural fungicides, insecticides and growth regulators. When preparing tank mixes, consult spray compatibility charts or State Cooperative Extension Service Specialists prior to actual use. Do not tank mix with any product which contains a prohibition on tank mixing. When two pesticides are tank mixed, the more restrictive label conditions apply.

The physical spray compatibility of specific tank mixes can be checked by preparing a sample suspension in a glass container and observing the suspension for any adverse effects, which might interfere with proper spray application.

Spray Adjuvants

The addition of a surfactant to sprays containing Fore 80WP Rainshield will improve initial spray deposits, fungicide redistribution and weatherability. Place Fore 80WP Rainshield into suspension prior to adding an adjuvant to the spray mixture. Read and carefully observe the precautionary statements and all other information appearing on both product labels prior to spray preparation. When an adjuvant is to be used with this product, Dow AgroSciences recommends the use of a Chemical Producers and Distributors Association certified adjuvant.

Application Directions

Fore 80WP Rainshield is a protectant fungicide. Optimum disease control is achieved with thorough coverage sprays applied on a regular schedule.

Disease Scouting

Fore 80WP Rainshield is a broad-spectrum protectant fungicide. If not applied on a routine protectant spray schedule, frequently observe ornamental plants for disease signs or symptoms. Apply fungicide at the required label use rate and spray schedule, at the first sign of disease, report of disease in the area, or during environmental conditions favorable for disease development.

Ornamentals

Restrictions:

- Cut flowers and greenhouse grown ornamentals: Limit to 20 applications per year.
- Do not use any part of treated plants, such as fruit, nuts, bark or sap, for food or feed purposes.
- Do not use on pachysandra.
- Do not apply more than a total of 8.4 lb of Fore 80WP Rainshield per acre per year.
- Not for use in residential greenhouses.

Neither the manufacturer nor the seller has determined the effects of using Fore 80WP Rainshield on ornamentals not specified on this label. Prior to any large-scale applications on such plants, the user should determine the effects of Fore 80WP Rainshield by testing a small section of the type of plants treated. User assumes all risk arising out of application to unlabeled plants. The Conditions of Sale and Warranty apply to all uses.

For outdoor or greenhouse use, apply the equivalent of 1 1/2 lb of Fore 80WP Rainshield per 100 gallons of dilute spray (1 1/2 lb of Fore 80WP Rainshield per acre). The addition of a surfactant to spray solutions will improve performance.

Begin spraying when plants are well leaved out or at first sign of disease in a full coverage spray at 7- to 10-day intervals throughout season or follow State Extension Service recommended retreatment intervals for disease control on the following ornamental plants:

Crop	Diseases Controlled	Crop Specific Use Directions
African violet	botrytis blight	
ageratum	botrytis blight rust	
almond (ornamental)	leaf spot	
alyssum		
anthurium	anthracnose, spadix rot	
apple (ornamental)	fabraea leaf spot rust scab	
arborvitae	cercospora blight	
areca palm	leaf spot	
ash, mountain	entomosporium leaf spot guignardia leaf blotch	
ash, white	anthracnose cylindrosporium leaf spot	
aster	leaf spot	
aster, perennial	puccinia rusts	
aucuba, japonica	alternaria leaf spot anthracnose	
azalea	cylindrocladium rot petal blight phytophthora twig and bud blight	Apply in a full coverage spray, 2 to 3 times a week, while flowers are opening. Direct spray into flowers and thoroughly spray ground under bushes.
begonia	botrytis blight	
bougainvillea	leaf spot	
boxwood		
buffaloberry	cylindrosporium leaf spot	
camellias	petal blight	Refer to azalea.
carnation	rust septoria leaf spot	
cedar, red (juniper)	cercospora blight phomopsis blight	

chrysanthemum	ascochyta blight botrytis petal spot rust	Apply twice weekly during blooming period.
cockscomb (<i>Celosia</i>)	alternaria leaf spot	
conifers (Christmas trees)	lophodermium needle cast pine gall rust scirrhia brown spot	Begin application in spring or early summer before infection occurs. Repeat after heavy rains and at two-week intervals as long as needed.
cordyline	cercospora leaf spot	
crabapple (ornamental)	cedar-apple rust scab sphaeropsis leaf spot	
cypress, Arizona (<i>Cupressus</i> sp.)	cercospora blight monochaetia canker	
dahlia	botrytis blight	
delphinium		
dieffenbachia	leptosphaeria brown spot	
dogwood, flowering	anthracnose elsinoe leaf spot septoria leaf spot	Apply when buds begin to open, when bracts have fallen, 4 weeks later and again in late summer after flower buds for next season have formed.
dracaena	fusarium leaf spot	
elm	black leaf spot	
euonymus	anthracnose	
fatsia		
fern	rhizoctonia blight	
figus	cercospora leaf spot	
fig	cyllindrocladium leaf spot	
firethorn (<i>pyracantha</i>)	fusicladium scab	
fir, Douglas	Swiss needle cast	
fir, fraser		
fuchsia	botrytis blight rust	
geranium	rust	

Crop (Cont.)	Diseases Controlled	Crop Specific Use Directions
gladiolus	botrytis blossom blight curvularia leaf spot	Make regular weekly applications starting before diseases appear and increase to 2 or 3 applications per week during periods of heavy disease and during rainy weather. On flower spikes, reduce spray concentration to 3/4 lb per 100 gallons.
gloxinia	botrytis blight	
gypsophila		
hawthorn	cedar-apple rust fabraea leaf spot frogeye leaf spot hawthorn rust scab	
hickory	gnomonium leaf spot	
holly	purple spot	
hollyhock	anthracnose cercospora leaf spot puccinia rust	
honeysuckle	herpobasidium blight	
horsechestnut, buckeye	alternaria leaf spot guignardia leaf blotch	
hydrangea	botrytis blight cercospora leaf spot	
impatiens	botrytis blight	
iris	didymellina leaf spot mycosphaerella leaf spot mystrosporium ink spot	(formerly didymellina)
juniper	phomopsis blight	
larkspur	rust	
laurel, mountain	cercospora leaf spot petal blight	Refer to azalea.
ligustrum	cercospora leaf spot	
lily	botrytis blight	
magnolia	gloeosporium leaf spot	
maple	alternaria leaf spot phyllosticta leaf spot	
marigold	botrytis blossom blight	Do not use on French dwarf double or Signet type marigold seedlings.
narcissus	botrytis blight (fire) smoulder	
oak	actinopelte leaf spot taphrina leaf blister	

orchid (<i>Dendrobium</i>)	botrytis blossom blight	
oxalis	rust	
pansy	anthracnose	
pears (ornamental)	fabraea leaf spot rust scab	
peony	botrytis blossom blight phytophthora blight	Apply in early spring and early fall, drenching soil around plants as well as the foliage. Promptly destroy all infected plant parts.
peperomia	cercospora leaf spot	
petunia	botrytis blight	
philodendron	dactylaria leaf spot phytophthora leaf spot	
phlox	leaf spot	
photinia	entomosporium leaf spot	
pine, Australia	cyclaneusma needle cast	
pine, Scotch	cyclaneusma needle cast gall rust	
pittosporum	alternaria leaf spot	
pleomele	fusarium leaf spot	
poinsettia	sphaceloma scab	
poplar	rust	
primrose	botrytis blight	
protea		
quince (ornamental)	fabraea leaf spot rust scab	
rhododendron	cercospora leaf spot discosia leaf spot petal blight	Refer to azalea.
rose	black spot cercospora leaf spot rust	
rosemary	rhizoctonia aerial blight	
schefflera	alternaria blight	
Scotts pine	needle cast	
skunkbush, sumac	cylindrosporium leaf spot	
snapdragon	rust	
spathiphyllum	myrothecium leaf spot	
statice	cercospora frogeye	
strawflower	rust	
syngonium	cephalosporium leaf spot	
thorn apple	rust	

tulip	botrytis blight (fire)	
venus, flytrap	anthracnose	
viburnum	downy mildew ramularia leaf spot	
walnut	anthracnose	
zinnia	alternaria leaf blight	

ATTENTION: This product contains mancozeb and ETU, chemicals known to the State of California to cause cancer. ETU is also known to the State of California to cause birth defects or other reproductive harm.

Terms and Conditions of Use

If terms of the following Warranty Disclaimer, Inherent Risks of Use and Limitation of Remedies are not acceptable, return unopened package at once to the seller for a full refund of purchase price paid. To the extent permitted by law, otherwise, use by the buyer or any other user constitutes acceptance of the terms under Warranty Disclaimer, Inherent Risks of Use and Limitation of Remedies.

Warranty Disclaimer

Dow AgroSciences warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions, subject to the inherent risks set forth below. **TO THE EXTENT PERMITTED BY LAW, Dow AgroSciences MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.**

Inherent Risks of Use

It is impossible to eliminate all risks associated with use of this product. Crop injury, lack of performance, or other unintended consequences may result because of such factors as use of the product contrary to label instructions (including conditions noted on the label, such as unfavorable temperatures, soil conditions, etc.), abnormal conditions (such as excessive rainfall, drought, tornadoes, hurricanes), presence of other materials, the manner of application, or other factors, all of which are beyond the control of Dow AgroSciences or the seller. To the extent permitted by law, all such risks shall be assumed by buyer.

Limitation of Remedies

To the extent permitted by law, the exclusive remedy for losses or damages resulting from this product (including claims based on contract, negligence, strict liability, or other legal theories), shall be limited to, at Dow AgroSciences' election, one of the following:

1. Refund of purchase price paid by buyer or user for product bought, or
2. Replacement of amount of product used.

To the extent permitted by law, Dow AgroSciences shall not be liable for losses or damages resulting from handling or use of this product unless Dow AgroSciences is promptly notified of such loss or damage in writing. To the extent permitted by law, in no case shall Dow AgroSciences be liable for consequential or incidental damages or losses.

The terms of the Warranty Disclaimer, Inherent Risks of Use and Limitation of Remedies cannot be varied by any written or verbal statements or agreements. No employee or sales agent of Dow AgroSciences or the seller is authorized to vary or exceed the terms of the Warranty Disclaimer or Limitation of Remedies in any manner.

®Trademark of The Dow Chemical Company ("Dow") or an affiliated company of Dow

EPA accepted 06/05/13

00173602



Dow AgroSciences

Fore[®] 80WP Rainshield[®]

SPECIALTY FUNGICIDE



300-002007