



Department of Agriculture  
STATE OF HAWAII



Department of Agriculture  
STATE OF HAWAII

**LICENSED**

PERIOD **2008-2010** LIC. NO.



**LICENSED**

PERIOD **2011-2013** LIC. NO.



Department of Agriculture  
STATE OF HAWAII

**LICENSED**

PERIOD **2014-2016** LIC. NO.

**9786.276**

# Accord<sup>®</sup> XRT II

## Herbicide

A non-selective broad-spectrum systemic herbicide for control of annual and perennial weeds and woody plants in noncrop areas and industrial sites, such as utility, railroad and roadside rights-of-way, airports, natural and production (plantations) forests for site preparation, mid-rotation release treatments, and timber stand improvement activities, wildlife and habitat management areas, wildlife openings, natural areas, such as wildlands, campgrounds, parks and recreational areas, wildlife refuges, rangeland and in and around seasonally dry wetlands including ditchbanks, dry ditches and dry canals and grazed areas on listed noncrop sites

**Avoid contact of herbicide with foliage, green stems, exposed non-woody roots or fruit of crops, desirable plants and trees, because severe injury or destruction may result.**

Active Ingredient:

glyphosate: N-(phosphonomethyl)glycine,	
dimethylamine salt .....	50.2%
Other Ingredients .....	49.8%
Total Ingredients .....	100.0%

Contains 5.4 pounds per gallon glyphosate, dimethylamine salt (4 pounds per gallon glyphosate acid).

**Keep Out of Reach of Children**  
**CAUTION**

### Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. Refer to the label booklet under "Agricultural Use Requirements" in the Directions for Use section for information about this standard.

**For additional Precautionary Statements, First Aid, Storage and Disposal and other use information see inside this label.**

**Notice:** Read the entire label. Use only according to label directions. **Before using this product, read Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies at end of label booklet. If terms are unacceptable, return at once unopened.**

In case of emergency endangering health or the environment involving this product, call 1-800-992-5994. If you wish to obtain additional product information, visit our web site at [www.dowagro.com](http://www.dowagro.com).

Agricultural Chemical: Do not ship or store with food, feeds, drugs or clothing.

EPA Reg. No. 62719-556

EPA Est. 34704-MS-1  
900-014750 / 00292630

©Trademark of Dow AgroSciences LLC  
Produced for  
Dow AgroSciences LLC  
9330 Zionsville Road  
Indianapolis, IN 46268

Net Contents **2.5 gal**

## Contents

## Page

Precautionary Statements .....	1
Environmental Hazards .....	2
Physical Or Chemical Hazards .....	2
Agricultural Use Requirements .....	2
Non-agricultural Use Requirements .....	2
Storage and Disposal .....	2
General Information	
(How this product Works) .....	3
Mixing .....	4
Mixing With Water .....	4
Tank Mixing Procedure .....	4
Ammonium Sulfate .....	4
Colorants or Dyes .....	4
Drift Control Additives .....	4
Application Equipment and Techniques .....	4
Aerial Equipment .....	5
Spray Drift Management .....	5
Ground Broadcast Equipment .....	6
Hand-Held and High-Volume Equipment .....	6
Mixing For Hand-Held Sprayers .....	6
Selective Equipment (Wipers, etc.) .....	6
Injection Systems .....	7
CDA Equipment .....	7
Injection and Frill Application (Woody Brush and Trees) .....	7
Cut Stump Application .....	7
Forestry Site Preparation .....	8
Mid-Rotation Conifer Release and Spot Treatments for Crop Tree Release and Timber Stand Improvement .....	9
Wildlife Habitat Management .....	9
Wildlife Food Plots .....	9
Parks, Recreational and Residential Areas .....	9
General Noncrop Areas and Industrial Sites .....	9
Utility Sites .....	9
Railroads .....	9
Bare Ground, Ballast and Shoulders, Crossings, and Spot Treatment .....	9
Brush Control .....	10
Roadsides .....	10
Guardrails and other Obstacles to Mowing .....	10
Spot Treatment .....	10
Tank mixtures .....	10
Release of Bermudagrass or Bahiagrass Dormant Applications .....	10
Actively Growing Bermudagrass .....	10
Actively Growing Bahiagrass .....	11
General Nonselective Weed Control, Trim-and-Edge and Bare Ground .....	11
Chemical Mowing .....	11
Dormant Turfgrass .....	11
Actively Growing Bermudagrass .....	12
Turfgrass Renovation, Seed, or Sod Production .....	12
Ornamentals, Plant Nurseries and Christmas Trees .....	12
Greenhouse / Shadehouse .....	12
Annual Weeds Rate Tables (Alphabetically by Species) .....	12
Perennial Weeds Rate Table (Alphabetically by Species) .....	16
Woody Brush and Trees Rate Table (Alphabetically by Species) .....	23
Terms and Conditions of Use .....	25
Warranty Disclaimer .....	25
Inherent Risks of Use .....	25
Limitation of Remedies .....	26

## Precautionary Statements

### Hazards to Humans and Domestic Animals

# CAUTION

**Causes Moderate Eye Irritation • Prolonged or Frequently Repeated Skin Contact May Cause Allergic Reactions in Some Individuals**

**Avoid contact with eyes or clothing.**

### Personal Protective Equipment (PPE)

Some of the materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for category A on an EPA chemical-resistance category selection chart.

#### Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Chemical-resistant gloves made of any waterproof material such as natural rubber
- Shoes plus socks

Follow manufacturer's instructions for cleaning/maintaining PPE (Personal Protective Equipment). If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

### Engineering Controls

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d) (4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

### User Safety Recommendations

Users should wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.

### First Aid

**If in eyes:** Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing. Call a poison control center or doctor for treatment advice.

**If on skin:** Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact 1-800-992-5994 for emergency medical treatment information.

**Domestic Animals:** This product is considered to be relatively nontoxic to dogs and other domestic animals; however, ingestion of this product or large amounts of freshly sprayed vegetation may result in temporary gastrointestinal irritation (vomiting, diarrhea, colic, etc.). If such symptoms are observed, provide the animal with plenty of fluids to prevent dehydration. Call a veterinarian if symptoms persist for more than 24 hours.

## Environmental Hazards

Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when cleaning equipment or disposing of equipment washwaters.

## Physical or Chemical Hazards

Spray solutions of this product should be mixed, stored and applied using only stainless steel, aluminum, fiberglass, plastic or plastic-lined steel containers.

**Do not mix, store or apply this product or spray solutions of this product in galvanized steel or unlined steel (except stainless steel) containers or spray tanks.** This product, or spray solutions of this product react with such containers and tanks to produce hydrogen gas that may form a highly combustible gas mixture. This gas mixture could flash or explode, causing serious personal injury, if ignited by open flame, spark, welder's torch, lighted cigarette or other ignition source.

## Directions for Use

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Read all Directions for Use carefully before applying.

**This is an end-use product. Dow AgroSciences does not intend and has not registered it for reformulation.**

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your state or tribe, consult the agency responsible for pesticide regulation.

## Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 4 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls
- Chemical-resistant gloves made of any waterproof material such as natural rubber
- Shoes plus socks

## Non-Agricultural Use Requirements

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries or greenhouses.

Keep people and pets off treated areas until spray solution has dried.

## Storage and Disposal

**Pesticide Storage:** Do not contaminate water, food, feed or seed by storage or disposal.

**Pesticide Disposal:** Wastes of this pesticide may cause eye and skin irritation and may be dangerous. Improper disposal of excess pesticide, spray mixtures, or rinsate is a violation of Federal law. If these wastes cannot be disposed of according to label use instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

**Nonrefillable containers 5 gallons or less:**

**Container Reuse:** Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available. Triple rinse or pressure rinse container (or equivalent) promptly after emptying. **Triple rinse** as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container  $\frac{1}{4}$  full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times.

**Pressure rinse** as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 psi for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

**Refillable containers larger than 5 gallons:**

**Container Reuse:** Refillable container. Refill this container with pesticide only. Do not reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from this container into application equipment or a mix tank. Fill the container about 10% full with water and, if possible, spray all sides while adding water. If practical, agitate vigorously or recirculate water with the pump for two minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times.

**Nonrefillable containers larger than 5 gallons:**

**Container Reuse:** Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available.

## Storage and Disposal (Cont.)

Triple rinse or pressure rinse container (or equivalent) promptly after emptying. **Triple rinse** as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. **Pressure rinse** as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 psi for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

## General Information

### (How this product works)

Accord® XRT II herbicide is a broad-spectrum systemic herbicide with no soil residual activity and is intended for control of annual and perennial weeds and woody plants in noncrop areas and industrial sites, such as utility, railroad and roadside rights-of-way, airports, natural and production (plantations) forests for site preparation, mid-rotation release treatments, and timber stand improvement activities, wildlife and habitat management areas, wildlife openings, natural areas, such as wildlands, campgrounds, parks and recreational areas, wildlife refuges, rangeland and in and around seasonally dry wetlands including ditchbanks, dry ditches and dry canals and grazed areas on listed noncrop sites.

It is permissible to treat non-irrigation ditch banks, seasonally dry wetlands, flood plains, deltas, marshes, swamps, bogs, and transitional areas between upland and lowland sites. Do not apply to open water such as lakes, reservoirs, rivers, streams, creeks, salt water bays, or estuaries.

This product is generally non-selective and gives broad-spectrum control of many annual weeds, perennial weeds, woody brush and trees. It is formulated as a water-soluble liquid. It may be applied through most standard industrial or field-type sprayers after dilution and thorough mixing with water or other carriers according to label instructions.

**Although not generally required, surfactant may be added to highly dilute spray solutions or when the application rate being used is at the low end of the effective rate range.**

Nonionic surfactants that are labeled for use with herbicides may be used. Do not reduce rates of this product when adding surfactant. When using additional surfactant, a surfactant concentration of 0.125 to 0.25 percent (1 to 2 pints per 100 gallons of spray solution) is specified for surfactants containing 70 percent or more active ingredient. Read and follow the precautionary

statements and applicable use directions on the label of the surfactant product.

**Time to Symptoms:** The active ingredient in this product moves through the plant from the point of foliage contact to and into the root system. Visible effects on most annual weeds occur within 2 to 4 days, but on most perennial weeds may not occur for 7 days or more. Extremely cool or cloudy weather following treatment may slow activity of this product and delay development of visual symptoms. Visible effects are a gradual wilting and yellowing of the plant that advances to complete browning of above ground growth and deterioration of underground plant parts.

**Stage of Weeds:** Annual weeds are easiest to control when they are small. Best control of most perennial weeds is obtained when treatment is made at late growth stages approaching maturity. Refer to the annual, perennial, woody brush and trees rate tables for directions for specific weeds.

Always use the higher rate of this product per acre within the specified range when weed growth is heavy or dense or weeds are growing in an undisturbed (noncultivated) area.

Do not treat weeds under poor growing conditions such as drought stress, disease or insect damage, as reduced weed control may result. Reduced herbicidal activity may also occur when treating weeds heavily covered with dust.

**Cultural Considerations:** Reduced control may result when applications are made to annual or perennial weeds that have been mowed, grazed, or cut, and have not been allowed to regrow to the specified stage for treatment.

**Rainfastness:** Heavy rainfall soon after application may wash this product off of the foliage and a repeat application may be required for adequate control.

**Spray Coverage:** For best results, spray coverage should be uniform and complete. Do not spray weed foliage to the point of runoff.

**Mode of Action:** The active ingredient in this product inhibits an enzyme found only in plants that is essential to formation of specific amino acids.

**No Soil Activity:** Weeds must be emerged at the time of application to be controlled by this product. Weeds germinating from seed after application will not be controlled. Unemerged plants arising from unattached underground rhizomes or rootstocks of perennials will not be affected by the herbicide and will continue to grow.

**Biological Degradation:** Degradation of this product is primarily a biological process carried out by soil microbes.

**Tank Mixing:** This product does not provide residual weed control. For subsequent residual weed control, follow a label-approved herbicide program. Read and carefully observe the cautionary statements and all other information appearing on the labels of all herbicides used. Use according to the most restrictive label directions for each product in the mixture.

Buyer and all users are responsible for all loss or damage in connection with the use or handling of mixtures of this product with herbicides or other materials that are not expressly directed in this labeling. Mixing this product with herbicides or other materials not specified on this label may result in reduced performance.

**Annual Maximum Use Rate:** For noncrop uses, the combined total of all treatments must not exceed 8 quarts of this product

per acre per year. The maximum use rates stated throughout this product's labeling apply to this product combined with the use of all other herbicides containing glyphosate or sulfosate as the active ingredient, whether applied as mixtures or separately. Calculate the application rates and ensure that the total use of this and other glyphosate or sulfosate containing products does not exceed stated use rate.

## Attention

**Avoid contact of herbicide with foliage, green stems, exposed non-woody roots or fruit of crops, desirable plants and trees, because severe injury or destruction may result.**

**AVOID DRIFT. Extreme care must be used when applying this product to prevent injury to desirable plants and crops.**

Do not allow the herbicide solution to mist, drip, drift or splash onto desirable vegetation since minute quantities of this product can cause severe damage or destruction to the crop, plants or other areas on which treatment was not intended. The likelihood of injury occurring from the use of this product increases when winds are gusty, as wind velocity increases, when wind direction is constantly changing or when there are other meteorological conditions that favor spray drift. When spraying, avoid combinations of pressure and nozzle type that will result in splatter or fine particles (mist) which are likely to drift. **Avoid applying at excessive speed or pressure.**

**NOTE:** Use of this product in any manner not consistent with this label may result in injury to persons, animals or crops, or other unintended consequences. Keep container closed to prevent spills and contamination.

## Mixing

Clean sprayer parts immediately after using this product by thoroughly flushing with water.

**NOTE:** Reduced results may occur if water containing soil is used, such as visibly muddy water or water from ponds and ditches that is not clear.

## Mixing with Water

This product mixes readily with water. Mix spray solutions of this product as follows: Fill the mixing or spray tank with the required amount of water. Add the specified amount of this product near the end of the filling process and mix well. Use caution to avoid siphoning back into the carrier source. Use approved anti-back-siphoning devices where required by state or local regulations. During mixing and application, foaming of the spray solution may occur. To prevent or minimize foam, avoid the use of mechanical agitators, terminate by-pass and return lines at the bottom of the tank and, if needed, use an approved anti-foam or defoaming agent.

## Tank Mixing Procedure

Mix labeled tank mixtures of this product with water as follows:

1. Place a 20 to 35 mesh screen or wetting basket over filling port.
2. Through the screen, fill the spray tank one-half full with water and start agitation.
3. If a wettable powder is used, make a slurry with the water carrier, and add it **slowly** through the screen into the tank. Continue agitation.

4. If a flowable formulation is used, premix one part flowable with one part water. Add diluted mixture **slowly** through the screen into the tank. Continue agitation.
5. If an emulsifiable concentrate formulation is used, premix one part emulsifiable concentrate with two parts water. Add diluted mixture slowly through the screen into the tank. Continue agitation.
6. Continue filling the spray tank with water and add the required amount of this product near the end of the filling process.
7. Add individual formulations to the spray tank as follows: wettable powder, flowable, drift control additive, water-soluble liquid and emulsifiable concentrate.

Maintain good agitation at all times until the contents of the tank are sprayed. If the spray mixture is allowed to settle, thorough agitation is required to resuspend the mixture before spraying is resumed.

Keep by-pass line on or near the bottom of the tank to minimize foaming. Screen size in nozzle or line strainers should be no finer than 50 mesh.

Always predetermine the compatibility of labeled tank mixtures of this product with water carrier by mixing small proportional quantities in advance.

Refer to the "Tank Mixing" section under "General Information" for additional precautions.

## Ammonium Sulfate

The addition of 1 to 2 percent dry ammonium sulfate by weight or 8.5 to 17 pounds per 100 gallons of water may increase the performance of this product, particularly when tank mixed with certain residual herbicides on annual and perennial weeds. The equivalent rate of ammonium sulfate in a liquid formulation may also be used. Ensure that ammonium sulfate is completely dissolved in the spray tank before adding herbicides. Thoroughly rinse the spray system with clean water after use to reduce corrosion.

**Note:** When using ammonium sulfate, apply this product at rates listed in this label. Lower rates will result in reduced performance.

## Colorants or Dyes

Agriculturally-approved colorants or marking dyes may be added to this product. Colorants or dyes used in spray solutions of this product may reduce performance, especially at lower rates or dilutions. Use colorants or dyes according to the manufacturer's directions.

## Drift Control Additives

Drift control additives may be used with all equipment types, except wiper applicators, sponge bars and CDA equipment. When a drift control additive is used, read and carefully observe the cautionary statements and all other information appearing on the additive label.

## Application Equipment and Techniques

Do not apply this product through any type of irrigation system.

This product may be applied with the following application equipment:

**Aerial:** Fixed Wing and Helicopter

**Ground Broadcast Spray:** Boom or boomless systems, pull-type sprayer, floaters, pick-up sprayers, spray coupes and other ground broadcast equipment.

**Hand-Held and High-Volume Spray Equipment:** Knapsack and backpack sprayers, pump-up pressure sprayers, handguns, hand wands, lances and other hand-held and motorized spray equipment used to direct the spray onto weed foliage.

**Selective Equipment:** Recirculating sprayers, shielded and hooded sprayers, wiper applicators and sponge bars.

**Injection Systems:** Aerial or ground injection sprayers.

**Controlled Droplet Applicator (CDA):** Hand-held or boom-mounted applicators, which produce a spray, consisting of a narrow range of droplet sizes.

**Apply these spray solutions in properly maintained and calibrated equipment capable of delivering desired volumes.**

**Injection and Frill Application (Woody Brush and Trees):** Use suitable equipment that will deliver this product into the living tissue of trees and brush.

**Cut Stump Application:** Apply using suitable equipment to ensure coverage of the entire cambium of cut stems.

## Aerial Equipment

**Do not apply this product using aerial spray equipment except under conditions as specified within this label.**

**For aerial application in California, refer to the federal supplemental label entitled "For Aerial Application in California Only" for aerial applications in that state for specific instructions, restrictions and requirements. In California, aerial application may be made for forestry site preparation and in noncrop areas. In California, this product is allowed for aerial application by helicopter only.**

Tank mixtures of this product plus Oust, dicamba or 2,4-D herbicide may not be applied by air in California.

Use the specified rates of this herbicide in 3 to 15 gallons of water per acre unless otherwise specified on this label. Unless otherwise specified, do not exceed 24 fluid ounces per acre. Refer to the individual use area sections of this label for volumes and application rates.

Avoid direct application to any body of water.

**AVOID DRIFT: Do not apply during low-level inversion conditions, when winds are gusty or under any other condition which favors drift. Drift may cause damage to any vegetation contacted to which treatment is not intended. To prevent injury to adjacent desirable vegetation, appropriate buffer zones must be maintained.**

Coarse sprays are less likely to drift; therefore, do not use nozzles or nozzle configurations that dispense spray as fine spray droplets. Do not angle nozzles forward into the airstream and do not increase spray volume by increasing nozzle pressure.

**Ensure uniform application:** To avoid streaked, uneven or overlapped application, use appropriate marking devices.

Thoroughly wash aircraft, especially landing gear, after each day of spraying to remove residues of this product accumulated

during spraying or from spills. **Prolonged exposure of this product to uncoated steel surfaces may result in corrosion and possible failure of the part. Landing gear are most susceptible.** The maintenance of an organic coating (paint), which meets aerospace specification MIL-C-38413, may prevent corrosion.

## Spray Drift Management

Avoiding spray drift at the application site is the responsibility of the applicator. The interaction of many equipment-and-weather-related factors determine the potential for spray drift. The applicator and the grower are responsible for considering all these factors when making decisions. The following drift management requirements must be followed to avoid off-target drift movement from aerial applications to agricultural field crops. These requirements do not apply to forestry applications, public health uses or to applications using dry formulations.

1. The distance of the outer most nozzles on the boom must not exceed  $\frac{3}{4}$  the length of the wingspan or rotor.
2. Nozzles must always point backward parallel with the air stream and never be pointed downwards more than 45 degrees.

Where states have more stringent regulations, they should be observed.

The applicator should be familiar with and take into account the information covered in the following **Aerial Drift Reduction Advisory Information:**

**Importance of Droplet Size:** The most effective way to reduce drift potential is to apply large droplets. The best drift management strategy is to apply the largest droplets that provide sufficient coverage and control. Applying larger droplets reduces drift potential, but will not prevent drift if applications are made improperly, or under unfavorable environmental conditions (see Wind, Temperature and Humidity, and Temperature Inversion section of this label).

**Controlling Droplet Size:**

**Volume** - Use high flow rate nozzles to apply the highest practical spray volume. Nozzles with higher rated flows produce larger droplets.

**Pressure** - Use the lower spray pressures for the nozzle. Higher pressure reduces droplet size and does not improve canopy penetration. When higher flow rates are needed, use higher flow rate nozzles instead of increasing pressure.

**Number of nozzles** - Use the minimum number of nozzles that provide uniform coverage.

**Nozzle Orientation** - Orienting nozzles so that the spray is released backwards, parallel to the airstream will produce larger droplets than other orientations. Significant deflection from the horizontal will reduce droplet size and increase drift potential.

**Nozzle Type** - Use a nozzle type that is designed for the intended application. With most nozzle types, narrower spray angles produce larger droplets. Consider using low-drift nozzles. Solid stream nozzles oriented straight back produce larger droplets than other nozzle types.

**Boom Length** - For some use patterns, reducing the effective boom length to less than  $\frac{3}{4}$  of the wingspan or rotor length may further reduce drift without reducing swath width.

**Application** - Applications should not be made at a height greater than 10 feet above the top of the largest plants unless a greater height is required for aircraft safety. Making applications at the lowest height that is safe reduces exposure of droplets to evaporation and wind.

**Swath Adjustment:** When applications are made with a crosswind, the swath will be displaced downwind. Therefore, on the up and downwind edges of the field, the applicator must compensate for this displacement by adjusting the path of the aircraft upwind. Swath adjustment distance should increase, with increasing drift potential (higher wind, smaller drops, etc.).

**Wind:** Drift potential is lowest between wind speeds of 2-10 mph. However, many factors, including droplet size and equipment type determine drift potential at any given speed. Application should be avoided below 2 mph due to variable wind direction and high inversion potential. Note: Local terrain can influence wind patterns. Every applicator should be familiar with local wind patterns and how they affect drift.

**Temperature and Humidity:** When making applications in low relative humidity, set up equipment to produce larger droplets to compensate for evaporation. Droplet evaporation is most severe when conditions are both hot and dry.

**Temperature Inversions:** Applications should not occur during a temperature inversion, because drift potential is high. Temperature inversions restrict vertical air mixing, which causes small suspended droplets to remain in a concentrated cloud. This cloud can move in unpredictable directions due to the light variable winds common during inversions. Temperature inversions are characterized by increasing temperatures with altitude and are common on nights with limited cloud cover and light to no wind. They begin to form as the sun sets and often continue into the morning. Their presence can be indicated by ground fog; however, if fog is not present, inversions can also be identified by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a connected cloud (under low wind conditions) indicates an inversion, while smoke that moves upwards and rapidly dissipates indicates good vertical air mixing.

**Sensitive Areas:** The pesticide should only be applied when the potential for drift to adjacent sensitive areas (e.g., residential areas, bodies of water, known habitat for threatened or endangered species, non-target crops) is minimal (e.g., when wind is blowing away from the sensitive areas).

## Ground Broadcast Equipment

Use the specified rates of this product in sufficient spray volume per acre as a broadcast spray to provide complete and uniform coverage. As density of weeds increases, spray volume should be increased to ensure complete coverage. Use proper nozzles to avoid spraying a fine mist. For best results with ground application equipment, use flat fan nozzles. Check for even distribution of spray droplets.

## Hand-Held and High-Volume Equipment

For applications made on a spray-to-wet basis, apply directly to foliage of vegetation to be controlled. Spray coverage should be uniform and complete. Do not spray to the point of runoff. Coarse sprays are required to reduce potential drift.

For control of weeds listed in the annual weeds rate tables, apply a 0.5 percent solution of this product to weeds less than 6 inches in height or runner length. Apply prior to seedhead formation in grass or bud formation in broadleaf weeds. For annual weeds over 6 inches tall, or unless otherwise specified, use a 1 percent solution.

Use a 1.5 percent solution on harder-to-control perennials, woody brush species and vines.

**For low volume directed spray applications,** use a 3.75 to 7.5 percent solution of this product for control or suppression of annual weeds, perennial weeds, or woody brush and trees. Spray coverage should be uniform with at least 50% of the foliage contacted. Coverage of the top one-half of the plant, including growing tip, is important for best results (over-the-top and top down coverage). To ensure adequate spray coverage, spray all sides of large or tall woody brush and trees, when foliage is thick and dense, or where there are dense or tall sprouts.

## Mixing for Hand-held Sprayers

Prepare the desired volume of spray solution by mixing the amount of this product in water as shown in the following table:

### Spray Solution

Spray Concentration (percent)	Amount of This Product for Desired Volume:		
	1 gal	25 gal	100 gal
0.5%	<sup>2</sup> / <sub>3</sub> fl oz	1 pt	2 qt
0.75%	1 fl oz	24 fl oz	3 qt
1.0%	1 <sup>1</sup> / <sub>3</sub> fl oz	1 qt	1 gal
1.5%	2 fl oz	1 <sup>1</sup> / <sub>2</sub> qt	1 <sup>1</sup> / <sub>2</sub> gal
2.0%	2 <sup>2</sup> / <sub>3</sub> fl oz	2 qt	2 gal
3.75%	5 fl oz	3 <sup>3</sup> / <sub>4</sub> qt	3 <sup>3</sup> / <sub>4</sub> gal
5.0%	6 <sup>1</sup> / <sub>2</sub> fl oz	5 qt	5 gal
10.0%	13 fl oz	10 qt	10 gal

2 tablespoons = 1 fluid ounce

For use in knapsack sprayers, it is suggested that the specified amount of this product be mixed with water in a larger container. Fill sprayer with the mixed solution.

## Selective Equipment

This product may be mixed as directed and applied through recirculating spray systems, shielded sprayers, hooded sprayers, wiper applicators or sponge bars to listed weeds growing in any noncrop site specified on this label.

A recirculating spray system directs the spray solution onto weeds growing above desirable vegetation, while spray solution not intercepted by weeds is collected and returned to the spray tank for reuse.

A shielded or hooded applicator directs the herbicide solution onto weeds, while shielding desirable vegetation from the herbicide.

A wiper or sponge applicator applies the herbicide solution onto weeds by rubbing the weed with an absorbent material containing the herbicide solution.

## Avoid contact of herbicide with desirable vegetation.

Contact of the herbicide solution with desirable vegetation may result in damage or destruction. Applicators used above desirable vegetation should be adjusted so that the lowest spray stream or wiper contact point is at least 2 inches above the desirable vegetation. Droplets, mist, foam or splatter of the herbicide solution settling on desirable vegetation may result in discoloration, stunting or destruction.

Applications made above desirable plants should be made when the weeds are a minimum of 6 inches above the desirable vegetation. Better results may be obtained when more of the weed is exposed to the herbicide solution. Weeds not contacted by the herbicide solution will not be affected. This may occur in dense clumps, severe infestations or when the height of the weeds varies so that not all weeds are contacted. In these instances, repeat treatment may be necessary.

### Shielded and hooded applicators

Use nozzles that provide uniform coverage within the treated area. Keep shields on these sprayers adjusted to protect desirable vegetation. **Extreme care must be exercised to avoid contact of herbicide with desirable vegetation.**

### Wiper applicators and sponge bars

Wiper applicators are devices that physically wipe appropriate amounts of this product directly onto the weed.

Equipment must be designed, maintained and operated to prevent the herbicide solution from contacting desirable vegetation. Operate this equipment at ground speeds no greater than 5 mph. Performance may be improved by reducing speed in areas of heavy weed infestations to ensure adequate wiper saturation. Better results may be obtained if 2 applications are made in opposite directions.

Avoid leakage or dripping onto desirable vegetation. Adjust height of applicator to ensure adequate contact with weeds. Keep wiping surfaces clean. Be aware that, on sloping ground, the herbicide solution may migrate, causing dripping on the lower end and drying of the wicks on the upper end of a wiper applicator.

Do not use wiper equipment when weeds are wet.

Mix only the amount of solution to be used during a 1-day period, as reduced activity may result from use of leftover solutions. Clean wiper parts immediately after using this product by thoroughly flushing with water.

**For Rope or Sponge Wick Applicators:** Mix 3 quarts of this product in 2 gallons of water to prepare a 25 percent solution. Apply this solution to weeds listed in this section.

**For Porous-Plastic Applicators:** Solutions ranging from 25 to 100 percent of this product in water may be used in porous-plastic wiper applicators.

## Injection Systems

This product may be used in aerial or ground injection spray systems. It may be used as a liquid concentrate or diluted prior to injecting into the spray stream. Do not mix this product with the concentrate of other products when using injection systems.

## CDA Equipment

The rate of this product applied per acre by vehicle-mounted controlled droplet application (CDA) equipment should not be less than the amount specified in this label for conventional broadcast applications. For vehicle-mounted and hand-held CDA equipment, use spray volumes and application techniques specified by the manufacturer.

**Note:** Controlled droplet application equipment produces a spray pattern that is not easily visible. Extreme care must be exercised to avoid spray or drift contacting the foliage or any other green tissue of desirable vegetation, as damage or destruction may result.

## Injection and Frill Application (Woody Brush and Trees)

**Types of Application:** Injection and frill application may be used in any noncrop site listed on this label

This product may be used to control woody brush and trees by injection or frill applications. Apply this product using suitable equipment that must penetrate into the living tissue. Apply the equivalent of 1 ml of this product per each 2 to 3 inches of trunk diameter at breast height (DBH). This is best achieved by applying a 40 to 100 percent concentration of this product either to a continuous frill around the tree or as cuts evenly spaced around the tree below all branches. As tree diameter increases in size, better results are achieved by applying diluted material to a continuous frill or more closely spaced cuttings. Avoid application techniques that allow runoff to occur from frilled or cut areas in species that exude sap freely. In species such as this, make the frill or cuts at an oblique angle to produce a cupping effect and use a 100 percent concentration of this product. For best results, applications should be made during periods of active growth and after full leaf expansion.

## Cut Stump Application

Woody vegetation may be controlled by treating freshly cut stumps of trees and resprouts with this product.

**Types of Application:** Treating cut stumps in any noncrop site listed on this label

**Specific Use:** This product will control regrowth of cut stumps and resprouts of many types of woody brush and tree species, some of which are listed below. Apply this product using suitable equipment to ensure coverage of the entire cambium. Cut trees or resprouts close to the soil surface. Apply a 40 to 100 percent solution of this product to the freshly cut surface **immediately** after cutting. Delays in application may result in reduced performance. For best results, applications should be made during periods of active growth and full leaf expansion.

**Precautions and Restrictions:** Do not make cut stump applications when the roots of desirable woody brush or trees may be grafted to the roots of the cut stump. Injury resulting from root grafting can occur in adjacent woody brush or trees of the same or closely related species. Different woody species growing in close proximity do not typically form root grafts.

## Forestry Site Preparation

This product herbicide is specified for the control or partial control of woody brush, trees and herbaceous weeds in forestry. This product is also specified for use in preparing or establishing wildlife openings within these sites and maintaining logging roads.

In forestry sites, this product is specified for use in site preparation prior to planting any tree species, including Christmas trees, eucalyptus, hybrid tree cultivars and silvicultural nursery sites. Unless otherwise specified, applications of this product may be made for control or partial control of herbaceous weeds, woody brush and trees listed in the "Weeds Controlled" section of the product label for this product.

### Application Rates:

Method of Application	Application Rate	Spray Volume (gal/acre)
<b>Broadcast</b>		
Aerial	1.5 to 8.0 qt/acre	5 to 30
Ground	1.5 to 8.0 qt/acre	10 to 60
<b>Spray-to-Wet</b>		
Handgun	0.75 to 1.5% by volume	spray-to-wet
Backpack		
<b>Low Volume Directed Spray ††</b>		
Handgun	3.75% to 7.5% by volume	partial coverage
Backpack		

†† For low volume directed spray applications, coverage should be uniform with at least 50 percent of the foliage contacted. For best results, coverage of the top one-half of the plant including the growing tip is important (over-the-top and down coverage). To ensure adequate spray coverage, spray all sides of large or tall woody brush and trees, when foliage is thick and dense, or where there are multiple stems or tall sprouts.

Use higher rates of this product within the specified rate ranges for control or partial control of woody brush, trees and hard-to-control perennial herbaceous weeds. For best results, apply to actively growing woody brush and trees after full leaf expansion and before fall color and leaf drop. Use increased rates within the specified rate range to control perennial herbaceous weeds from emergence up to the appearance of seedheads, flowers or berries. Use lower rates within the rate range to control annual herbaceous weeds and actively growing perennial herbaceous weeds after seedheads, flowers or berries appear. Apply to foliage of actively growing annual herbaceous weeds anytime after emergence.

This product has no herbicidal or residual activity in the soil. Where repeat applications are necessary, do not exceed 8 quarts per acre per year.

### Tank Mixtures

This product may be used in tank mix combination with other herbicide products to broaden the spectrum of vegetation controlled. When tank mixing, read and observe applicable use directions, precautions and limitations on the respective product labels. Use according to the most restrictive precautionary statements for each product in the mixture. Any specified rate of this product may be used in a tank mix.

**Note:** For forestry site preparation, make sure the tank mix product is approved for use prior to planting the desired species. Observe planting interval restrictions.

Any specified rate of this product may be used in a tank mix with the following products for forestry site preparation:

Product	Method of Application and Use Rates (Broadcast) ††
Garlon® 3A † herbicide	1 to 4 qt/acre
Garlon 4 herbicide	1 to 4 qt/acre
Arsenal Applicators Concentrate	2 to 16 fl oz/acre
Escort herbicide	1/2 to 1 1/2 oz/acre
Chopper herbicide	4 to 32 fl oz/acre
Oust herbicide	1 to 4 oz/acre
	<b>Spray-to-Wet Rates</b>
Arsenal Applicators Concentrate	1/32% to 1/2% by volume
	<b>Low Volume Directed Spray Rates</b>
Arsenal Applicators Concentrate	1/8% to 1/2% by volume

† Ensure that Garlon 3A is thoroughly mixed with water before adding this product. Agitation is required while mixing this product with Garlon 3A to avoid compatibility problems.

†† For a tank mix partner, up to the maximum labeled rate for a treatment site may be applied in combination with this product.

For control of herbaceous weeds, use the lower specified tank mixture rates. For control of dense stands or difficult-to-control woody brush and trees, use the higher specified rates.

### Aerial Equipment

For aerial application, apply this product in forestry sites by helicopter only. For details on aerial application, refer to "Aerial Equipment" in the "Application Equipment and Techniques" section of this label.

### Ground Broadcast Equipment

This product is specified for broadcast applications using suitable ground equipment in forestry sites. For details on ground broadcast application, refer to "Ground Broadcast Equipment" in the "Application Equipment and Techniques" section of this label. Apply the specified rates of this product as a broadcast spray in sufficient spray volume to provide complete and uniform coverage of plant foliage, unless otherwise specified. Check for even distribution throughout the spray pattern.

### Backpack and Handgun Equipment

This product is for application through backpack and handgun equipment. For details, refer to "Hand-Held and High Volume Equipment" in the "Application Equipment and Techniques" section of this label.

For spray-to-wet applications, coverage should be uniform and complete, but not to the point of runoff.

This product may be used for low volume directed sprays for spot treatment of trees and brush. It is most effective in areas where there is a low density of undesirable trees or brush. For

flat fan and cone nozzles, spray the foliage of the targeted vegetation. Small, open branched trees need only be treated from one side. If the foliage is thick or there are multiple root sprouts, application must be made from several sides to ensure adequate spray coverage.

---

## Mid-Rotation Conifer Release and Spot Treatments for Crop Tree Release and Timber Stand Improvement

---

This product is applied as a ground broadcast or directed spray application for mid-rotation release applications under the canopy of pines (and other conifers) and hardwoods. Applications must be made using application techniques that prevent or minimize direct contact to the foliage of crop trees (such as in stands of pine, other conifers or hardwood). This may be accomplished using directed sprays and ground equipment with nozzles oriented to target only undesirable understory vegetation below the crop tree canopy. This product is applied as a spot, individual plant treatment (see Hand-Held and High-Volume Equipment section of this label) for woody and herbaceous weeds. When making spot applications, do not allow spray to contact the foliage of desirable crop trees.

---

## Wildlife Habitat Management and Restoration

---

**Types of Uses:** Small, habitat restoration and maintenance, wildlife food plots

### Habitat restoration and maintenance

**Specific Use:** This product may be used to control exotic and other undesirable vegetation in habitat management and natural areas, including rangeland and wildlife refuges. Applications can be made to allow recovery of native plant species, prior to planting desirable native species, and for similar broad-spectrum vegetation control requirements. Spot treatments can be made to selectively remove unwanted plants for habitat maintenance and enhancement.

### Wildlife food plots

**Specific Use:** This product may be used as a site preparation treatment to control annual and perennial weeds prior to planting wildlife food plots. Any wildlife food species may be planted after applying this product, or native species may be allowed to repopulate the area. If tillage is needed to prepare a seedbed, wait 7 days after application before tillage.

---

## Parks, Recreational and Residential Areas

---

This product may be used in parks, recreational and residential areas. It may be applied with any application equipment described in this label. This product may be used to trim-and-edge around trees, fences, paths, around buildings, sidewalks, and other objects in these areas. This product may be used for spot treatment of unwanted vegetation. This product may be used to eliminate unwanted weeds growing in established shrub

beds or ornamental plantings. This product may be used prior to planting an area to ornamentals, flowers, turfgrass (sod or seed), or prior to laying asphalt or beginning construction projects.

All of the instructions in the "**General Noncrop Areas and Industrial Sites**" section apply to park and recreational areas.

---

## General Noncrop Areas and Industrial Sites

---

**Labeled Use Sites:** This product may be used in areas such as airports, apartment complexes, Christmas tree farms, ditch banks, dry ditches, dry canals, fencerows, golf courses, industrial sites, lumberyards, manufacturing sites, office complexes, ornamental nurseries, parks, parking areas, petroleum tank farms and pumping installations, railroads, recreational areas, residential areas, roadsides, sod or turf seed farms, schools, storage areas, utility substations, warehouse areas, other public areas, and similar industrial and noncrop sites and wildlife habitat management areas.

**Types of Applications:** General nonselective weed control, trim-and-edge, chemical mowing, cut stumps, injection and frill, habitat management.

This product may be used in general noncrop areas. It may be applied with any application equipment described in this label. This product may be used to trim-and-edge around objects in noncrop sites, for spot treatment of unwanted vegetation and to eliminate unwanted weeds growing in established shrub beds or ornamental plantings. This product may be used prior to planting an area to ornamentals, flowers, turfgrass (sod or seed), or prior to laying asphalt or beginning construction projects.

## Utility Sites

**Labeled Use Sites:** This product may be used in areas such as electrical power, pipeline, and telephone rights-of-way, and in other sites associated with these rights-of-way such as substations, roadsides, railroads or similar rights-of-way that run in conjunction with utilities and in general noncrop areas.

**Types of Applications:** General nonselective weed control, trim-and-edge, chemical mowing, cut stumps, injection and frill, habitat management.

This product may be applied with any application equipment described in this label. This product may be used to trim-and-edge around objects in these sites, for spot treatment of unwanted vegetation and to eliminate unwanted weeds, brush, vines and other vegetation growing in established shrub beds or ornamental plantings. This product may be used prior to planting an area or prior to laying asphalt or beginning construction projects.

## Railroads

All of the instructions in the "**General Noncrop Areas and Industrial Sites**" section apply to railroads.

### Bare ground, Ballast and Shoulders, Crossings, and Spot treatment

This product may be used to maintain bare ground on railroad ballast and shoulders. Repeat applications of this product may be used, as weeds emerge, to maintain bare ground. This product may be used to control tall-growing weeds to improve line-of-sight at railroad crossings and reduce the need for

mowing along rights-of-way. Apply the specified rate in sufficient spray volume to provide complete and uniform coverage of target vegetation, unless otherwise specified.

This product may be used in tank mix combination with other herbicide products labeled for use on non-cropland areas and industrial sites to broaden the spectrum of vegetation controlled. Follow applicable use directions, precautions and limitations on the respective product labels. Use according to the most restrictive precautionary statements for each product in the mixture.

### Brush control

This product may be used to control woody brush and trees on railroad rights-of-way. Apply 3 to 8 quarts of this product per acre as a broadcast spray, using boom-type or boomless nozzles. Apply the specified rate in sufficient spray volume to provide complete and uniform coverage of target vegetation, unless otherwise specified. Apply a 3/4 to 2 percent solution of this product when using high-volume spray-to-wet applications. Apply a 5 to 10 percent solution of this product when using low volume directed sprays for spot treatment. This product may be mixed with the following herbicide products for enhanced control of woody brush and trees:

Arsenal	Garlon 4
Escort	Tordon* K
Garlon 3A	

### Bermudagrass release

This product may be used to control or partially control many annual and perennial weeds for effective release of actively growing bermudagrass. Apply 0.75 to 2.25 pints of this product in up to 80 gallons of spray solution per acre. Use the lower rate when treating annual weeds below 6 inches in height (or runner length). Use the higher rate as weeds increase in size or as they approach flower or seedhead formation. These rates will also provide partial control of the following perennial species:

Bahiagrass	Johnsongrass
Bluestem, silver	Trumpetcreeper
Fescue, tall	Vaseygrass

This product may be tank-mixed with Oust. If tank-mixed, use no more than 0.75 to 2.25 pints of this product with 1 to 2 ounces of Oust per acre. Use the lower rates of each product to control annual weeds less than 6 inches in height (or runner length) that are listed in this label and the Oust label. Use the higher rates as annual weeds increase in size and approach the flower or seedhead stages. These rates will also provide partial control of the following perennial weeds:

Bahiagrass	Fescue, tall
Blackberry	Johnsongrass
Bluestem, silver	Poorjoe
Broomsedge	Raspberry
Dallisgrass	Trumpetcreeper
Dewberry	Vaseygrass
Dock, curly	Vervain, blue
Dogfennel	

Use only on well-established bermudagrass. Bermudagrass injury may result from the treatment, but regrowth will occur under moist conditions. Repeat applications in the same season are not specified, since severe injury may occur.

## Roadsides

All of the instructions in the "General Noncrop Areas and Industrial Sites" section apply to roadsides.

### Roadside Shoulder and Median treatments

This product may be used on road shoulders. It may be applied with boom sprayers, manifold nozzle systems, shielded boom sprayers, off-center nozzles and OC nozzle clusters, under-deck mowing + herbicide systems, hand-held equipment, and similar equipment.

### Guardrails and other obstacles to mowing

This product may be used to control weeds growing under guardrails and around signposts and other objects along the roadside.

### Spot treatment

This product may be used as a spot treatment to control unwanted vegetation growing along roadsides.

### Tank mixtures

This product may be used in tank mix combination with other herbicide products labeled for use on non-cropland areas and industrial sites to broaden the spectrum of vegetation controlled. Follow applicable use directions, precautions and limitations on the respective product labels. Use according to the most restrictive precautionary statements for each product in the mixture.

### Release of Bermudagrass or Bahiagrass Dormant applications

This product may be used to partially control many winter annual weeds and tall fescue for effective release of dormant bermudagrass or bahiagrass. Treat only when turf is dormant and prior to spring greenup. This product may also be tank-mixed with Oust for residual control. Tank mixtures of this product with Oust may delay greenup of bermudagrass. To avoid delay in greenup or severe damage to bahiagrass, use no more than 0.5 oz of Oust in tank mix combination with this product.

For best results on winter annuals, treat when plants are in an early growth stage (below 6 inches in height) after most have germinated. For best results on tall fescue, treat when fescue is at or beyond the 4- to 6-leaf stage.

Apply 6 to 48 fluid ounces of this product per acre alone or in a tank mixture with 1/4 to 1 ounce per acre of Oust. Use sufficient spray volume for uniform coverage of target vegetation. Use only in areas where bermudagrass or bahiagrass are desirable ground covers and where some temporary injury or discoloration can be tolerated. To avoid delays in greenup and minimize injury, add no more than 1 ounce of Oust per acre on bermudagrass and no more than 0.5 ounce of Oust per acre on bahiagrass and avoid treatments when these grasses are in a semi-dormant condition.

### Actively growing bermudagrass

This product may be used to control or partially control many annual and perennial weeds for effective release of actively growing bermudagrass. Apply up to 2.25 pints of this product in 10 to 40 gallons of spray solution per acre. Use the lower rate when treating annual weeds below 6 inches in height (or runner length). Use the higher rate as weeds increase in size or as they

approach flower or seedhead formation. These rates will also provide partial control of the following perennial species:

Bahiagrass	Johnsongrass
Bluestem, silver	Trumpetcreeper
Fescue, tall	Vaseygrass

This product may be tank-mixed with Oust. If tank-mixed, use up to 1.5 pints of this product with 1 to 2 ounces of Oust per acre. Use the lower rates of each product to control annual weeds less than 6 inches in height (or runner length) that are listed in this label and the Oust label. Use the higher rates as annual weeds increase in size and approach the flower or seedhead stages. These rates will also provide partial control of the following perennial weeds:

Bahiagrass	Fescue, tall
Bluestem, silver	Johnsongrass
Broomsedge	Poorjoe
Dallisgrass	Trumpetcreeper
Dock, curly	Vaseygrass
Dogfennel	Vervain, blue

Use only on well-established bermudagrass. Bermudagrass injury may result from the treatment, but regrowth will occur under moist conditions. Repeat applications of the tank mix in the same season are not specified, since severe injury may occur.

### Actively growing bahiagrass

For suppression of vegetable growth and seedhead inhibition of bahiagrass for approximately 45 days, apply 4.5 fluid ounces of this product in 10 to 40 gallons of water per acre. Apply 1 to 2 weeks after full greenup or after mowing to a uniform height of 3 to 4 inches. This application must be made prior to seedhead emergence.

For suppression up to 120 days, apply 3 fluid ounces of this product per acre, followed by an application of 1.5 to 3 fluid ounces per acre about 45 days later. Make no more than 2 applications per year.

A tank mixture of this product plus Oust may be used. Apply 6 fluid ounces of this product plus 0.25 ounces of Oust per acre 1 to 2 weeks following an initial spring mowing. Make only one application per year.

## General nonselective weed control, Trim-and-edge and Bare Ground

This product may be tank mixed with the following herbicide products. Refer to these product labels for labeled application sites and application rates. This product may be tank mixed with the products listed provided the product tank mixed is registered for use on this site. For annual weeds, use 1.5 pints per acre of this product when weeds are less than 6 inches tall and 2.25 pints per acre when weeds are greater than 6 inches tall. If weed growth is heavy or dense and/or growing in an undisturbed (non-cultivated) area and/or growing under stress, up to 3 quarts per acre may be applied. For perennial weeds, apply 1.5 to 3.75 quarts per acre in these tank mixes. For tank mixtures of this product with these products through backpack sprayers, handguns or other high-volume spray-to-wet applications, see the "Hand-Held and High Volume Equipment" section of this label for specified rates.

Arsenal	Plateau
Banvel (dicamba) †	Princep DF
Barricade 65WG	Princep Liquid
diuron †	Ronstar 50WP
Endurance	Sahara
Escort	simazine
Karmex DF	Surflan
Krovir I DF	Telar
Oust	Vanquish
Pendulum 3.3 EC	2,4-D †
Pendulum WDG	

† This product may be tank mixed with this product provided the label includes use on non-cropland and industrial sites.

Tank mixtures of this product with Oust, Banvel and 2,4-D may not be applied by air in California.

When applied as a tank mixture for bare ground, this product provides control of the emerged annual weeds and control or suppression of emerged perennial weeds, woody brush and trees.

For control or suppression of the following perennial weeds, apply 1.5 to 3 pints of this product plus 2 to 4 ounces of Oust per acre.

Bahiagrass	Fescue, tall
Bermudagrass	Johnsongrass
Broomsedge	Poorjoe
Dallisgrass	Quackgrass
Dock, curly	Vaseygrass
Dogfennel	Vervain, blue

## Chemical mowing

**Perennials:** This product will suppress perennial grasses listed in this section to serve as a substitute for mowing. Apply this product at a rate of 4.5 to 6 fluid ounces per acre. Use 6 fluid ounces of this product per acre when treating tall fescue, fine fescue, orchardgrass or quackgrass covers. Use 4.6 fluid ounces of this product per acre when treating Kentucky bluegrass. Apply treatments in 10 to 40 gallons of spray solution per acre.

**Precautions and Restrictions:** Use only in areas where some temporary injury or discoloration of perennial grasses can be tolerated.

**Annuals:** For growth suppression of some annual grasses, such as annual ryegrass, wild barley and wild oats growing in coarse turf on roadsides or other industrial areas, apply 3 to 3.75 fluid ounces of this product in 10 to 40 gallons of spray solution per acre. Applications should be made when annual grasses are actively growing and before the seedheads are in the boot stage of development. Treatments may cause injury to the desired grasses.

## Dormant turfgrass

This product may be used to control or suppress many winter annual weeds and tall fescue for effective release of dormant bermudagrass and bahiagrass turf. Treat only when turf is dormant and prior to spring greenup.

Apply 6 to 48 fluid ounces of this product per acre. Apply the specified rates in 10 to 40 gallons of water per acre. Use only in areas where bermudagrass or bahiagrass are desirable ground covers and where some temporary injury or discoloration can be tolerated.

Treatments in excess of 12 fluid ounces per acre may result in injury or delayed greenup in highly maintained areas, such as golf courses and lawns. **Do not** apply tank mixtures of this product plus Oust in highly maintained turfgrass areas. For further uses, refer to the “Roadsides” section of this label, which gives rates for dormant bermudagrass and bahiagrass treatments.

## Actively growing bermudagrass

This product may be used to control or partially control many annual and perennial weeds for effective release of actively growing bermudagrass. **Do not** apply more than 12 fluid ounces of this product per acre in highly maintained turfgrass areas. **Do not** apply tank mixtures of this product plus Oust in highly maintained turfgrass areas. For further uses, refer to the “Roadsides” section of this label, which gives rates for bermudagrass treatments. Use only in areas where some temporary injury or discoloration can be tolerated.

## Turfgrass Renovation, Seed, or Sod Production

This product controls most existing vegetation prior to renovating turfgrass areas or establishing turfgrass grown for seed or sod. For maximum control of existing vegetation, delay planting or sodding to determine if any regrowth from escaped underground plant parts occurs. When repeat treatments are necessary, sufficient regrowth must be attained prior to application. For warm-season grasses such as bermudagrass, summer or fall applications provide the best control. Where existing vegetation is growing under mowed turfgrass management, apply this product after omitting at least one regular mowing to allow sufficient growth for good interception of the spray.

Do not disturb soil or underground plant parts before treatment. Tillage or renovation techniques such as vertical mowing, coring or slicing should be delayed for 7 days after application to allow translocation into underground plant parts.

Desirable turfgrasses may be planted following the above procedures.

Hand-held equipment may be used for spot treatment of unwanted vegetation growing in existing turfgrass. Broadcast or hand-held equipment may be used to control sod remnants or other unwanted vegetation after sod is harvested.

Do not feed or graze turfgrass grown for seed or sod production for 8 weeks following application.

## Ornamentals, Plant Nurseries and Christmas Trees

**Post-direct, Trim-and-edge:** This product may be used as a post-directed spray around established woody ornamental species such as arborvitae, azalea, boxwood, crabapple, euonymus, fir, douglas fir, jobba, hollies, lilac, magnolia, maple, oak, privet, pine, spruce and yew. This product may also be used to trim and edge around trees, buildings, sidewalks and roads, potted plants and other objects in a nursery setting.

Desirable plants may be protected from the spray solution by using shields or coverings made of cardboard or other impermeable material. **This product is NOT specified for use as any over-the-top broadcast spray in ornamentals and Christmas trees.** Care must be exercised to avoid contact of spray, drift or mist with foliage or green bark of established ornamental species.

**Site preparation:** This product may be used prior to planting any ornamental, nursery or Christmas tree species.

**Greenhouse/Shadehouse:** This product may be used to control weeds growing in and around greenhouses and shadehouses. Desirable vegetation must not be present during application and air circulation fans must be turned off.

## Annual Weeds Rate Tables (Alphabetically By Species)

Use the specified rate for this product in sufficient broadcast spray volume per acre to ensure complete and uniform coverage of target vegetations. See “Application Equipment and Techniques” section for specified spray volume and coverage recommendations.

Apply to actively growing annual weeds.

Do not tank mix with soil residual herbicides when using these rates unless otherwise specified.

For weeds that have been mowed, grazed or cut, allow regrowth to occur prior to treatment.

For those rates less than 36 fluid ounces per acre, this product may be used up to 36 fluid ounces per acre where heavy weed densities exist. See following table for rate information for specific weeds.

Refer to this map for location of the regions listed in the annual weed tables below.



## Annual Weeds Rate Table, North and South Regions

		Rate of this product† (Fluid Ounces Per Acre)					
		9	12	18	24	30	36
Weed Species	Region	Maximum Height / Length					
annoda, spurred		-	1"	2"	3"	5"	8"
barley		-	18"	18"+	-	-	-
barnyardgrass	South	-	3"	5"	7"	9"	12"
	North	-	-	6"	12"	-	-
bittercress		-	12"	20"	-	-	-
bluegrass, annual		-	10"	-	-	-	-
bassia, fivehook		-	-	-	6"	-	-
brome, downy		6"	-	-	-	-	-
brome, Japanese		-	6"	-	24"	-	-
browntop panicum		-	6"	8"	12"	-	24"
burcucumber		-	6"	12"	-	-	-
buttercup		-	12"	20"	-	-	-
Carolina foxtail		-	20"	-	-	-	-
Carolina geranium		-	-	-	4"	-	9"
carpetweed		-	-	6"	12"	-	-
cheat		-	6"	20"	-	-	-
chervil		-	20"	-	-	-	-
chickweed		-	12"	18"	-	-	-
cocklebur		-	12"	18"	24"	-	-
copperleaf, hophornbeam		-	1"	2"	3"	4"	6"
copperleaf, Virginia		-	1"	2"	3"	4"	6"
corn		-	12"	20"	-	-	-
corn speedwell		-	12"	-	-	-	-
crabgrass		-	12"	18"	-	-	-
cutleaf evening primrose		-	-	-	3"	3"	6"
dwarf dandelion		-	20"	-	-	-	-
eastern mannagrass		-	8"	12"	-	-	-
eclipta		-	4"	8"	12"	-	-
fall panicum	South	-	4"	6"	8"	12"	24"
	North	-	6"	12"	18"	-	-
falsedandelion		-	20"	-	-	-	-
falseflax, smallseed		-	12"	-	-	-	-
fiddleneck		-	-	-	6"	6"	12"
field pennycress		-	6"	12"	-	-	-
filaree		-	-	-	-	-	12"
fleabane, annual		-	6"	20"	-	-	-
fleabane, hairy ( <i>conyza bonariensis</i> )		-	6"	-	-	-	-

## Annual Weeds Rate Table, North and South Regions (Cont.)

		Rate of this product† (Fluid Ounces Per Acre)					
		9	12	18	24	30	36
Weed Species	Region	Maximum Height / Length					
fleabane, rough		-	3"	6"	12"	-	-
Florida pusley		-	-	-	4"	4"	6"
foxtail	South	-	8"	12"	20"	-	-
	North	18"	18"+	-	-	-	-
goatgrass, jointed		-	6"	-	-	-	-
goosegrass		-	3"	5"	8"	-	18"
grain sorghum (milo)		-	6"	12"	20"	-	-
groundsel, common		-	6"	-	-	-	-
hemp sesbania		-	-	2"	4"	6"	8"
henbit		-	-	-	6"	-	20"
horseweed / marestalk ( <i>conyza canadensis</i> )	South	-	-	12"	30"	-	-
	North	-	6"	12"	18"	-	-
itchgrass		-	6"	12"	18"	-	-
jimsonweed		-	-	-	6"	6"	12"
johnsongrass (seedling)	South	-	-	-	18"	-	-
	North	-	12"	18"	-	-	-
junglerice		-	3"	5"	7"	9"	12"
knotweed		-	3"	8"	12"	-	20"
kochia <sup>1</sup>		-	3"-6"	12"	-	-	-
lambquarters		-	6"	8"	12"	-	20"
little barley		-	20"	-	-	-	-
London rocket		-	6"	-	-	-	-
mayweed		-	-	2"	6"	12"	18"
morningglory ( <i>ipomoea spp.</i> )		-	-	2"	4"	-	6"
Mustard, blue		6"	-	-	-	-	-
Mustard, tansy		6"	12"	20"	-	-	-
Mustard, tumble		6"	-	-	-	-	-
Mustard, wild		6"	12"	18"	-	-	-
nightshade, black		6"	12"	-	-	-	-
nightshade, hairy		-	6"	12"	-	-	-
Oats		-	-	6"	20"	-	-
Pigweed		-	12"	18"	24"	-	-
prickly lettuce		-	6"	12"	20"	-	-
purslane		-	-	-	6"	6"	12"
ragweed, common	South	-	4"	6"	8"	-	11"
	North	-	6"	12"	18"	-	-
ragweed, giant		-	-	4"	6"	-	11"

## Annual Weeds Rate Table, North and South Regions (Cont.)

Weed Species	Region	Rate of this product† (Fluid Ounces Per Acre)					
		9	12	18	24	30	36
		Maximum Height / Length					
red rice		-	-	-	4"	-	-
Russian thistle		-	6"	-	-	-	-
Rye	South	-	6"	20"	60"	-	-
	North	-	18"	18"+	-	-	-
ryegrass		-	-	-	6"	-	7+"
Sandbur, field		12"	-	-	-	-	-
shattercane		-	12"	18"	-	-	-
shepherd's-purse		-	6"	12"	-	-	-
sicklepod		-	-	2"	4"	-	8"
signalgrass, broadleaf		-	3"	5"	7"	9"	12"
smartweed, ladythumb		-	4"	6"	8"	-	12"
smartweed, Pennsylvania		-	4"	6"	8"	-	12"
sowthistle, annual		-	-	-	6"	-	12"
spanishneedles		-	-	-	8"	-	18"
speedwell, purslane		-	12"	-	-	-	-
sprangletop		-	6"	12"	20"	-	-
spurge, prostrate		-	6"	12"	20"	-	-
spurge, spotted		-	6"	12"	20"	-	-
spurry, umbrella		6"	-	-	-	-	-
stinkgrass		12"	-	-	-	-	-
sunflower		-	12"	18"	-	-	-
teaweed / prickly sida		1"	2"	3"	4"	6"	-
Texas panicum		6"	8"	12"	-	24"	-
velvetleaf	South	-	2"	3"	4"	5"	8"
	North	-	3"	6"	12"	-	-
Virginia pepperweed		-	18"	-	-	-	-
waterhemp		-	-	6"	12"	-	-
Wheat	South	-	6"	30"	-	-	-
	North	-	18"	18"+	-	-	-
wheat (over-wintered)		-	6"	18"	-	-	-
wild oats		-	12"	-	-	-	-
wild proso millet		-	-	6"	12"	12"	18"
witchgrass		-	12"	-	-	-	-
woolly cupgrass		-	6"	12"	-	-	-
yellow rocket		-	-	12"	20"	-	-

† Do not treat kochia in the button stage.

† If weed growth is heavy or dense and / or growing in an undisturbed (non-cultivated) area and / or growing under stress, up to 3 quarts per acre may be applied.

## Annual Weeds Rate Table, West Region

	Rate of this product † (Fluid Ounces Per Acre)				
	9	12	18	24	36
<b>Weed Species</b>	<b>Maximum Height/Length</b>				
Barley	12"	-	-	-	-
barnyardgrass	6"	-	-	-	-
bluegrass, annual	6"	-	-	-	-
bluegrass, bulbous	-	6"	-	-	-
brome, downy <sup>1</sup>	6"	-	-	-	-
Buttercup	-	12"	-	-	-
Cheat	-	6"	-	-	-
Chickweed	-	6"	-	-	-
Cocklebur	-	12"	-	-	-
Corn	-	6"	-	-	-
Crabgrass	-	12"	-	-	-
Dwarfandelion	-	12"	-	-	-
fall panicum	-	12"	-	-	-
falseflax, smallseed	-	12"	-	-	-
field pennycress	-	6"	-	-	-
Filaree	-	-	-	-	12
fleabane, hairy ( <i>conyza bonariensis</i> )	-	6"	-	-	-
Florida pusley	-	-	-	12"	-
Foxtail	(8 fl. oz. for up to 12")				
goatgrass, jointed	-	6"	-	-	-
groundsel, common	-	6"	-	-	-
Henbit	-	6"	-	-	-
horseweed/marestail ( <i>conyza canadensis</i> )	-	6"	-	-	-
johnsongrass, seedling	-	12"	-	-	-
Lambsquarters	-	6"	-	-	-
London rocket	-	6"	-	-	-
morningglory ( <i>ipomoea spp.</i> )	-	2"	-	-	-
mustard, blue	6"	-	-	-	-
mustard, tansy	6"	-	-	-	-
mustard, tumble	6"	-	-	-	-

	Rate of this product † (Fluid Ounces Per Acre)				
	9	12	18	24	36
<b>Weed Species (Cont.)</b>	<b>Maximum Height/Length</b>				
mustard, wild	6"	-	-	-	-
Pigweed	-	12"	-	-	-
Rye	12"	-	-	-	-
ryegrass, Italian	-	6"	-	-	-
sandbur, field	12"	-	-	-	-
Shattercane	12"	-	-	-	-
shepherd's-purse	-	6"	-	-	-
sowthistle, annual	-	6"	-	-	-
spurge, annual	-	6"	-	-	-
Stinkgrass	12"	-	-	-	-
Texas panicum	-	12"	-	-	-
Wheat	18"	-	-	-	-
wild oats	-	12"	-	-	-
Witchgrass	-	12"	-	-	-

<sup>1</sup> For control of downy brome in no-till systems, use 16 fluid ounces per acre.

† If weed growth is heavy or dense and/or growing in an undisturbed (non-cultivated) area and/or growing under stress, up to 3 quarts per acre may be applied.

## Perennial Weeds Rate Table (Alphabetically By Species)

Apply to actively growing perennial weeds.

**NOTE:** If weeds have been mowed or tilled, do not treat until plants have resumed active growth and have reached the specified stages.

Repeat treatments may be necessary to control weeds regenerating from underground parts or seed. Repeat treatments must be made prior to crop emergence.

Unless otherwise stated, allow 7 or more days after application before tillage.

Best results are obtained when soil moisture is adequate for active weed growth.

For difficult to control perennial weeds and woody brush and trees, where plants are growing under stressed conditions, or where infestations are dense, this product may be used at 3.75 to 7.5 quarts per acre for enhanced results. The annual maximum use rate for this product is 8 qt per acre per year.

Weed Species	Rate (pt/acre)	Water Volume (gpa)	Hand-Held (% Solution)
<b>Alfalfa</b>	<b>1.5 - 3</b>	<b>3 - 10</b>	<b>1.5%</b>
Make applications after the last hay cutting in the fall. Allow alfalfa to regrow to a height of 6 to 8 inches or more prior to treatment. Applications should be followed with deep tillage at least 7 days after treatment, but before soil freeze-up.			
<b>Alligatorweed</b>	<b>6</b>	<b>3 -20</b>	<b>1.25%</b>
Suppression. Apply when most of the plants are in bloom. Repeat applications will be required to maintain control.			
<b>Anise (fennel)</b>	<b>—</b>	<b>—</b>	<b>0.75 - 1.5%</b>
Apply as a spray-to-wet treatment. Optimum results are obtained when plants are treated at the bud to full-bloom stage of growth.			
<b>Bahiagrass</b>	<b>4.5 - 7.5</b>	<b>3 - 20</b>	<b>1.5%</b>
Apply when most plants have reached the early head stage.			
<b>Bentgrass</b>	<b>2.25</b>	<b>10 - 20</b>	<b>1.5%</b>
For suppression in grass seed production areas. For ground applications only. Ensure entire crown area has resumed growth prior to a fall application. Bentgrass should have at least 3 inches of growth. Tillage prior to treatment should be avoided. Tillage 7 to 10 days after application provides best results.			
<b>Bermudagrass</b>	<b>4.5 - 7.5</b>	<b>3 - 20</b>	<b>1.5%</b>
For control, apply 7.5 pints of this product per acre. For suppression, apply 4.5 pints per acre. Treat when bermudagrass is actively growing and seedheads are present. Retreatment may be necessary to maintain control.			
<b>Bermudagrass, water (knotgrass)</b>	<b>1.5 - 2.25</b>	<b>5 - 10</b>	<b>1.5%</b>
Apply 2.25 pints of this product in 5 to 10 gallons of water per acre. Apply when water bermudagrass is 12 to 18 inches in length. Allow 7 or more days before tilling, flushing or flooding the field.			
<b>Fall applications only:</b> Apply 1.5 pints of this product in 5 to 10 gallons of water per acre. Fallow fields should be tilled prior to application. Apply prior to frost on water bermudagrass that is 12 to 18 inches in length.			
<b>This product is not registered in California for use on water bermudagrass.</b>			
<b>Bindweed, field</b>	<b>0.75 - 7.5</b>	<b>3 - 20</b>	<b>1.5%</b>
Do not treat when weeds are under drought stress as good soil moisture is necessary for active growth.			
For control, apply 6 to 7.5 pints of this product per acre west of the Mississippi River and 4.5 to 6 pints east of the Mississippi River. Apply when the weeds are at or beyond full bloom. For best results, apply in late summer or fall. Fall treatments must be applied before a killing frost.			
Also for control, apply 3 pints of this product plus 0.5 pound a.i. of dicamba in 10 to 20 gallons of water per acre. Do not apply by air.			
For suppression on irrigated agricultural land, apply 1.5 to 3 pints of this product plus 1 pound a.i. of 2,4-D in 10 to 20 gallons of water per acre with ground equipment only. Applications should be made following harvest or in fall fallow ground when the bindweed is actively growing and the majority of runners are 12 inches or more in length. The use of at least one irrigation will promote active bindweed growth.			
For suppression, apply 12 fluid ounces of this product plus 0.5 pound a.i. of 2,4-D or 0.25 pound a.i. of dicamba in 3 to 10 gallons of water per acre for ground applications and 3 to 5 gallons of water per acre for aerial applications. Apply by air in fallow and reduced tillage systems only. Applications should be delayed until maximum emergence has occurred and when vines are between 6 to 18 inches in length.			
<b>In California only,</b> apply 1.5 to 7.5 pints of this product per acre. The actual rate needed for suppression or control will vary within this range depending on local conditions. For suppression on irrigated land where annual tillage is performed, apply 1.5 pints of this product in 3 to 10 gallons of water per acre. Apply to bindweed that has reached a length of 12 inches or greater. Allow maximum weed emergence and runner growth. Allow 3 or more days after application before tillage.			
<b>Bluegrass, Kentucky</b>	<b>1.5 - 3</b>	<b>3 - 40</b>	<b>1.5%</b>
Apply 3 pints of this product in 10 to 40 gallons of water per acre when most plants have reached boot-to-early seedhead stage of development. For suppression, in pasture or hay crop renovation, apply 1.5 to 2.25 pints of this product in 3 to 10 gallons of water per acre. Apply to actively growing plants when most have reached 4 to 12 inches in height.			

<b>Weed Species (Cont.)</b>	<b>Rate (pt/acre)</b>	<b>Water Volume (gpa)</b>	<b>Hand-Held (% Solution)</b>
<b>Blueweed, Texas</b>	<b>4.5 - 7.5</b>	<b>3 - 40</b>	<b>1.5%</b>
Apply 6 to 7.5 pints of this product per acre west of the Mississippi River and 4.5 to 6 pints per acre east of the Mississippi River. Apply when plants are at or beyond full bloom. New leaf development indicates active growth. For best results, apply in late summer or fall. Fall treatments must be applied before a killing frost.			
<b>Brackenfern</b>	<b>4.5 - 6</b>	<b>3 - 40</b>	<b>0.75 - 1.5%</b>
Apply to fully expanded fronds, which are at least 18 inches long.			
<b>Bromegrass, smooth</b>	<b>1.5 - 3</b>	<b>3 - 40</b>	<b>1.5%</b>
Apply 3 pints of this product in 10 to 40 gallons of water per acre when most plants have reached boot-to-early seedhead stage of development. For suppression, in pasture or hay crop renovation, apply 1.5 to 2.25 pints of this product in 3 to 10 gallons of water per acre. Apply to actively growing plants when most have reached 4 to 12 inches in height.			
<b>Bursage, woolly-leaved</b>	<b>—</b>	<b>3 - 20</b>	<b>1.5%</b>
For control, apply 3 pints of this product plus 0.5 lb a.i. of dicamba per acre. For suppression, apply 1.5 pints of this product plus 0.5 lb a.i. of dicamba per acre. Apply when plants are producing new active growth, which has been initiated by moisture for at least 2 weeks and when plants are at or beyond flowering.			
<b>Canarygrass, reed</b>	<b>3 - 4.5</b>	<b>3 - 40</b>	<b>1.5%</b>
For best results, apply when most plants have reached the boot-to-head stage of growth.			
<b>Cattail</b>	<b>4.5 - 7.5</b>	<b>3 - 40</b>	<b>1.5%</b>
Apply when most plants have reached the early head stage.			
<b>Clover; red, white</b>	<b>4.5 - 7.5</b>	<b>3 - 20</b>	<b>1.5%</b>
Apply when most plants have reached the early bud stage.			
<b>Cogongrass</b>	<b>4.5 - 7.5</b>	<b>10 - 40</b>	<b>1.5%</b>
Apply when cogongrass is at least 18 inches tall in late summer or fall. Due to uneven stages of growth and the dense nature of vegetation preventing good spray coverage, repeat treatments may be necessary to maintain control.			
<b>Dallisgrass</b>	<b>4.5 - 7.5</b>	<b>2 - 20</b>	<b>1.5%</b>
Apply when most plants have reached the early head stage.			
<b>Dandelion</b>	<b>4.5 - 7.5</b>	<b>3 - 40</b>	<b>1.5%</b>
Apply when most plants have reached the early bud stage of growth.			
Also for control, apply 12 fluid ounces of this product plus 0.5 pound a.i. 2,4-D in 3 to 10 gallons of water per acre.			
<b>Dock, curly</b>	<b>4.5 - 7.5</b>	<b>3 - 40</b>	<b>1.5%</b>
Apply when most plants have reached the early bud stage of growth.			
Also for control, apply 12 fluid ounces of this product plus 0.5 pound a.i. 2,4-D in 3 to 10 gallons of water per acre.			
<b>Dogbane, hemp</b>	<b>6</b>	<b>3 - 40</b>	<b>1.5%</b>
Apply when most plants have reached the late bud to flower stage of growth. Following crop harvest or mowing, allow weeds to regrow to a mature stage prior to treatment. For best results, apply in late summer or fall.			
For suppression, apply 12 fluid ounces of this product plus 0.5 pound a.i. of 2,4-D in 3 to 10 gallons of water per acre for ground applications and 3 to 5 gallons of water per acre for aerial applications. Delay applications until maximum emergence of dogbane has occurred.			
<b>Fescue (Except tall)</b>	<b>4.5 - 7.5</b>	<b>3 - 20</b>	<b>1.5%</b>
Apply when most plants have reached the early head stage.			
<b>Fescue, tall</b>	<b>1.5 - 4.5</b>	<b>3 - 40</b>	<b>1.5%</b>
Apply 4.5 pints of this product per acre when most plants have reached boot-to-early seedhead stage of development.			
Fall applications only: Apply 1.5 pints of this product in 3 to 10 gallons of water per acre. Apply to fescue in the fall when plants have 6 to 12 inches of new growth. A sequential application of 12 fluid ounces per acre of this product will improve long-term control and control seedlings germinating after fall treatments or the following spring.			

<b>Weed Species (Cont.)</b>	<b>Rate (pt/acre)</b>	<b>Water Volume (gpa)</b>	<b>Hand-Held (% Solution)</b>
<b>Guineagrass</b>	<b>4.5</b>	<b>3 - 40</b>	<b>0.75%</b>
Apply when most plants have reached at least the 7-leaf stage of growth. Ensure thorough coverage when using hand-held equipment.			
<b>Horsenettle</b>	<b>4.5 - 7.5</b>	<b>3 - 20</b>	<b>1.5%</b>
Apply when most plants have reached the early bud stage.			
<b>Horseradish</b>	<b>6</b>	<b>3 - 40</b>	<b>1.5%</b>
Apply when most plants have reached the late bud to flower stage of growth. For best results, apply in late summer or fall.			
<b>Iceplant</b>	<b>—</b>	<b>—</b>	<b>1.5%</b>
Iceplant should be at or beyond the early bud stage of growth. Thorough coverage is necessary for best control.			
<b>Jerusalem artichoke</b>	<b>4.5 - 7.5</b>	<b>3 - 20</b>	<b>1.5%</b>
Apply when most plants are in the early bud stage.			
<b>Johnsongrass</b>	<b>0.75 - 4.5</b>	<b>3 - 40</b>	<b>0.75%</b>
In annual cropping systems apply 1.5 to 3 pints of this product per acre. Apply 1.5 pints of this product in 3 to 10 gallons of water per acre. Use 3 pints of this product when applying 10 to 40 gallons of water per acre. In noncrop or areas where annual tillage (no-till) is not practiced, apply 3 to 4.5 pints of this product in 10 to 40 gallons of water per acre.			
For best results, apply when most plants have reached the boot-to-head stage of growth or in the fall prior to frost. Allow 7 or more days after application before tillage. Do not tank mix with residual herbicides when using the 1.5 pint per acre rate.			
For burndown of Johnsongrass, apply 12 fluid ounces of this product in 3 to 10 gallons of water per acre before the plants reach a height of 12 inches. For this use, allow at least 3 days after treatment before tillage.			
Spot treatment (suppression): Apply a 0.75% solution of this product when Johnsongrass is 12 to 18 inches in height. Coverage should be uniform and complete.			
<b>Kikuyugrass</b>	<b>3 - 4.5</b>	<b>3-40</b>	<b>1.5%</b>
Spray when most kikuyugrass is at least 8 inches in height (3 or 4-leaf stage of growth). Allow 3 or more days after application before tillage.			
<b>Knapweed</b>	<b>6</b>	<b>3-40</b>	<b>1.5%</b>
Apply when most plants have reached the late bud to flower stage of growth. For best results, apply in late summer or fall.			
<b>Lantana</b>	<b>-</b>	<b>-</b>	<b>0.75 - 1%</b>
Apply at or beyond the bloom stage of growth. Use the higher application rate for plants that have reached the woody stage of growth.			
<b>Lespedeza</b>	<b>4.5 - 7.5</b>	<b>3 - 20</b>	<b>1.5%</b>
Apply when most plants have reached the early bud stage.			
<b>Milkweed, common</b>	<b>4.5</b>	<b>3 - 40</b>	<b>1.5%</b>
Apply when most plants have reached the late bud to flower stage of growth.			
<b>Muhly, wirestem</b>	<b>1.5 - 3</b>	<b>3 - 40</b>	<b>1.5%</b>
Use 1.5 pints of this product in 3 to 10 gallons of water per acre. Use 3 pints of this product when applying 10 to 40 gallons of water per acre or in pasture, sod, or noncrop areas. Spray when the wirestem muhly is 8 inches or more in height. Do not till between harvest and fall applications or in the fall or spring prior to spring applications. Allow 3 or more days after application before tillage.			
<b>Mullein, common</b>	<b>4.5 - 7.5</b>	<b>3 - 20</b>	<b>1.5%</b>
Apply when most plants are in the early bud stage.			
<b>Napiergrass</b>	<b>4.5 - 7.5</b>	<b>3 - 20</b>	<b>1.5%</b>
Apply when most plants are in the early head stage.			
<b>Nightshade, silverleaf</b>	<b>3</b>	<b>3 - 10</b>	<b>1.5%</b>
Applications should be made when at least 60 percent of the plants have berries. Fall treatments must be applied before a killing frost.			

<b>Weed Species (Cont.)</b>	<b>Rate (pt/acre)</b>	<b>Water Volume (gpa)</b>	<b>Hand-Held (% Solution)</b>
<b>Nutsedge; purple, yellow</b>	<b>0.75 - 4.5</b>	<b>3 - 40</b>	<b>0.75 - 1.5%</b>
Apply 4.5 pints of this product per acre or apply a 0.75 to 1.5% solution for control of nutsedge plants and immature nutlets attached to treated plants. Treat when plants are in flower or when new nutlets can be found at rhizome tips. Nutlets, which have not germinated, will not be controlled and may germinate following treatment. Repeat treatments will be required for long-term control of ungerminated tubers.			
Sequential applications: 1.5 to 3 pints of this product in 3 to 10 gallons of water per acre will also provide control. Make applications when a majority of the plants are in the 3 to 5-leaf stage (less than 6 inches tall). Repeat this application, as necessary, when newly emerging plants reach the 3 to 5-leaf stage. Subsequent applications will be necessary for long-term control.			
For suppression of existing plants, apply 12 fluid ounces to 3 pints of this product in 3 to 40 gallons of water per acre. Treat when plants have 3 to 5 leaves and most are less than 6 inches tall. Repeat treatments will be required to control subsequent emerging plants or regrowth of existing plants.			
<b>Orchardgrass</b>	<b>1.5 - 3</b>	<b>3 - 40</b>	<b>1.5%</b>
Apply 3 pints of this product in 10 to 40 gallons of water per acre when most plants have reached boot-to-early seedhead stage of development. For suppression in pasture or hay crop renovation, apply 1.5 to 2.25 pints of this product in 3 to 10 gallons of water per acre. Apply to actively growing plants when most have reached 4 to 12 inches in height.			
<b>Orchardgrass sods going to no-till corn:</b> Apply 1.5 to 2.25 pints of this product in 3 to 10 gallons of water per acre. Apply to orchardgrass that is a minimum of 12 inches tall for spring applications and 6 inches tall for fall applications. Allow at least 3 days following application before planting. A sequential application of atrazine will be necessary for optimum results.			
<b>Pampasgrass</b>	—	—	<b>1.5%</b>
Pampasgrass should be at or beyond the boot stage of growth. Thorough coverage is necessary for best control.			
<b>Paragrass</b>	<b>4.5 - 7.5</b>	<b>3 - 20</b>	<b>1.5%</b>
Apply when most plants are in the early head stage.			
<b>Phragmites</b>	<b>4.5 - 7.5</b>	<b>10 - 40</b>	<b>0.75 - 1.5%</b>
For suppression. For best results, treat during late summer or fall months or when plants are actively growing and in full bloom. Treatment before or after this stage may lead to reduced control. Due to the dense nature of the vegetation, which may prevent good spray coverage or uneven stages of growth, repeat treatments may be necessary to maintain control. Visual control symptoms will be slow to develop.			
<b>Poison hemlock</b>	—	—	<b>0.75 - 1.5%</b>
Apply as a spray-to-wet treatment. Optimum results are obtained when plants are treated at the bud to full-bloom stage of growth.			
<b>Pokeweed, common</b>	<b>1.5</b>	<b>3 - 40</b>	<b>1.5%</b>
Apply to actively growing plants up to 24 inches tall.			
<b>Quackgrass</b>	<b>1.5 - 4.5</b>	<b>3 - 40</b>	<b>1.5%</b>
In annual cropping systems or in pastures and sods followed by deep tillage: Apply 1.5 pints of this product in 3 to 10 gallons of water per acre. For 10 to 40 gallons of water per acre, apply 3 pints of this product. Do not tank mix with residual herbicides when using the 1.5 pint rate. Spray when quackgrass is 6 to 8 inches in height. Do not till between harvest and fall applications or in fall or spring prior to spring application. Allow 3 or more days after application before tillage. In pastures or sods, use a moldboard plow for best results.			
In pastures, sods or noncrop areas where deep tillage does not follow application: Apply 3 to 4.5 pints of this product in 10 to 40 gallons of water per acre when the quackgrass is greater than 8 inches tall.			
<b>Redvine</b>	<b>1.25 - 3</b>	<b>5 - 10</b>	<b>1.5%</b>
For suppression, apply 18 fluid ounces of this product per acre at each of two applications 7 to 14 days apart or a single application of 3 pints per acre. Apply specified rates in 5 to 10 gallons of water per acre. Apply in late September or early October to plants that are at least 18 inches tall and have been growing 45 to 60 days since the last tillage operation. Make applications at least 1 week before a killing frost.			
<b>Reed, giant</b>	—	—	<b>1.5%</b>
Best results are obtained when applications are made in late summer to fall.			

<b>Weed Species (Cont.)</b>	<b>Rate (pt/acre)</b>	<b>Water Volume (gpa)</b>	<b>Hand-Held (% Solution)</b>
<b>Ryegrass, perennial</b>	<b>1.5 - 4.5</b>	<b>3 - 40</b>	<b>0.75%</b>
In annual cropping systems apply 1.5 to 3 pints of this product per acre. Apply 1.5 pints of this product in 3 to 10 gallons of water per acre. Use 3 pints of this product when applying 10 to 40 gallons of water per acre. In noncrop or areas where annual tillage (no-till) is not practiced, apply 3 to 4.5 pints of this product in 10 to 40 gallons of water per acre.			
For best results, apply when most plants have reached the boot-to-head stage of growth or in the fall prior to frost. Do not tank-mix with residual herbicides when using the 1.5 pint per acre rate.			
<b>Smartweed, swamp</b>	<b>4.5 - 7.5</b>	<b>3 - 40</b>	<b>1.5%</b>
Apply when most plants have reached the early bud stage of growth.			
Also for control, apply 12 fluid ounces of this product plus 0.5 pound a.i. of 2,4-D in 3 to 10 gallons of water per acre in the late summer or fall.			
<b>Sowthistle, perennial</b>	<b>3 - 4.5</b>	<b>3 - 40</b>	<b>1.5%</b>
Apply when most plants are at or beyond the bud stage of growth. After harvest, mowing or tillage in the late summer or fall, allow at least 4 weeks for initiation of active growth and rosette development prior to the application of this product. Fall treatments must be applied before a killing frost. Allow 3 or more days after application before tillage.			
<b>Spurge, leafy</b>	<b>—</b>	<b>3 - 10</b>	<b>1.5%</b>
For suppression, apply 12 fluid ounces of this product plus 0.5 pound a.i. 2,4-D in 3 to 10 gallons of water per acre in the late summer or fall. If mowing has occurred prior to treatment, apply when most of the plants are 12 inches tall.			
<b>Starthistle, yellow</b>	<b>3</b>	<b>10 - 40</b>	<b>1.5%</b>
Best results are obtained when applications are made during the rosette, bolting and early flower stages.			
<b>Sweet potato, wild</b>	<b>—</b>	<b>—</b>	<b>1.5%</b>
Suppression. Apply to plants that are at or beyond the bloom stage of growth. Repeat applications may be required.			
<b>Thistle, artichoke</b>	<b>—</b>	<b>—</b>	<b>1.5%</b>
Suppression. Apply to plants that are at or beyond the bloom stage of growth. Repeat applications may be required.			
<b>Thistle, Canada</b>	<b>3 - 4.5</b>	<b>3 - 40</b>	<b>1.5%</b>
Apply when most plants are at or beyond the bud stage of growth. After harvest, mowing or tillage in the late summer or fall, allow at least 4 weeks for initiation of active growth and rosette development prior to the application of this product. Fall treatments must be applied before a killing frost. Allow 3 or more days after application before tillage.			
For suppression, apply 1.5 pints of this product, or 12 fluid ounces of this product plus 0.5 pound a.i. 2,4-D, in 3 to 10 gallons of water per acre in the late summer or fall after harvest, mowing or tillage. Allow rosette regrowth to a minimum of 6 inches in diameter before treating. Applications can be made as long as leaves are still green and plants are actively growing at the time of application. Allow 3 or more days after application before tillage.			
<b>Timothy</b>	<b>3 - 4.5</b>	<b>3 - 40</b>	<b>1.5%</b>
For best results, apply when most plants have reached the boot-to-head stage of growth.			
<b>Torpedograss</b>	<b>6 - 7.5</b>	<b>3 - 40</b>	<b>1.5%</b>
For suppression. Apply when most plants are at or beyond the seedhead stage of growth. Repeat applications will be required to maintain control. Fall treatments must be applied before frost.			
<b>Trumpetcreeper</b>	<b>3</b>	<b>5 - 10</b>	<b>1.5%</b>
Suppression. Apply in late September or October, to plants that are at least 18 inches tall and have been growing 45 to 60 days since the last tillage operation. Make applications at least 1 week before a killing frost.			
<b>Vaseygrass</b>	<b>4.5 - 7.5</b>	<b>3 - 20</b>	<b>1.5%</b>
Apply when most plants are in the early head stage.			
<b>Velvetgrass</b>	<b>4.5 - 7.5</b>	<b>3 - 20</b>	<b>1.5%</b>
Apply when most plants are in the early head stage.			
<b>Wheatgrass, western</b>	<b>3 - 4.5</b>	<b>3 - 40</b>	<b>1.5%</b>
For best results, apply when most plants have reached the boot-to-head stage of growth.			

## Woody Brush And Trees Rate Table (Alphabetically By Species)

Apply this product after full leaf expansion, unless otherwise directed. Use the higher rate for larger plants and/or dense areas of growth. On vines, use the higher rate for plants that have reached the woody stage of growth. Best results are obtained when application is made in late summer or fall after fruit formation.

In arid areas, best results are obtained when applications are made in the spring to early summer when brush species are at high moisture content and are flowering.

Ensure thorough coverage when using hand-held equipment. Symptoms may not appear prior to frost or senescence with fall treatments.

Allow 7 or more days after application before tillage, mowing or removal. Repeat treatments may be necessary to control plants regenerating from underground parts or seed. Some autumn colors on undesirable deciduous species are acceptable provided no major leaf drop has occurred. Reduced performance may result if fall treatments are made following a frost.

For difficult to control perennial weeds and woody brush and trees, where plants are growing under stressed conditions, or where infestations are dense, this product may be used at 5 to 10 quarts per acre for enhanced results. The annual maximum use rate for this product is 10.6 qt per acre per year.

<b>Weed Species</b>	<b>Rate (pt / acre)</b>	<b>Water Volume (gpa)</b>	<b>Hand-Held (% Solution)</b>
<b>Alder</b>	<b>4.5 - 6</b>	<b>3 - 40</b>	<b>0.75 - 1.5%</b>
For control			
<b>Ash</b>	<b>3 - 7.5</b>	<b>3 - 40</b>	<b>0.75 - 1.5%</b>
For suppression			
<b>Aspen, quaking</b>	<b>3 - 4.5</b>	<b>3 - 40</b>	<b>0.75 - 1.5%</b>
For control			
<b>Bearmat (Bearclover)</b>	<b>3 - 7.5</b>	<b>3 - 40</b>	<b>0.75 - 1.5%</b>
For suppression			
<b>Beech</b>	<b>3 - 7.5</b>	<b>3 - 40</b>	<b>0.75 - 1.5%</b>
For suppression			
<b>Birch</b>	<b>3</b>	<b>3 - 40</b>	<b>0.75%</b>
For control			
<b>Blackberry</b>	<b>4.5 - 6</b>	<b>10 - 40</b>	<b>0.75 - 1.5%</b>
For control. Make applications after plants have reached full leaf maturity. Best results are obtained when applications are made in late summer or fall. Applications may also be made after leaf drop and until a killing frost or as long as stems are green. After berries have set or dropped in late fall, blackberry can be controlled by applying a 0.75% solution of this product. For control of blackberries after leaf drop and until killing frost or as long as stems are green, apply 4.5 to 6 pints of this product in 10 to 40 gallons of water per acre.			
<b>Blackgum</b>	<b>3 - 7.5</b>	<b>3 - 40</b>	<b>0.75 - 1.5%</b>
For control			
<b>Bracken</b>	<b>3 - 7.5</b>	<b>3 - 40</b>	<b>0.75 - 1.5%</b>
For control			
<b>Broom; French, Scotch</b>	<b>-</b>	<b>-</b>	<b>1.5%</b>
For control			
<b>Buckwheat, California</b>	<b>-</b>	<b>-</b>	<b>0.75 - 1.5%</b>
For suppression. Thorough coverage of foliage is necessary for best results.			
<b>Cascara</b>	<b>3 - 7.5</b>	<b>3 - 40</b>	<b>0.75 - 1.5%</b>
Suppression			
<b>Catsclaw</b>	<b>-</b>	<b>-</b>	<b>0.75 - 1.5%</b>
Suppression			
<b>Ceanothus</b>	<b>3 - 7.5</b>	<b>3 - 40</b>	<b>0.75 - 1.5%</b>
Suppression			

<b>Weed Species (Cont.)</b>	<b>Rate (pt / acre)</b>	<b>Water Volume (gpa)</b>	<b>Hand-Held (% Solution)</b>
<b>Chamise</b>	-	-	0.75%
For control. Thorough coverage of foliage is necessary for best results.			
<b>Cherry; bitter, black, pin</b>	3 - 4.5	3 - 40	0.75 - 1.5%
For control			
<b>Coyote brush</b>	-	-	1.5%
For control. Apply when at least 50 percent of the new leaves are fully developed.			
<b>Dogwood</b>	3 - 7.5	3 - 40	0.75 - 1.5%
Suppression			
<b>Elderberry</b>	3	3 - 40	0.75%
For control			
<b>Elm</b>	3 - 7.5	3 - 40	0.75 - 1.5%
Suppression			
<b>Eucalyptus</b>	-	-	1.5%
For control of eucalyptus resprouts, apply when resprouts are 6 to 12 feet tall. Ensure complete coverage. Avoid application to drought-stressed plants.			
<b>Florida holly (Brazilian Peppertree)</b>	3 - 7.5	3 - 40	0.75 - 1.5%
Suppression			
<b>Gorse</b>	3 - 7.5	3 - 40	0.75 - 1.5%
Suppression			
<b>Hasardia</b>	-	-	0.75 - 1.5%
Suppression. Thorough coverage of foliage is necessary for best results.			
<b>Hawthorn</b>	3 - 4.5	3 - 40	0.75 - 1.5%
For control			
<b>Hazel</b>	3	3 - 40	0.75%
For control			
<b>Hickory</b>	3 - 7.5	3 - 40	0.75 - 1.5%
Suppression			
<b>Honeysuckle</b>	3 - 6	3 - 40	0.75 - 1.5%
For control			
<b>Hornbeam, American</b>	3 - 7.5	3 - 40	0.75 - 1.5%
Suppression			
<b>Kudzu</b>	6	3 - 40	1.5%
For control. Repeat applications may be required to maintain control.			
<b>Locust, black</b>	3 - 6	3 - 40	0.75 - 1.5%
Suppression			
<b>Madrone resprouts</b>	-	-	1.5%
Suppression. Apply to resprouts that are 3 to 6 feet tall. Best results are obtained with spring/early summer treatments.			
<b>Manzanita</b>	3 - 7.5	3 - 40	0.75 - 1.5%
Suppression			

<b>Weed Species (Cont.)</b>	<b>Rate (pt / acre)</b>	<b>Water Volume (gpa)</b>	<b>Hand-Held (% Solution)</b>
<b>Maple, red</b>	<b>3 - 6</b>	<b>3 - 40</b>	<b>0.75 - 1.5%</b>
For control, apply a 0.75 to 1.5 percent solution when at least 50 percent of the new leaves are fully developed. For suppression, apply 3 to 6 pints of this product per acre.			
<b>Maple, sugar</b>	-	-	<b>0.75 - 1.5%</b>
For control. Apply when at least 50 percent of the new leaves are fully developed.			
<b>Monkey flower</b>	-	-	<b>0.75 - 1.5%</b>
Suppression. Thorough coverage of foliage is necessary for best results.			
<b>Oak; black, white</b>	<b>3 - 6</b>	<b>3 - 40</b>	<b>0.75 - 1.5%</b>
Suppression			
<b>Oak, post</b>	<b>4.5 - 6</b>	<b>3 - 40</b>	<b>0.75 - 1.5%</b>
For control			
<b>Oak; northern, pin</b>	-	-	<b>0.75 - 1.5%</b>
For control. Apply when at least 50 percent of the new leaves are fully developed.			
<b>Oak; southern red</b>	<b>3 - 4.5</b>	<b>3 - 40</b>	<b>0.75 - 1.5%</b>
For control			
<b>Persimmon</b>	<b>3 - 7.5</b>	<b>3 - 40</b>	<b>0.75 - 1.5%</b>
Suppression			
<b>Pine</b>	<b>3 - 7.5</b>	<b>3 - 40</b>	<b>0.75 - 1.5%</b>
For control			
<b>Poison ivy/ Poison oak</b>	<b>6 - 7.5</b>	<b>3 - 40</b>	<b>1.5%</b>
For control. Repeat applications may be required to maintain control. Fall treatments must be applied before leaves lose green color.			
<b>Poplar, yellow</b>	<b>3 - 7.5</b>	<b>3 - 40</b>	<b>0.75 - 1.5%</b>
Suppression			
<b>Redbud, eastern</b>	<b>3 - 7.5</b>	<b>3 - 40</b>	<b>0.75 - 1.5%</b>
For control			
<b>Rose, multiflora</b>	<b>3</b>	<b>3 - 40</b>	<b>0.75%</b>
For control. Treatments should be made prior to leaf deterioration by leaf-eating insects.			
<b>Russian olive</b>	<b>3 - 7.5</b>	<b>3 - 40</b>	<b>0.75 - 1.5%</b>
Suppression			
<b>Sage, black</b>	-	-	<b>0.75%</b>
For control. Thorough coverage of foliage is necessary for best results.			
<b>Sage, white</b>	<b>3 - 7.5</b>	<b>3 - 40</b>	<b>0.75 - 1.5%</b>
Suppression			
<b>Sage brush, California</b>	-	-	<b>0.75%</b>
For control. Thorough coverage of foliage is necessary for best results.			
<b>Salmonberry</b>	<b>3</b>	<b>3 - 40</b>	<b>0.75%</b>
For control			
<b>Salt-cedar</b>	<b>3 - 7.5</b>	<b>3 - 40</b>	<b>0.75 - 1.5%</b>
For control			
<b>Sassafras</b>	<b>3 - 7.5</b>	<b>3 - 40</b>	<b>0.75 - 1.5%</b>
Suppression			

<b>Weed Species (Cont.)</b>	<b>Rate (pt / acre)</b>	<b>Water Volume (gpa)</b>	<b>Hand-Held (% Solution)</b>
<b>Sourwood</b>	<b>3 - 7.5</b>	<b>3 - 40</b>	<b>0.75 - 1.5%</b>
Suppression			
<b>Sumac; poison, smooth, winged</b>	<b>3 - 6</b>	<b>3 - 40</b>	<b>0.75 - 1.5%</b>
Suppression			
<b>Sweetgum</b>	<b>3 - 4.5</b>	<b>3 - 40</b>	<b>0.75 - 1.5%</b>
For control			
<b>Swordfern</b>	<b>3 - 7.5</b>	<b>3 - 40</b>	<b>0.75 - 1.5%</b>
Suppression			
<b>Tallowtree, Chinese</b>	<b>-</b>	<b>-</b>	<b>0.75%</b>
For control. Thorough coverage of foliage is necessary for best results.			
<b>Tan oak resprouts</b>	<b>-</b>	<b>-</b>	<b>1.5%</b>
For suppression. Apply to resprouts that are less than 3 to 6 feet tall. Best results are obtained with fall applications.			
<b>Thimbleberry</b>	<b>3</b>	<b>3 - 40</b>	<b>0.75%</b>
For control			
<b>Tobacco, tree</b>	<b>-</b>	<b>-</b>	<b>0.75 - 1.5%</b>
Suppression			
<b>Trumpet creeper</b>	<b>3 - 4.5</b>	<b>3 - 40</b>	<b>0.75 - 1.5%</b>
For control			
<b>Vine maple</b>	<b>3 - 7.5</b>	<b>3 - 40</b>	<b>0.75 - 1.5%</b>
Suppression			
<b>Virginia creeper</b>	<b>3 - 7.5</b>	<b>3 - 40</b>	<b>0.75 - 1.5%</b>
For control			
<b>Waxmyrtle, southern</b>	<b>3 - 7.5</b>	<b>3 - 40</b>	<b>0.75 - 1.5%</b>
Suppression			
<b>Willow</b>	<b>3</b>	<b>3 - 40</b>	<b>0.75%</b>
For control			

## Terms and Conditions of Use

If terms of the following Warranty Disclaimer, Inherent Risks of Use and Limitation of Remedies are not acceptable, return unopened package at once to the seller for a full refund of purchase price paid. To the extent permitted by law, otherwise, use by the buyer or any other user constitutes acceptance of the terms under Warranty Disclaimer, Inherent Risks of Use and Limitation of Remedies.

## Warranty Disclaimer

Dow AgroSciences warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions, subject to the inherent risks set forth below. To the extent permitted by law, Dow AgroSciences MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.

## Inherent Risks of Use

It is impossible to eliminate all risks associated with use of this product. Crop injury, lack of performance, or other unintended consequences may result because of such factors as use of the product contrary to label instructions (including conditions noted on the label, such as unfavorable temperatures, soil conditions, etc.), abnormal conditions (such as excessive rainfall, drought, tornadoes, hurricanes), presence of other materials, the manner of application, or other factors, all of which are beyond the control of Dow AgroSciences or the seller. To the extent permitted by law, all such risks shall be assumed by buyer.

---

## Limitation of Remedies

To the extent permitted by law, the exclusive remedy for losses or damages resulting from this product (including claims based on contract, negligence, strict liability, or other legal theories), shall be limited to, at Dow AgroSciences' election, one of the following:

1. Refund of purchase price paid by buyer or user for product bought, or
2. Replacement of amount of product used.

To the extent permitted by law, Dow AgroSciences shall not be liable for losses or damages resulting from handling or use of this product unless Dow AgroSciences is promptly notified of such loss or damage in writing. To the extent permitted by law, in no case shall Dow AgroSciences be liable for consequential or incidental damages or losses.

The terms of the Warranty Disclaimer, Inherent Risks of Use and this Limitation of Remedies cannot be varied by any written or verbal statements or agreements. No employee or sales agent of Dow AgroSciences or the seller is authorized to vary or exceed the terms of the Warranty Disclaimer or this Limitation of Remedies in any manner.

®Trademark of Dow AgroSciences LLC  
EPA-accepted 05/12/08

# Accord<sup>®</sup> XRT II

## Herbicide

A non-selective broad-spectrum systemic herbicide for control of annual and perennial weeds and woody plants in noncrop areas and industrial sites, such as utility, railroad and roadside rights-of-way, airports, natural and production (plantations) forests for site preparation, mid-rotation release treatments, and timber stand improvement activities, wildlife and habitat management areas, wildlife openings, natural areas, such as wildlands, campgrounds, parks and recreational areas, wildlife refuges, rangeland and in and around seasonally dry wetlands including ditchbanks, dry ditches and dry canals and grazed areas on listed noncrop sites

**Avoid contact of herbicide with foliage, green stems, exposed non-woody roots or fruit of crops, desirable plants and trees, because severe injury or destruction may result.**

### Active Ingredient:

glyphosate: N-(phosphonomethyl)glycine, dimethylamine salt.....	50.2%
Other Ingredients.....	49.8%
Total Ingredients.....	100.0%

Contains 5.4 pounds per gallon glyphosate, dimethylamine salt (4 pounds per gallon glyphosate acid).

**Keep Out of Reach of Children**

## CAUTION

### Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. Refer to the label booklet under "Agricultural Use Requirements" in the Directions for Use section for information about this standard.

**For additional Precautionary Statements, First Aid, Storage and Disposal and other use information see inside this label.**

**Notice:** Read the entire label. Use only according to label directions. **Before using this product, read Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies at end of label booklet. If terms are unacceptable, return at once unopened.**

In case of emergency endangering health or the environment involving this product, call 1-800-992-5994. If you wish to obtain additional product information, visit our web site at [www.dowagro.com](http://www.dowagro.com).

Agricultural Chemical: Do not ship or store with food, feeds, drugs or clothing.

EPA Reg. No. 62719-556 900-014750 / 00292630

®Trademark of Dow AgroSciences LLC  
**Produced for**  
**Dow AgroSciences LLC**  
9330 Zionsville Road  
Indianapolis, IN 46268

**Net Contents 2.5 gal**



# Accord<sup>®</sup> XRT II

## Herbicide

A non-selective broad-spectrum systemic herbicide for control of annual and perennial weeds and woody plants in noncrop areas and industrial sites, such as utility, railroad and roadside rights-of-way, airports, natural and production (plantations) forests for site preparation, mid-rotation release treatments, and timber stand improvement activities, wildlife and habitat management areas, wildlife openings, natural areas, such as wildlands, campgrounds, parks and recreational areas, wildlife refuges, rangeland and in and around seasonally dry wetlands including ditchbanks, dry ditches and dry canals and grazed areas on listed noncrop sites

Avoid contact of herbicide with foliage, green stems, exposed non-woody roots or fruit of crops, desirable plants and trees, because severe injury or destruction may result.

Active Ingredient:  
glyphosate: N-(phosphonomethyl)glycine, dimethylamine salt .....50.2%  
Other Ingredients .....49.8%  
Total Ingredients .....100.0%

Contains 5.4 pounds per gallon glyphosate, dimethylamine salt (4 pounds per gallon glyphosate acid).

## Keep Out of Reach of Children CAUTION

### Precautionary Statements

#### Hazards to Humans and Domestic Animals

Causes Moderate Eye Irritation • Prolonged or Frequently Repeated Skin Contact May Cause Allergic Reactions in Some Individuals

Avoid contact with skin, eyes or clothing.

#### Personal Protective Equipment (PPE)

Some of the materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for category A on an EPA chemical-resistance category selection chart.

#### Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Chemical-resistant gloves made of any waterproof material such as natural rubber
- Shoes plus socks

Follow manufacturer's instructions for cleaning/maintaining PPE (Personal Protective Equipment). If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

#### Engineering Controls

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in

#### Engineering Controls (Cont.)

Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d) (4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

#### User Safety Recommendations

Users should wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.

#### First Aid

**If in eyes:** Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing. Call a poison control center or doctor for treatment advice.

**If on skin:** Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact 1-800-992-5994 for emergency medical treatment information.

**Domestic Animals:** This product is considered to be relatively nontoxic to dogs and other domestic animals; however, ingestion of this product or large amounts of freshly sprayed vegetation may result in temporary gastrointestinal irritation (vomiting, diarrhea, colic, etc.). If such symptoms are observed, provide the animal with plenty of fluids to prevent dehydration. Call a veterinarian if symptoms persist for more than 24 hours.

#### Environmental Hazards

Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when cleaning equipment or disposing of equipment washwaters.

#### Physical or Chemical Hazards

Spray solutions of this product should be mixed, stored and applied using only stainless steel, aluminum, fiberglass, plastic or plastic-lined steel containers.

**Do not mix, store or apply this product or spray solutions of this product in galvanized steel or unlined steel (except stainless steel) containers or spray tanks.**

This product, or spray solutions of this product react with such containers and tanks to produce hydrogen gas that may form a highly combustible gas mixture. This gas mixture could flash or explode, causing serious personal injury, if ignited by open flame, spark, welder's torch, lighted cigarette or other ignition source.

#### Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. Refer to the label booklet under "Agricultural Use Requirements" in the Directions for Use section for information about this standard.

#### Storage and Disposal

**Pesticide Storage:** Do not contaminate water, food, feed or seed by storage or disposal.

**Pesticide Disposal:** Wastes of this pesticide may cause eye and skin irritation and may be dangerous. Improper disposal of excess pesticide, spray mixtures, or rinsate is a violation of Federal law. If these wastes cannot be disposed of according to label use instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

**Container Reuse:** Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available.

Triple rinse or pressure rinse container (or equivalent) promptly after emptying. **Triple rinse** as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times.

**Pressure rinse** as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank and collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 psi for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

#### Refer to label booklet for Directions for Use.

**Notice:** Read the entire label. Use only according to label directions. **Before using this product, read Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies at end of label booklet. If terms are unacceptable, return at once unopened.**

In case of emergency endangering health or the environment involving this product, call 1-800-992-5994. If you wish to obtain additional product information, visit our web site at [www.dowagro.com](http://www.dowagro.com).


Agricultural Chemical: Do not ship or store with food, feeds, drugs or clothing.

EPA Reg. No. 62719-556 EPA Est. 34704-MS-1

®Trademark of Dow AgroSciences LLC

Produced for  
Dow AgroSciences LLC  
9330 Zionsville Road  
Indianapolis, IN 46268

00293833



# Accord<sup>®</sup> XRT II

2 x 2.5 gal

Herbicide

2 x 2.5 gal



# Accord<sup>®</sup> XRT II

Herbicide

2 x 2.5 gal

00293833



# Accord<sup>®</sup> XRT II

Herbicide

2 x 2.5 gal

4G/Z31.2/S/08  
USA/+AA2111

# Accord<sup>®</sup> XRT II

## Herbicide

A non-selective broad-spectrum systemic herbicide for control of annual and perennial weeds and woody plants in noncrop areas and industrial sites, such as utility, railroad and roadside rights-of-way, airports, natural and production (plantations) forests for site preparation, mid-rotation release treatments, and timber stand improvement activities, wildlife and habitat management areas, wildlife openings, natural areas, such as wildlands, campgrounds, parks and recreational areas, wildlife refuges, rangeland and in and around seasonally dry wetlands including ditchbanks, dry ditches and dry canals and grazed areas on listed noncrop sites

Avoid contact of herbicide with foliage, green stems, exposed non-woody roots or fruit of crops, desirable plants and trees, because severe injury or destruction may result.

Active Ingredient:

glyphosate: N-(phosphonomethyl)glycine, dimethylamine salt .....	50.2%
Other Ingredients .....	49.8%
Total Ingredients .....	100.0%

Contains 5.4 pounds per gallon glyphosate, dimethylamine salt (4 pounds per gallon glyphosate acid).

## Keep Out of Reach of Children

# CAUTION

### Precautionary Statements

### Hazards to Humans and Domestic Animals

**Causes Moderate Eye Irritation • Prolonged or Frequently Repeated Skin Contact May Cause Allergic Reactions in Some Individuals**

Avoid contact with skin, eyes or clothing.

### Personal Protective Equipment (PPE)

Some of the materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for category A on an EPA chemical-resistance category selection chart.

#### Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Chemical-resistant gloves made of any waterproof material such as natural rubber
- Shoes plus socks

Follow manufacturer's instructions for cleaning/maintaining PPE (Personal Protective Equipment). If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

### Engineering Controls

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in

### Engineering Controls (Cont.)

Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d) (4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

### User Safety Recommendations

Users should wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.

### First Aid

**If in eyes:** Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing. Call a poison control center or doctor for treatment advice.

**If on skin:** Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact 1-800-992-5994 for emergency medical treatment information.

**Domestic Animals:** This product is considered to be relatively nontoxic to dogs and other domestic animals; however, ingestion of this product or large amounts of freshly sprayed vegetation may result in temporary gastrointestinal irritation (vomiting, diarrhea, colic, etc.). If such symptoms are observed, provide the animal with plenty of fluids to prevent dehydration. Call a veterinarian if symptoms persist for more than 24 hours.

### Environmental Hazards

Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when cleaning equipment or disposing of equipment washwaters.

### Physical or Chemical Hazards

Spray solutions of this product should be mixed, stored and applied using only stainless steel, aluminum, fiberglass, plastic or plastic-lined steel containers.

**Do not mix, store or apply this product or spray solutions of this product in galvanized steel or unlined steel (except stainless steel) containers or spray tanks.** This product, or spray solutions of this product react with such containers and tanks to produce hydrogen gas that may form a highly combustible gas mixture. This gas mixture could flash or explode, causing serious personal injury, if ignited by open flame, spark, welder's torch, lighted cigarette or other ignition source.

### Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. Refer to the label booklet under "Agricultural Use Requirements" in the Directions for Use section for information about this standard.

### Storage and Disposal

**Pesticide Storage:** Do not contaminate water, food, feed or seed by storage or disposal.

**Pesticide Disposal:** Wastes of this pesticide may cause eye and skin irritation and may be dangerous. Improper disposal of excess pesticide, spray mixtures, or rinsate is a violation of Federal law. If these wastes cannot be disposed of according to label use instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

**Container Reuse:** Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available. Triple rinse or pressure rinse container (or equivalent) promptly after emptying. **Triple rinse** as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. **Pressure rinse** as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 psi for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

### Refer to label booklet for Directions for Use.

**Notice:** Read the entire label. Use only according to label directions. **Before using this product, read Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies at end of label booklet. If terms are unacceptable, return at once unopened.**

In case of emergency endangering health or the environment involving this product, call 1-800-992-5994. If you wish to obtain additional product information, visit our web site at [www.dowagro.com](http://www.dowagro.com).

Agricultural Chemical: Do not ship or store with food, feeds, drugs or clothing.

EPA Reg. No. 62719-556

EPA Est. 34704-MS-1

®Trademark of Dow AgroSciences LLC  
**Produced for**  
**Dow AgroSciences LLC**  
9330 Zionville Road  
Indianapolis, IN 46268

# 2 x 2.5 gal

# Supplemental Labeling



Dow AgroSciences LLC

9330 Zionsville Road

Indianapolis, IN 46268-1054 USA

## Accord<sup>®</sup> XRT II

EPA Reg. No. 62719-556

### Control and Management of Glyphosate-Resistant Horseweed (*Marestail*, *Conyza canadensis*) in Noncrop Areas and Industrial Sites (Not for Use or Distribution Within the State of California)

#### ATTENTION

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application.
- Read the label affixed to the container for this product before applying. Carefully follow all precautionary statements and applicable use directions.
- Use of this product according to this supplemental labeling is subject to all use precautions and limitations imposed by the label affixed to the product container.

#### Directions for Use

Read the Product Information, Herbicide Resistance Management, Spray Drift Management, Mixing Directions, and Application Equipment and Application Methods on the label attached to this product's container.

Use this product to control and manage glyphosate-resistant horseweed (*marestail*, *Conyza canadensis*). Apply 24 fl oz of this product per acre before *marestail* is more than 6 inches in height. Make applications when horseweed is still in the rosette stage of growth to enhance control.

**Tank Mixes:** The following products may be tank mixed with this product provided that the product is labeled for application at the target site. Refer to the label of the tank mix partner for use sites and applications rates. Read and carefully follow the label directions on the tank mix partner. Use according to the most restrictive precautionary statements for each product in the mixture.

2,4-D <sup>1</sup>	Landmark II MP	Telar DF
Crossbow L	Landmark MP	Transline
dicamba <sup>1</sup>	Milestone	Vanquish
Gallery 75 DF	Overdrive	Velpar DF
Krovar I DF		

<sup>1</sup>Tank mixes with products containing this generic active ingredient may be made provided the specific product is labeled for application on the use site.

®Trademark of Dow AgroSciences LLC

R351-014  
EPA accepted 11/26/10  
Replaces R351-007

# Supplemental Labeling



Dow AgroSciences LLC

9330 Zionsville Road

Indianapolis, IN 46268-1054 USA

## Accord<sup>®</sup> XRT II

EPA Reg. No. 62719-556

### Control and Management of Glyphosate-Resistant Ryegrass (Not for Use or Distribution Within the State of California)

#### ATTENTION

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application.
- Read the label affixed to the container for this product before applying. Carefully follow all precautionary statements and applicable use directions.
- Use of this product according to this supplemental labeling is subject to all use precautions and limitations imposed by the label affixed to the product container.

#### Directions for Use

Read the Product Information, Herbicide Resistance Management, Spray Drift Management, Mixing Directions, and Application Equipment and Application Methods on the label attached to this product's container.

**Preemergence:** To control other emerged weeds, apply this product in a tank mix with a preemergence herbicide labeled for control of ryegrass.

**Preemergence and Postemergence:** To control other emerged weeds, apply this product in a tank mix with a residual preemergence herbicide and a postemergence herbicide (other than glyphosate) labeled for control of ryegrass. Apply before ryegrass is more than 4 inches in height.

**Postemergence:** To control other emerged weeds, apply this product in a tank mix with another postemergence herbicide labeled for control of ryegrass. Apply before ryegrass is more than 4 inches in height.

Not all herbicides are registered in each state or for all use sites (orchards, noncrop areas, or ditch banks) or crops for the management of ryegrass. When using this product in a tank mix, refer to each product's label and observe the most restrictive label's precautions and limitations.

<sup>®</sup>Trademark of Dow AgroSciences LLC

R351-015  
EPA accepted 11/26/10  
Replaces R351-008

# Supplemental Labeling



Dow AgroSciences LLC

9330 Zionsville Road

Indianapolis, IN 46268-1054 USA

## Accord<sup>®</sup> XRT II

EPA Reg. No. 62719-556

**Tank Mix Products for Noncrop Areas**  
(Not for Use or Distribution Within the State of California)

### ATTENTION

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application.
- Read the label affixed to the container for this product before applying. Carefully follow all precautionary statements and applicable use directions.
- Use of this product according to this supplemental labeling is subject to all use precautions and limitations imposed by the label affixed to the product container.

### Directions for Use

Read the Product Information, Herbicide Resistance Management, Spray Drift Management, Mixing Directions, and Application Equipment and Application Methods on the label attached to this product's container.

See the rate tables in the Annual Weeds, Perennial Weeds, and Woody Brush and Trees sections of the the label affixed to the product container for specific application rates. For improved results, apply 4 to 7 quarts of this product per acre may be used for difficult to control species, where dense stands occur, or where conditions for control are not ideal. The following herbicides, when applied as a tank mix, provide preemergence and/or postemergence control of the weeds listed in the individual product tables.

The following list of products may be tank mixed with this product provided that the product is labeled for use on the target site. Any specified rate of this product may be used in a tank mix with these products. Refer to the product labels for specified sites and application rates.

Tank Mix Product	Rate per Acre
Arsenal <sup>1</sup>	0.5 – 4 pints
2,4-D	0.5 – 1 lb
Garlon 3A	1 – 6 pints
Garlon 4	1 – 6 pints
diuron	4 – 8 lb
diuron + 2,4-D	4 – 8 lb + 0.5 – 1 lb
diuron + Garlon 3A	4 – 10 lb + 1 – 2 pints
diuron + Garlon 4	4 – 10 lb + 1 – 2 pints
Hyvar X	4 – 8 lb
Hyvar X + 2,4-D	4 – 8 lb + 0.5 – 1 lb
Hyvar X + Garlon 3A	4 – 8 lb + 1 – 2 pints
Hyvar X + Garlon 4	4 – 8 lb + 1 – 2 pints
Krovar I DF	4 – 6 lb
Krovar I DF + 2,4-D	4 – 6 lb + 0.5 – 1 lb
Krovar I DF + Garlon 3A	4 – 6 lb + 1 – 2 pints
Krovar I DF + Garlon 4	4 – 6 lb + 1 – 2 pints
Oust	2 – 6 oz
Oust + 2,4-D	2 – 6 oz + 0.5 – 1 lb
Oust + Garlon 3A	2 – 6 oz + 1 – 2 pints

<b>Tank Mix Product</b>	<b>Rate per Acre</b>
Oust + Garlon 4	2 – 6 oz + 1 – 2 pints
Spike 80W	2 – 5 lb
Spike 80W + 2,4-D	2 – 5 lb + 0.5 – 1 lb
Spike 80W + Garlon 3A	2 – 5 lb + 1 – 2 pints
Spike 80W + Garlon 4	2 – 5 lb + 1 – 2 pints

<sup>1</sup>Arsenal is not approved for use in the state of California.

Read and carefully follow the label claims, precautionary statements, specified use rates and all other information on the labels of the products used in the tank mix. Use according to the most restrictive label directions for each product in the mixture.

During the mixing process and during application, maintain good agitation at all times. Ensure that the tank mix product(s) is well mixed with the spray solution before adding this product. Mix only the amount of spray solution that will be used during the same day. Reduced weed control may result if a tank mixture is allowed to stand overnight. If the spray mix is allowed to settle, thorough agitation is required to resuspend the mixture before spraying is resumed.

®Trademark of Dow AgroSciences LLC

R351-016  
EPA accepted 11/26/10  
Replaces R351-009

# Supplemental Labeling



Dow AgroSciences LLC

9330 Zionsville Road

Indianapolis, IN 46268-1054 USA

## Accord<sup>®</sup> XRT II

EPA Reg. No. 62719-556

**Tank Mixtures for Improved Control of Bentgrass (*Agrostis* spp.)**  
(Not for Use or Distribution Within the State of California)

### ATTENTION

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application.
- Read the label affixed to the container for this product before applying. Carefully follow all precautionary statements and applicable use directions.
- Use of this product according to this supplemental labeling is subject to all use precautions and limitations imposed by the label affixed to the product container.

### Directions for Use

Read the Product Information, Herbicide Resistance Management, Spray Drift Management, Mixing Directions, and Application Equipment and Application Methods on the label attached to this product's container.

For improved control of bentgrass (*Agrostis* spp.), the following products may be tank mixed with this product: Envoy, Fusion, Fusilade II, Vantage. When tank mixing products, read and carefully observe label directions, precautionary statements and all information on the labels of each product in the mixture. Refer to each product label for the approved use sites.

Dry ammonium sulfate, at 1 to 2 percent by weight, may also be added to the spray solution. The equivalent rate of ammonium sulfate in a liquid formulation may also be used. Completely dissolve the ammonium sulfate in the spray tank before adding herbicides. Thoroughly rinse the spray system with clean water after use to reduce corrosion.

**Broadcast Application:** Apply 2 to 2.5 quarts of this product per acre plus

- 34 fl oz of Envoy per acre in 20 to 40 gallons per acre of spray solution.
- 24 fl oz of Fusilade II per acre in 20 to 40 gallons per acre of spray solution.
- 60 fl oz of Vantage per acre in 20 to 40 gallons per acre of spray solution.
- 9 fl oz of Fusion per acre in 20 to 40 gallons per acre of spray solution.

In the event of incomplete control, re-treatment may be necessary.

**Spot Treatment:** Mix 2 fl oz of this product with

- 1.3 fl oz of Envoy in 1 gallon of water and spray to wet.
- 0.75 fl oz of Fusilade II in 1 gallon of water and spray to wet.
- 3 fl oz of Vantage in 1 gallon of water and spray to wet.
- 0.25 fl oz of Fusion in 1 gallon of water and spray to wet.

**Attention:** Avoid drift. Use extreme care when applying this product to prevent injury to desirable plants and crops.

®Trademark of Dow AgroSciences LLC

R351-017

EPA accepted 11/26/10

Replaces R351-010

# Supplemental Labeling

LICENSED

LICENSED

PERIOD 2011-2013 LIC. NO. PERIOD 2014-2016 LIC. NO. 9786.276



Dow AgroSciences LLC

9330 Zionsville Road

Indianapolis, IN 46268-1054 USA

## Accord<sup>®</sup> XRT II

EPA Reg. No. 62719-556

### Hollow Stem Injection

#### ATTENTION

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application.
- Read the label affixed to the container for this product before applying. Carefully follow all precautionary statements and applicable use directions.
- Use of this product according to this supplemental labeling is subject to all use precautions and limitations imposed by the label affixed to the product container.

#### Directions for Use

Read the Product Information, Herbicide Resistance Management, Spray Drift Management, Mixing Directions, and Application Equipment and Application Methods on the label attached to this product's container.

Apply this product through hand-held injection devices that deliver the specified amount of this product into targeted hollow stem plants growing in any noncrop site listed on this label. To control the following hollow stem plants, follow the use directions below:

Target Plants		Use Directions
Common Name	Scientific Name	
castorbean	<i>Ricinus communis</i>	Inject 4 mL of this product per plant into the lower portion of the main stem
hemlock poison	<i>Conium maculatum</i>	Inject one leaf cane per plant, 10 to 12 inches above the root crown, with 5 mL of a 5 percent by volume solution of this product.
hogweed, giant	<i>Hercleum mantegazzianum</i>	Inject one leaf cane per plant 12 inches above the root crown with 5 mL of a 5 percent by volume solution of this product.
horsetail field	<i>Equisetum arvense</i>	Inject one segment above the root crown with 0.5 mL of this product per stem using a low volume syringe capable of accurately delivering this amount of product.
knotweed, bohemian	<i>Polygonum bohemicum</i>	Inject 5 mL of this product per stem between the second and third internode.
knotweed, Japanese	<i>Polygonum cuspidatum</i>	
reed, giant	<i>Arundo donax</i>	Inject 6 mL of this product per stem between the second and third internode.
thistle, Canada	<i>Cirsium arvense</i>	Cut 8 to 9 of the tallest plants at bud stage in a clump with clippers. Use a cavity needle pushed into the stem center and then slowly removed as 0.5 mL of this product per stem is injected into the stem.

**Precautions and Restrictions:**

- Do not apply more than a total of 8 quarts of this product per acre for all treatments combined. At 5 mL per stem, 8 quarts will treat approximately 1300 stems per acre.

®Trademark of Dow AgroSciences LLC

R351-011  
EPA accepted 11/26/10  
Initial printing

# Supplemental Labeling



Dow AgroSciences LLC

9330 Zionsville Road

Indianapolis, IN 46268-1054 USA

## Accord<sup>®</sup> XRT II

EPA Reg. No. 62719-556

### Weed Control in Non-Food Tree, Shrub, or Vine Production Sites (Not for Use or Distribution Within the State of California)

#### ATTENTION

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application.
- Read the label affixed to the container for this product before applying. Carefully follow all precautionary statements and applicable use directions.
- Use of this product according to this supplemental labeling is subject to all use precautions and limitations imposed by the label affixed to the product container.

#### Directions for Use

Read the Product Information, Herbicide Resistance Management, Spray Drift Management, Mixing Directions, and Application Equipment and Application Methods on the label attached to this product's container.

**Types of Applications:** Site preparation, post-directed, trim and edge, wiper application

This product may be used for general weed control prior to the planting of and around established ornamentals or any woody tree, shrub, or vine species including arborvitae, azalea, boxwood, crabapple, eucalyptus, euonymus, fir, douglas fir, jojoba, hollies, lilac, magnolia, maple, oak, poplar, privet, pine, spruce and yew, growing in plant nurseries, on Christmas tree farms, or on other non-food tree production sites.

Do not use this product as an over the top broadcast spray in ornamentals and Christmas trees unless otherwise directed. Take care to avoid contact of spray, drift, or mist with foliage or green bark of desirable ornamental species.

Use this product to control weeds growing in and around greenhouses and shadehouses. During application, desirable vegetation must not be present. Air circulation fans must be turned off until after the application has dried.

#### Site Preparation

Use this product prior to planting any tree, shrub or vine, including Christmas tree species, in a nursery or production setting.

#### Post-Directed and Trim and Edge

Use this product as a post-directed spray around established woody ornamental species or to trim and edge around trees, buildings, sidewalks, roads, potted plants and other objects in a production setting. Protect desirable plants from the spray solution by using shields or coverings made of cardboard or other impermeable material.

**Wiper Application**

Use this product through wick or other suitable wiper applicators to control or partially control undesirable vegetation around established trees, shrubs, or vines. See Selective Equipment section of this label for further information about the proper use of wiper applicators.

®Trademark of Dow AgroSciences LLC

R351-018

EPA accepted 11/26/10

Replaces R351-012