



Dow AgroSciences



STATE OF HAWAII
Department of Agriculture

ACCEPTED

LICENSE NO.

9786.298

Gallery[®]

SC

SPECIALTY HERBICIDE

A preemergence herbicide for control of certain broadleaf weeds in:

- Established Turfgrass
- Landscape Ornamentals
- Container Grown Ornamentals
- Field Grown Ornamentals
- Groundcovers/ Perennials
- Non-Cropland
- Ornamental Bulbs
- Christmas Tree/Conifer Plantations

For additional Precautionary Statements, First Aid, Storage and Disposal and other use information see inside this label.

Notice: Read the entire label. Use only according to label directions. **Before using this product, read Warranty Disclaimer, Inherent Risks of Use and Limitation of Remedies at end of label booklet.** If terms are unacceptable, return at once unopened.

In case of emergency endangering health or the environment involving this product, call 1-800-992-5994.

Agricultural Chemical: Do not ship or store with food, feeds, drugs or clothing.

Shake Well Before Use – Avoid Freezing

EPA Reg. No. 62719-658

EPA Est. 11773-IA-001
900-021178 / 11079455

Group	21	HERBICIDE
-------	----	-----------

Active Ingredient:

isoxaben: N-[3-(1-ethyl-1-methylpropyl)-5-isoxazolyl]-2,6-dimethoxybenzamide and isomers	45.45%
Other Ingredients	54.55%
Total	100%

Contains 4.16 lb active ingredient per gallon.

Keep Out of Reach of Children CAUTION

Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. Refer to the label booklet under "Agricultural Use Requirements" in the Directions for Use section for information about this standard.

[®]Trademark of The Dow Chemical Company ("Dow") or an affiliated company of Dow

Produced for
Dow AgroSciences LLC
9330 Zionsville Road
Indianapolis, IN 46268

NET CONTENTS 2.5 GAL



Precautionary Statements

Hazards to Humans and Domestic Animals

CAUTION

Prolonged Or Frequently Repeated Skin Contact May Cause Allergic Reactions In Some Individuals

Personal Protective Equipment (PPE)

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Shoes plus socks
- Chemical-resistant gloves

Follow the manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

Environmental Hazards

Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Drift may result in reduced germination or emergence of non-target plants adjacent to treated area. Do not contaminate water when cleaning equipment or disposing of equipment washwaters.

Ground Water Advisory: This pesticide has properties and characteristics associated with chemicals detected in ground water. This chemical may leach into ground water if used in areas where soils are permeable, particularly where the water table is shallow.

Surface Water Advisory: This pesticide may impact surface water quality due to runoff of rain water. This is especially true for poorly draining soil and soils with shallow ground water. This product is classified as having a high potential for reaching surface water via runoff for several months or more after application.

Directions for Use

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Read all Directions for Use carefully before applying.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your state or tribe, consult the agency responsible for pesticide regulation.

Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls
- Chemical resistant gloves made of any waterproof material
- Shoes plus socks

Non-Agricultural Use Requirements

The requirements of this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Entry Restrictions for Non-WPS Uses: When this product is applied to turf and ornamental plantings in landscape settings and non-cropland areas, do not allow entry into treated areas until sprays have dried unless wearing coveralls, waterproof gloves, and shoes plus socks.

Storage and Disposal

Do not contaminate water, food or feed by storage or disposal.

Pesticide Storage: Store in original container. Do not store in direct sunlight. Do not store at temperatures above 120°F. In case of leak or spill, contain material and dispose as waste.

Pesticide Disposal: Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.

Container Handling: Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures allowed by state and local authorities. Triple rinse or pressure rinse container (or equivalent) promptly after emptying. **Triple rinse** as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times.

Pressure rinse as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain

Storage and Disposal (Cont.)

for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinseate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 psi for at least 30 seconds. Drain for 10 seconds after the flow begins to drip. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures allowed by state and local authorities.

Product Information

Gallery® SC specialty herbicide is a preemergence product for control of certain broadleaf weeds in established turfgrass, landscape ornamentals, container grown ornamentals, field grown ornamentals, groundcovers/perennials, ornamental bulbs, Christmas tree/conifer plantations and non-cropland areas. Apply Gallery SC in late summer to early fall, in early spring, or anytime prior to germination of target weeds, or immediately after cultivation.

Use Precautions and Restrictions

Gallery SC controls weeds germinating from seed. Gallery SC does not control established weeds, or weeds growing from stolons, rhizomes, or root pieces. Existing weeds should be controlled by cultivation or with postemergence herbicides. Weed residues, prunings, and trash should be removed or thoroughly mixed into the soil prior to application. Soil in non-turfgrass areas should be in good condition and free of clods at the time of application. Gallery SC is stable on the soil surface for up to 21 days, but must be incorporated by moisture to be effective. A single rainfall or sprinkler irrigation of 0.5 inches or more, or flood irrigation after application, is necessary to activate Gallery SC. If Gallery SC is not activated by rainfall or irrigation within 21 days after application, erratic weed control may result. In non-turfgrass areas, if weeds emerge due to lack of rainfall or irrigation, shallow cultivation to a depth of 1 to 2 inches will incorporate the herbicide and destroy existing weeds.

Treatment of Turfgrass or Ornamental Species Not Listed on the Label

Although this label contains a large number of ornamental species, it is not possible to include all of the ornamental plants that may be encountered in nursery or landscape settings. Users who wish to use Gallery SC on a plant species not listed on this label may determine the suitability for such use by treating a small area or small number of plants at a specified rate. Prior to treatment of larger areas, the treated area/plants should be observed for any sign of herbicidal injury during 30 to 60 days of typical growing conditions. The user assumes the responsibility for any plant damage or other liability resulting from use of Gallery SC on species not listed on this label.

Chemigation: Do not apply Gallery SC through any type of irrigation system.

Not for sale, use or distribution in New York State.

Weed Resistance Management:

Isoxaben, the active ingredient in this product, is a Group 21 herbicide based on the mode of action classification system of the Weed Science Society of America. Any weed

population may contain plants naturally resistant to Group 21 herbicides. Such resistant weed plants may not be effectively managed using Group 21 herbicides but may be effectively managed utilizing another herbicide alone or in mixtures from a different Group and/or by using cultural or mechanical practices. However, a herbicide mode of action classification by itself may not adequately address specific weeds that are resistant to specific herbicides. Consult your local company representative, state cooperative extension service, professional consultants or other qualified authorities to determine appropriate actions for treating specific resistant weeds.

Best Management Practices:

Proactively implementing diversified weed control strategies to minimize selection for weed populations resistant to one or more herbicides is recommended. A diversified weed management program may include the use of multiple herbicides with different modes of action and overlapping weed spectrum with or without tillage operations and/or other cultural practices. Research has demonstrated that using the labeled rate and directions for use is important to delay the selection for resistance. Scouting after a herbicide application is important because it can facilitate the early identification of weed shifts and/or weed resistance and thus provide direction on future weed management practices. One of the best ways to contain resistant populations is to implement measures to avoid allowing weeds to reproduce by seed or to proliferate vegetatively. Cleaning equipment between sites and avoiding movement of plant material between sites will greatly aid in retarding the spread of resistant weed seed.

Spray Drift Management:

Spray equipment and weather affect spray drift. Consider all factors when making application decisions. Where states have more stringent regulations, they must be observed. Avoiding spray drift is the responsibility of the applicator or grower. To reduce the potential for drift, the application equipment must be set to apply medium to coarse droplets (i.e., ASAE Standard 572) with corresponding spray pressure. Use high flow rate nozzles to apply the highest practical spray volume. With most nozzle types, narrower spray angles produce larger droplets. Follow the nozzle manufacturer's directions on pressure, orientation, spray volume, etc. in order to minimize drift and optimize coverage and control.

Wind: Avoid making applications when spray particles may be carried by air currents to areas where sensitive crops and non-target plants are growing. Do not spray near sensitive plants if wind is gusty, below 2 mph, or in excess of 10 mph and moving in the direction of adjacent areas of sensitive areas. Local terrain may influence wind patterns; the applicator must be familiar with local conditions and understand how they may impact spray drift.

Sensitive Areas: Sensitive areas to this product are defined as bodies of water (ponds, lakes, rivers, streams, and ditches), wetlands, habitats of endangered species, and non-labeled agricultural crop areas. Applicators must take all precautions necessary to keep spray drift from reaching those areas.

Temperature Inversion: A surface temperature inversion (i.e., increasing temperature with increasing altitude) greatly increases the potential for drift. Presence of ground fog is a good indicator of a surface temperature inversion. Do not apply during temperature inversions. Always make applications when there is

some air movement to determine the direction and distance of possible spray drift.

Boom Height: Set the boom and make applications at the lowest height that safely permits uniform coverage of the soil and minimizes droplet evaporation. Boom or nozzle shielding can reduce the effects of wind or air currents on drift. Verify that the shields do not interfere with uniform deposition of product prior to application.

Application Directions

Apply Gallery SC with a properly calibrated low pressure herbicide sprayer that provides uniform spray distribution. Nozzle screens should be no finer than 50 mesh (50 mesh is finer than 16 mesh). In-line screens and strainers should be no finer than 16 mesh. Apply Gallery SC in 10 gallons or more of water carrier per acre. As the spray volume decreases, the importance of accurate calibration and uniform application increases. Take precautions to avoid spray drift when applying Gallery SC. Drift may result in reduced germination or emergence of non-target plants adjacent to the treated area. Maintain agitation from mixing through application. Avoid boom overlaps that will increase rates above those specified. Calibrate application equipment prior to use according to manufacturer's directions. Check calibration frequently to be sure equipment is working properly and distributing spray uniformly.

Mixing Directions

Gallery SC - Alone

Check to be sure spray equipment is clean and not contaminated with other herbicides. Using clean water, fill the tank to 1/2 of the final volume required and start agitation. Slowly add the required quantity of Gallery SC to the spray tank, continue agitation and complete filling the tank. Maintain agitation during filling and throughout application. Agitation should be sufficient to create a rippling or rolling action on the liquid's surface. Sparger pipe agitation generally provides the best agitation. In-line screens and strainers should be no finer than 16 mesh.

If spraying and agitation is stopped, Gallery SC may settle to the bottom of the spray tank. If settling occurs, material must be re-suspended before continuing spray application. Clean the spray tank, lines and screens thoroughly after use.

Application Rate Conversion Table for Gallery SC

lb ai/A	fl oz per acre	fl oz per 1000 sq ft	mls per 1000 sq ft
0.50	16	0.3	10
0.75	23	0.5	16
1.00	31	0.7	21

To avoid crop injury, do not repeat applications of 31 fl oz per acre Gallery SC sooner than 60 days after a previous application of Gallery SC. Do not apply more than a total of 124 fl oz/A of Gallery SC per acre within a 12-month period.

Gallery SC - Tank Mix

Gallery SC may be applied in tank mix combination with labeled rates of other products provided (1) the tank mix product is labeled for the timing and method of application for the use site to be treated; (2) tank mixing with Gallery SC is not

prohibited by the label of the tank mix product; and (3) the tank mix combination is compatible as determined by a "jar test" described in the Tank Mix Compatibility Testing section below.

Add Gallery SC to the spray tank as described above and fill the spray tank to 3/4 of the final volume required. Add formulations in this order: (1) dry flowables, (2) wettable powders, (3) aqueous suspensions (such as Gallery SC), (4) flowables, (5) liquids, (6) solutions and emulsifiable or liquid concentrates. Allow two to three minutes between the addition of each product and agitate continuously until each product is completely dispersed in water, and add water to the final volume. Maintain agitation during filling and through application. If a build up of materials is observed on the walls of the spray tank, wash the tank with soapy water between fillings, rinse and then continue the spraying operation. Follow label directions for each material added to the tank.

Tank Mix Compatibility Testing: A jar test is recommended prior to tank mixing to ensure compatibility of Gallery SC and other pesticides. Use a clear glass quart jar with lid and mix the tank mix ingredients in their relative proportions. Invert the jar containing the mixture several times and observe the mixture for approximately 1/2 hour. If the mixture balls-up, forms flakes, sludges, jels, oily films or layers, or other precipitates, it is not compatible and the tank mix combination should not be used.

Weeds Controlled or Suppressed

Weeds controlled when applied at 16 fl oz per acre (0.3 fl oz or 10 mls per 1000 sq ft):

Common Name

aster, slender
bursage, annual
burweed, lawn
celery, wild
chickweed, common
clover, white
cludweed, purple
fiddleneck, coast
filaree, redstem
fleabane, blackleaved
fleabane, dwarf
groundcherry, lanceleaf
henbit
horseweed
knotweed, prostrate
lambquarters, common
mallow, little
mustard, Indian
mustard, wild
nightshade, black
pepperweed, Virginia
pigweed
pineappleweed
plantain, slender
purslane, common
radish, wild
ragweed, common
rocket, London
shepherd's-purse
sibara
smartweed, Pennsylvania
sowthistle, annual
speedwell, purslane
telegraphplant
thistle, Russian

Scientific Name

Symphotrichum divaricatum
Ambrosia acanthicarpa
Soliva sessilis
Cyclosporum leptophyllum
Stellaria media
Trifolium repens
Gnaphalium purpureum
Amsinckia menziesii var. *intermedia*
Erodium cicutarium
Conyza bonariensis
Conyza ramosissima
Physalis angulata
Lamium amplexicaule
Conyza canadensis
Polygonum aviculare
Chenopodium album
Malva parviflora
Brassica juncea
Sinapis arvensis
Solanum nigrum
Lepidium virginicum
Amaranthus spp.
Matricaria discoidea
Plantago heterophylla
Portulaca oleracea
Raphanus raphanistrum
Ambrosia artemisiifolia
Sisymbrium irio
Capsella bursa-pastoris
Sibara virginica
Polygonum pensylvanicum
Sonchus oleraceus
Veronica peregrina
Heterotheca grandiflora
Salsola tragus

**Weeds controlled when applied at 23 fl oz per acre
(0.5 fl oz or 16 mis per 1000 sq ft):**

Common Name	Scientific Name
aster, heath	<i>Symphytichum ericoides</i>
bittercrass, little	<i>Cardamine oligosperma</i>
bittercrass, hairy	<i>Cardamine hirsuta</i>
brassbuttons, southern	<i>Cotula australis</i>
carrot, wild	<i>Daucus carota</i>
chamber-bitter	<i>Phyllanthus urinaria</i>
chickweed, mouseear	<i>Cerastium fontanum</i> ssp. <i>vulgare</i>
dandelion	<i>Taraxacum officinale</i>
eclipta	<i>Eclipta prostrata</i>
galinsoga, hairy	<i>Galinsoga quadriradiata</i>
geranium, Carolina	<i>Geranium carolinianum</i>
ladysthumb	<i>Polygonum persicaria</i>
lettuce, prickly	<i>Lactuca serriola</i>
mallow, common	<i>Malva neglecta</i>
marestail (see horseweed)	
mayweed, chamomile	<i>Anthemis cotula</i>
morningglory, ivyleaf	<i>Ipomoea hederacea</i>
mustard, black	<i>Brassica nigra</i>
pennywort	<i>Hydrocotyle</i> spp.
plantain, bracted	<i>Plantago aristata</i>
plantain, broadleaf	<i>Plantago major</i>
plantain, buckhorn	<i>Plantago lanceolata</i>
pokeweed, common	<i>Phytolacca americana</i>
rockpurslane, redmaids	<i>Calandrinia ciliata</i>
sida, prickly	<i>Sida spinosa</i>
sorrel, red	<i>Rumex acetosella</i>
speedwell, thymeleaf	<i>Veronica serpyllifolia</i>
spurge, hyssop	<i>Chamaesyce hyssopifolia</i>
spurge, spotted	<i>Chamaesyce maculata</i>
sweetclover, yellow	<i>Mellilotus officinalis</i>
tansymustard, green	<i>Descurainia pinnata</i> ssp. <i>brachycarpa</i>
woodsorrel, yellow	<i>Oxalis stricta</i>

**Weeds controlled when applied at 31 fl oz per acre
(0.7 fl oz or 21 mis per 1000 sq ft):**

Common Name	Scientific Name
burclover, California	<i>Medicago polymorpha</i>
dogfennel	<i>Eupatorium capillifolium</i>
eveningprimrose	<i>Oenothera</i> spp.
fescue, rattail	<i>Vulpia myuros</i>
filaree, whitestem	<i>Erodium moschatum</i>
goosefoot, nettleleaf	<i>Chenopodium murale</i>
groundsel, common	<i>Senecio vulgaris</i>
jimsonweed	<i>Datura stramonium</i>
knotweed, silversheath	<i>Polygonum argyrocoleon</i>
kochia	<i>Kochia scoparia</i>
medic, black	<i>Medicago lupulina</i>
mullein, turkey	<i>Croton setigerus</i>
nettle, burning	<i>Urtica urens</i>
oxtongue, bristly	<i>Picris echioides</i>
pimpernel, scarlet	<i>Anagallis arvensis</i>
sowthistle, spiny	<i>Sonchus asper</i>
spurge, petty	<i>Euphorbia pepulus</i>
spurge, prostrate	<i>Chamaesyce humistrata</i>
sunflower	<i>Helianthus</i> spp.
swinecress	<i>Coronopus didymus</i>
thistle, musk	<i>Carduus nutans</i>
willoweed, panicle	<i>Epilobium brachycarpum</i>
woodsorrel, creeping	<i>Oxalis corniculata</i>

**Weeds partially controlled or suppressed when applied at
31 fl oz per acre (0.7 fl oz or 21 mis per 1000 sq ft):**

Common Name	Scientific Name
bindweed, field	<i>Convolvulus arvensis</i>
carpetweed	<i>Mollugo verticillata</i>
dock, curly	<i>Rumex crispus</i>
mallow, Venice	<i>Hibiscus trionum</i>
milkweed, honeyvine	<i>Cynanchum laeve</i>
morningglory, tall	<i>Ipomoea purpurea</i>
pusley, Florida	<i>Richardia scabra</i>

Uses

Established Turfgrass

Use Gallery SC as a preemergence treatment for control of certain broadleaf weeds in established cool season and warm season turfgrass.

Apply Gallery SC anytime prior to germination of target weeds.

Note: Refer to the Product Information section of this label for use precautions and restrictions and information on mixing and application, application rates, and weeds controlled prior to using this product.

Tank Mixing

Gallery SC may be tank mixed with Dimension® herbicide and applied as a preemergence treatment to broaden the spectrum of annual grass and broadleaf weed control. Gallery SC may also be applied as a separate treatment to supplement the effectiveness of Team® 2G herbicide in cool and warm season turfgrass. Gallery SC may be tank mixed with postemergence broadleaf herbicides registered for use on established turfgrass to control existing broadleaf weeds to provide residual preemergence broadleaf weed control. Applied as directed, Gallery SC in tank mix with other products registered for use on turfgrass will provide control of susceptible weed species listed on the respective labels. When using Gallery SC in tank mix combinations with other products, read and follow all applicable use directions, precautions, and limitations on the respective product labels. Refer to tank mix instructions for Gallery SC in the Mixing Directions section.

Specific Use Precautions and Restrictions:

Apply Gallery SC to newly seeded turfgrass (including overseeded turfgrass) **only** after seedlings are established (three leaf stage and tillering) and well rooted. Do not overseed established turfgrass sooner than 60 days following an application of Gallery SC.

- Do not apply Gallery SC to golf course putting greens.
- Do not apply Gallery SC to dichondra.
- Do not apply Gallery SC to turfgrass grown for seed.

Use Gallery SC on the following turfgrass species:

Common Name	Scientific Name
Established Cool Season Turfgrass	
bentgrass, creeping	<i>Agrostis stolonifera</i>
bentgrass, colonial	<i>Agrostis tenuis</i>
bluegrass, Kentucky	<i>Poa pratensis</i>
fescue, chewing	<i>Festuca rubra</i> var. <i>commutata</i>
fescue, creeping red	<i>Festuca rubra</i>
fescue, sheeps	<i>Festuca ovina</i>
fescue, tall	<i>Festuca arundinaceae</i>
ryegrass, perennial	<i>Lolium perenne</i>

Established Warm Season Turfgrass¹

bahiagrass	<i>Paspalum notatum</i>
bermudagrass	<i>Cynodon dactylon</i>
buffalograss	<i>Buchloe dactyloides</i>
centipedegrass	<i>Eremochloa ophiuroides</i>
fescue, tall (growing in warm season areas)	<i>Festuca ovina</i>
St. Augustinegrass	<i>Stenotaphrum secundatum</i>
zoysiagrass	<i>Zoysia japonica</i>
zoysiagrass	<i>Zoysia tenuifolia</i>

¹Sprigged Warm Season Turfgrass: Use Gallery SC post-sprigging as a preemergence treatment for control of certain broadleaf weeds in warm season turfgrass. Apply anytime after sprigging in the following turfgrass species: bermudagrass, bahiagrass, St. Augustinegrass, centipedegrass and buffalograss. Do not apply more than 23 fl oz of Gallery SC per acre during the establishment phase for newly sprigged warm season turfgrass. Do not apply Gallery SC to varieties of dwarf-type bermudagrass or to any turfgrass species being sprigged on golf course tees or greens.

Ornamental Plantings

Use Gallery SC as a preemergence treatment for control of certain broadleaf weeds in landscape ornamentals, container grown ornamentals, field grown ornamentals, and groundcovers/perennials.

Apply Gallery SC anytime prior to germination of target weeds or immediately after cultivation.

Note: Refer to the Product Information section of this label for use precautions and restrictions and information on mixing and application, application rates, and weeds controlled prior to using this product.

Tank Mixing

Gallery SC may be tank mixed with Roundup or other postemergence herbicides registered for control of existing unwanted vegetation in ornamental plantings and to provide residual preemergence broadleaf weed control. Gallery SC may also be tank mixed with Dimension and applied preemergence to provide broad spectrum control of annual grasses and broadleaf weeds in ornamental areas. Applied as directed, tank mixes of Gallery SC will provide control of susceptible weed species listed

on the respective labels. When using Gallery SC in tank mix combination with other products, read and follow all applicable use directions, precautions, and limitations on the respective product labels. Refer to tank mix instructions for Gallery SC in the Mixing Directions section.

Note: Do not apply sprays containing Roundup over the top of ornamental plants. Extreme care must be exercised to prevent contact of sprays containing Roundup with foliage or stems of turfgrass, trees, shrubs, or other desirable vegetation since severe damage or death may result. If spraying Roundup in areas adjacent to desirable plants, use a shield to prevent spray from contacting foliage or stems of desirable plants.

Specific Use Precautions and Restrictions:

- Injury may be incurred if Gallery SC is applied in the following manner. Grower assumes all risk if Gallery SC is applied to:
- Nursery, forest, or Christmas tree: seedling beds, cutting beds, or transplant beds
 - Unrooted liners or cuttings that have been planted in pots for the first time
 - Pots less than six inches wide
 - Groundcovers until they are established and well rooted
 - Bedding plants or areas where bedding plants will be planted or transplanted within one year after application

Do not apply Gallery SC to newly transplanted ornamentals, nursery stock, groundcovers, or ornamental bulbs, until soil or potting media has been settled by packing and irrigation or rainfall and no cracks are present or plant injury may occur.

Applications of Gallery SC over the top of plants with newly forming buds may cause injury. Possible plant injury may be avoided by application as a directed spray to the soil surface beneath ornamental plants.

When planting into a site treated with Gallery SC in the past 8 months, use untreated soil as fill around roots when replacing plants or injury may occur.

Note: Injury to certain ornamental plants has been observed following application of Gallery SC. To avoid plant injury, do not use Gallery SC for weed control in the following ornamental plant species:

Scientific Name	Common Name
<i>Ajuga</i> spp.	bugleweed or ajuga
<i>Brassica</i> spp.	mustard
<i>Echinacea purpurea</i>	purple coneflower
<i>Euonymus alatus</i> 'Compacta'	dwarf burning bush
<i>Euphorbia</i> spp.	spurge
<i>Hydrangea</i> spp.	hydrangea
<i>Iberis</i> spp.	candytuft
<i>Juniperus horizontalis</i> 'Prince of Wales'	Prince of Wales juniper
<i>Melaleuca quinquenervia</i>	cajeput tree
<i>Rhododendron carolinianum</i>	Carolina rhododendron
<i>Rhododendron catawbiense</i> 'Roseum elegans'	roseum elegans
<i>Sedum</i> spp.	stonecrop
<i>Yucca recurvifolia</i>	green yucca

Gallery SC may be used in the culture of the following established plant species: (Note: Limitations on treatment methods)

Trees

Scientific Name	Common Name	Treatment Method ¹	Scientific Name	Common Name	Treatment Method ¹
<i>Abies balsamea</i>	balsam fir	C, F	<i>Elaeagnus angustifolia</i>	Russian olive	C, F
<i>Abies concolor</i>	white or concolor fir	F	<i>Elaeagnus x ebbengei</i>	gilt edge elaeagnus	C, F
<i>Abutilon hybridum</i>	white or concolor fir	F	'Gilt edge'		
	albus-flowering maple	C, F	<i>Eucalyptus camaldulensis</i>	red gum eucalyptus	F
	luteus-flowering maple	C, F	<i>Eucalyptus cinerea</i>	mealy eucalyptus	F
	roseus-flowering maple	C, F		silver dollar eucalyptus	F
	tangerine-flowering maple	C, F	<i>Eucalyptus microtheca</i>	coolibah tree	C, F
	maple		<i>Eucalyptus sideroxylon</i>	red ironbark eucalyptus	C, F
	vesuvius red-flowering maple	F	<i>Fagus sylvatica</i>	European beech	C, F
			<i>Ficus benamina</i>	fig	C, F
<i>Acer ginnala</i>	flame maple	F		mini fig	C, F
<i>Acer rubrum</i>	red maple	F	<i>Fraxinus udhei</i>	shamel ash	C, F
	red sunset maple	F	<i>Ginkgo biloba</i>	ginkgo (maidenhair tree)	F
<i>Acer saccharinum</i>	silver maple	C, F	<i>Gleditsia triacanthos var. inermis</i>	thornless honeylocust	F
<i>Acoelorrhapha whrightii</i>	Everglades palm	C, F			
<i>Albizia julibrissin</i>	silk tree	C, F	<i>Gleditsia triacanthos var. inermis</i>	shademaster honeylocust	F
<i>Alsophila australis</i>	Australian tree fern	C, F			
<i>Archontophoenix cunninghamiana</i>	king palm	C, F	<i>Heteromeles arbutiflora</i>	toyon	F
<i>Areacstrum romanzoffianum</i>	queen palm	C, F	<i>Illicium floridanum</i>	Florida anise-tree	C, F
<i>Araucaria heterophylla</i>	Norfolk island pine	C, F	<i>Juniperus virginiana</i>	eastern redcedar	C, F
<i>Bauhinia galpinii</i>	red bauhinia	C, F	<i>Leptospermum scoparium</i>	New Zealand tea tree	C, F
<i>Betula nigra</i>	river birch	C, F		ruby glow New Zealand tea tree	F
<i>Betula papyrifera</i>	paper birch	F	<i>Liquidambar styraciflua</i>	tea tree	F
<i>Betula pendula</i>	European white birch	C, F	<i>Magnolia grandiflora</i>	American sweetgum	F
<i>Brachychiton populneus</i>	bottle tree	C, F	<i>Magnolia soulangeana</i>	southern magnolia	C, F
<i>Bucida buceras</i>	black olive	F	<i>Magnolia stellata</i>	saucer magnolia	C, F
<i>Cerantonia siliqua</i>	carob	F	<i>Malus sargentii</i>	royal star magnolia	C, F
<i>Cercis canadensis</i>	redbud	C, F	<i>Morus alba</i>	crabapple non-bearing	C, F
<i>Chamaecyparis obtusa</i>	filicoides-fernspray cypress	F	<i>Musa aluminata</i>	white mulberry	F
	gracilis-slender hinoki cypress	F	<i>Oxydendrum arboreum</i>	banana	C, F
			<i>Picea abies</i>	sourwood	C, F
<i>Chamaecyparis pisifera</i>	sawara-false cypress	F		pendula-weeping Norway spruce	C, F
	squarrosa-moss cypress	F	<i>Picea glauca</i>	Norway spruce	C, F
<i>Chamaedorea cataractarum</i>	cat palm	F	<i>Picea glauca 'Conica'</i>	white spruce	C, F
<i>Chamaedorea costaricana</i>	palm	C, F	<i>Picea pungens</i>	dwarf alberta spruce	F
<i>Chamaedorea elegans</i>	parlor palm	C, F	<i>Picea pungens 'Glaucia'</i>	Colorado spruce	C, F
<i>Chamaerops humilis</i>	Mediterranean fan palm	C, F	<i>Picea pungens 'Hoopsia'</i>	Colorado blue spruce	C, F
<i>Cornus florida</i>	cloud nine dogwood	C, F	<i>Picea pungens 'Koster'</i>	hoopsi blue spruce	C, F
	flowering dogwood	C, F	<i>Pinus aristata</i>	koster blue spruce	F
<i>Cornus kousa</i>	kousa dogwood	C, F	<i>Pinus canariensis</i>	bristlecone pine	F
<i>Crataegus viridis</i>	green hawthorn	F	<i>Pinus contorta</i>	canary Island pine	F
<i>Cryptomeria japonica</i>	Japanese cryptomeria	C, F	<i>Pinus ularia</i>	shore pine, beach pine	F
<i>Cupaniopsis anacardioides</i>	carrot wood	F	<i>Pinus ularia</i>	eldarica pine	C, F
<i>Cupressus arizonicus or glabra</i>	Arizona cypress	F	<i>Pinus leucodermis</i>	Bosnian pine	C, F
<i>Cupressus ariz 'Blue Pyramid'</i>	blue pyramid cypress	C, F	<i>Pinus mugo var. pumilio</i>	pumilio mugo pine	C, F
<i>Cupressocyparis leylandii 'Emerald Isle'</i>	emerald isle leyland cypress	C, F	<i>Pinus nigra</i>	Austrian black pine	C, F
<i>Cupressocyparis leylandii 'Naylor's Blue'</i>	Naylor's blue leyland cypress	C, F	<i>Pinus radiata</i>	monterey pine	F
<i>Cupressus sempervirens</i>	Italian cypress	C, F	<i>Pinus strobus</i>	eastern white pine	C, F
<i>Cupressus sempervirens 'Glaucia'</i>	glauca Italian cypress	C, F		white pine	C, F
<i>Cycas revoluta</i>	sago palm	C, F	<i>Pinus sylvestris</i>	columnar Scotch pine	C, F
				Scotch pine	C, F
			<i>Pinus thunbergii</i>	Japanese black pine	C, F
			<i>Platanus occidentalis</i>	American sycamore	F
			<i>Platanus racemosa</i>	California sycamore	F
			<i>Podocarpus</i> spp.	podocarpus	F
			<i>Populus deltoides</i>	cottonwood	F
			<i>Prosopis chilensis</i>	Chilean mesquite	C, F

Trees (Cont.)

Scientific Name	Common Name	Treatment Method ¹	Scientific Name	Common Name	Treatment Method ¹
<i>Prunus yedoensis</i>	voshino flowering cherry	F	<i>Berberis thunbergii</i> var. <i>atropurpurea</i>	crimson pygmy barberry	C, F
<i>Prunus caroliniana</i>	Carolina laurel cherry	C, F	'Crimson Pygmy'		
	bright 'n tight Carolina laurel cherry	C, F	<i>Berberis thunbergii</i> var. <i>atropurpurea</i>	rose glow red barberry	C, F
<i>Prunus laurocerasus</i>	English laurel	C, F	'Rose Glow'		
<i>Quercus ilicifolia</i>	bear oak	F	<i>Berberis thunbergii</i> var. <i>atropurpurea</i>	redleaf Japanese barberry	F
<i>Quercus palustris</i>	pin oak	F	<i>Berberis thunbergii</i> var. <i>atropurpurea</i>	cherry bomb barberry	C, F
<i>Quercus phellos</i>	willow oak	C, F	'Cherry Bomb'		
<i>Quercus rubra</i>	red oak	C, F	<i>Bougainvillea</i> spp.	Barbara karst	C, F
<i>Quercus virginiana</i>	live oak	C, F		California gold	C, F
<i>Ravenea rivularis</i>	majesty palm	C, F		pink pixie	C, F
<i>Salix babylonica</i>	Babylon weeping willow	F		scarlet o'hara	C, F
<i>Salix matsudana</i> 'Torulosa'	corkscrew willow	F		temple fire	C, F
<i>Sequoiadendron giganteum</i>	giant sequoia	F		Texas dawn	C, F
<i>Sequoia sempervirens</i>	coast redwood	C, F	<i>Bougainvillea</i> 'Purple Queen'	purple queen	C, F
<i>Swietenia mahogany</i>	mahogany	F	<i>Bougainvillea</i> 'Rosenska'	rosenka bougainvillea	C, F
<i>Syagrus romanzoffianum</i>	queen palm	C, F	<i>Buxus microphylla</i> var. <i>japonica</i> 'Green Beauty'	green beauty boxwood	C, F
<i>Tabebuia caraiiba</i>	yellow tab	F	<i>Buxus microphylla</i> var. <i>japonica</i>	Japanese littleleaf boxwood	F
<i>Trachycarpus fortunei</i>	windmill palm	C, F	<i>Buxus microphylla</i> var. <i>Koreana</i>	Korean boxwood	F
<i>Tsuga canadensis</i>	eastern hemlock	C, F	<i>Buxus sempervirens</i>	common boxwood	C, F
<i>Ulmus parvifolia</i>	Chinese elm	F	<i>Buxus x 'Green velvet'</i>	green velvet boxwood	C, F
<i>Washingtonia robusta</i>	California fan palm	C, F	<i>Callistemon citrinus</i>	lemon bottlebrush	F
<i>Washingtonia robusta</i>	Mexican fan palm	F	<i>Callistemon citrinus</i>	little John lemon	C, F
<i>Zamia furluracea</i>	cardboard palm	C, F	'Little John'	bottlebrush	

¹C=container grown, F=field grown

Shrubs

Scientific Name	Common Name	Treatment Method ¹	Scientific Name	Common Name	Treatment Method ¹
<i>Abelia x grandiflora</i>	Edward goucher abelia	C, F	<i>Callistemon viminalis</i>	weeping bottlebrush	C, F
	glossy abelia	C, F	<i>Calluna vulgaris</i>	spring torch Scotch heather	C, F
	sunrise variegated abelia	C, F	<i>Camellia japonica</i>	camellia	C, F
<i>Acacia abyssinica</i>	abyssinica acacia	C, F	<i>Caryopteris clandonensis</i>	blue mist bluebeard	C, F
<i>Acacia redolens</i>	prostrate acacia	C, F	<i>Caryopteris x clandonen</i>	dark knight bluebeard	C, F
<i>Acacia stenophylla</i>	shoestring acacia	C, F	'Blk night'		
<i>Acalypha wilkesiana</i>	copper leaf	C, F	<i>Cassia artemisioides</i>	feathery cassia	C, F
<i>Acer ginnala</i>	amur maple	C, F	<i>Cassia, eremophila</i>	senna	C, F
<i>Acer palmatum</i>	coral bark Japanese maple	C, F	<i>Ceanothus</i> spp.	wild lilac	F
	maple		<i>Cephalotaxus harringtonia</i> var. <i>drupacae</i>	Japanese plum yew	C, F
<i>Acer palmatum</i>	dwarf Japanese maple	C, F	<i>Cerastium tomentosum</i>	snow-in-summer	C, F
<i>Agave americana</i>	century plant	F	<i>Ceratostigma</i>	dwarf plumbago	C, F
<i>Amelanchier alnifolia</i>	Saskatoon serviceberry	C, F	<i>plumbaginoides</i>		
<i>Andromeda polifolia</i>	bog rosemary	C, F	<i>Ceratostigma willmottianum</i>	Chinese plumbago	C, F
<i>Anisodonta</i>	cape mallow	C, F	<i>Chaenomeles japonica</i>	orange flowering quince	C, F
			<i>Chamaecyparis obtusa</i>	kosteri cypress	C, F
<i>hypomandarum</i>				nana-dwarf hinoki cypress	C, F
<i>Arctostaphylos uva-ursi</i>	bearberry	C, F		torulosa cypress	C, F
<i>Ardisia japonica</i>	chirimen marlberry	C, F	<i>Chamaecyparis pisifera</i>	baileyi-dogwood	F
<i>Armeria maritima</i>	bloodstone sea thrift	C, F		flaviramea-dogwood	F
'Bloodstone'				sawara-false cypress	F
<i>Artemesia lactiflora</i>	white mugwort	C, F		squarrosa minima	C, F
<i>Athyrium nipponicum</i>	Japanese painted fern	C, F		cypress	
<i>Aucuba japonica</i>	Japanese laurel			filifera-thread cypress	C, F
'Goldstrike'			<i>Chrysalidocarpus lutescens</i>	areca palm	F
<i>Baccharis pilularis</i>	coyotebush	F			
<i>Berberis x gladwynensis</i>	William Penn barberry	C, F			
<i>Berberis mentorensis</i>	mentor barberry	C, F			
<i>Berberis thunbergii</i> 'Aurea'	golden Japanese barberry	F			

Shrubs (Cont.)

Scientific Name	Common Name	Treatment Method ¹	Scientific Name	Common Name	Treatment Method ¹
<i>Clethra alnifolia</i>	summersweet	C, F	<i>Euonymus vegetus</i>	bigleaf wintercreeper	C, F
<i>Cleyera japonica</i>	Japanese cleyera	C, F	<i>Fatsia japonica</i>	Japanese aralia	C, F
<i>Colonema pulchrum</i>	pink breath of heaven	C, F	<i>Felicia amelloides</i>	blue marguerite	C, F
<i>Convolvus cneorum</i>	bush morning glory	C, F	<i>Forsythia x intermedia</i>	border forsythia	C, F
<i>Convolvulus mauritanicus</i>	ground morningglory	C, F	<i>Forsythia x 'Spring glory'</i>	spring glory forsythia	C, F
<i>Cornus alba</i>	sibirica-Siberian	C, F	<i>Fuchsia x "Santa Claus"</i>	Santa Claus fuchsia	C, F
	dogwood		<i>Gardenia jasminoides</i>	August beauty gardenia	C, F
<i>Cornus sericea</i>	baileyi redosier dogwood	F		gardenia	C, F
	flaviramea yellowtwig	F		miniature gardenia	C, F
	dogwood		<i>Gaultheria procumbens</i>	wintergreen	C, F
<i>Corylus americanus</i>	Harry Lauder's walking stick	C, F	<i>Gaultheria shallon</i>	sala/lemon leaf	C, F
"Contorta"			<i>Gelsemium sempervirens</i>	Carolina jessamine	C, F
<i>Cotinus coggygria</i>	royal purple smoke tree	C, F	<i>Genista pilosa</i>	woadwaxen	C, F
<i>Cotinus dammeri</i>	coral beauty smoke tree	C, F	<i>Hamamelis virginiana</i>	common witch hazel	C, F
	eichholz smoke tree	C, F	<i>Hardenbergia violacea</i>	lilac vine	C, F
<i>Cotoneaster adpressus</i>	praecox-early cotoneaster	C, F	<i>Hebe buxifolia</i>	boxleaf hebe	C, F
			<i>Hibiscus rosa-sinensis</i>	ross estey-hibiscus	C, F
<i>Cotoneaster apiculatus</i>	cranberry cotoneaster	C, F	<i>Hibiscus syriacus</i>	red bird rose of sharon	C, F
<i>Cotoneaster congestus</i>	Pyrenees cotoneaster	C, F		red heart rose of sharon	F
<i>Cotoneaster dammeri</i>	bearberry cotoneaster	C, F		woodbridge rose of sharon	C, F
<i>Cotoneaster himalayan</i>	Himalayan cotoneaster	C, F			
<i>Cotoneaster horizontalis</i>	rock cotoneaster	C, F	<i>Ilex aquifolium</i>	Balkans English holly	F
<i>Cycas revoluta</i>	sago palm	C, F		gold coast English holly	F
<i>Cyrtomium fortunei</i>	holly fern	C, F	<i>Ilex x aquipernyi</i>	San Jose holly	C, F
<i>Cytisus praecox</i>	hollandia-warminster broom	C, F	<i>Ilex x attenuata</i>	Savannah holly	C, F
			<i>Ilex cassine</i>	cassine holly	C, F
<i>Cytisus scoparius</i>	lena-Scotch broom	C, F	<i>Ilex cornuta</i>	burford holly	C, F
<i>Cytisus spp.</i>	holandia-Scotch broom	F		dwarf burford holly	C, F
	allgold warminster broom	C, F		needlepoint holly	C, F
			<i>Ilex crenata</i>	Chinese holly	C, F
	lilac time broom	C, F		compacta-dwarf	C, F
<i>Dalea greggii</i>	trailing indigo bush	C, F		Japanese holly	
<i>Daphne cneorum</i>	rose daphne	C, F		convexa holly	C, F
<i>Daphne odora</i>	fragrant daphne	C, F		dwarf Chinese holly	C, F
<i>Deutzia crenata</i>	nakiana-dwarf deutzia	C, F		green luster holly	C, F
<i>Deutzia gracilis</i>	slender gracilis	C, F		helleri-heller's	C, F
<i>Dodonea viscosa</i>	hopseed bush	F		Japanese holly	
<i>Enkianthus companulatus</i>	red-veined enkianthus	C, F		hetzii's Japanese holly	C, F
<i>Elaeagnus pungens</i>	fruitland silver berry	C, F	<i>Ilex crenata 'Steeds'</i>	steeds Japanese holly	C, F
<i>Erica cinerea</i>	purple bell heather	C, F		stokesii Japanese holly	C, F
<i>Erica vagans</i>	cornish heather	C, F	<i>Ilex glabra</i>	compacta-compact	C, F
<i>Erica x darleyensa</i>	Mediterranean pink heather	C, F		inkberry holly	
			<i>Ilex glabra</i>	nordica-inkberry holly	C, F
<i>Escalonia spp.</i>	escallonia	C, F	<i>Ilex x meserveae</i>	blue boy holly	C, F
<i>Eugenia myrtifolia</i>	teenie genie brushcherry	C, F		blue girl holly	C, F
<i>Eugenia myrtifolia 'Globulus'</i>	dwarf brush cherry	C, F		China boy holly	C, F
				China girl holly	C, F
<i>Euonymus fortunei</i>	canadale gold euonymus	C, F		ebony magic holly	F
	emerald 'n gold euonymus	F	<i>Ilex x 'Nellie Stevens'</i>	Nellie R. Stevens holly	C, F
			<i>Ilex opaca</i>	American holly	C, F
	sunspot euonymus	C, F	<i>Ilex vomitoria</i>	nana-dwarf yaupon holly	C, F
<i>Euonymus japonicus</i>	silver king euonymus	F		pendula-weeping	C, F
	chollipo euonymus	C, F		yaupon holly	
	gold spot euonymus	C, F		yaupon holly	C, F
	silver princess euonymus	C, F	<i>Illicium annisatum</i>	mystery gardenia	C, F
	variegated evergreen euonymus	C, F	<i>Itea virginica</i>	Henry's garnet	C, F
				sweetspire	
<i>Euonymus kiatschovicus</i>	spreading euonymus	C, F	<i>Ixora collinea</i>	ixora	C, F
<i>Euonymus kiatschovicus 'Manhattan'</i>	Manhattan euonymus	C, F	<i>Juniperus chinensis 'Gold Coast'</i>	gold coast juniper	C, F

Shrubs (Cont.)

Scientific Name	Common Name	Treatment Method ¹	Scientific Name	Common Name	Treatment Method ¹	
<i>Juniperus chinensis</i>	hollywood juniper	C, F	<i>Mahonia repens</i>	creeping mahonia	C, F	
	media-old gold juniper	C, F	<i>Mandevilla splendens</i>	red riding hood	F	
	pfitzer juniper	C, F	'Red Riding Hood'	mandevilla		
	pfitzerana glauca-blue juniper	C, F	<i>Michelia figo</i>	banana shrub	C, F	
	pfitzerana-pfitzer juniper	C, F	<i>Myrica cerifera</i>	southern waxmyrtle	C, F	
	sea green juniper	F	<i>Myrica pennsylvanica</i>	bayberry	C, F	
	torulosa-hollywood juniper	C, F	<i>Myoporum parvifolium</i>	putah creek	C, F	
			<i>Nandina domestica</i>	compacta-dwarf	C, F	
				heavenly bamboo		
	<i>Juniperus conferta</i>	emerald sea shore juniper	C, F		harbour dwarf-heavenly bamboo	C, F
	shore juniper	C, F		heavenly bamboo (nandina)	C, F	
<i>Juniperus davurica</i>	parsonii juniper	C, F		nana compacta-heavenly bamboo	C, F	
<i>Juniperus horizontalis</i>	andorra juniper	C, F		nana purpurea-heavenly bamboo	C, F	
	bar harbor juniper	C, F		woods dwarf-heavenly bamboo	C, F	
	blue chip juniper	C, F				
	blue rug juniper	C, F				
	creeping juniper	C, F	<i>Nerium oleander</i>	hardy red oleander	C, F	
	dwarf andorra juniper	C, F		oleander	C, F	
	huntington blue juniper	C, F		ruby lace oleander	C, F	
	plumosa-andorra juniper	C, F		fortune's osmanthus	C, F	
	wiltonii-blue carpet juniper	C, F	<i>Osmanthus x fortunei</i>	sweet olive osmanthus	C, F	
			<i>Osmathus fragrans</i>	purple fountain grass	C, F	
<i>Juniperus procumbens</i>	nana-dwarf Japanese garden juniper	C, F	<i>Pennisetum setaceum</i>	'Rubrum'		
	prostrata juniper	C, F				
<i>Juniperus prostrata</i>	broadmoor juniper	C, F	<i>Phoenix roebelenii</i>	pigmy date palm	C, F	
<i>Juniperus sabina</i>	foemina-hicks juniper	C, F	<i>Photinia x fraseri</i>	fraser photinia	C, F	
	savin juniper	C, F	<i>Physocarpus opulifolius</i>	dwarf ninebark	C, F	
		tamariscifolia-tam juniper	C, F	<i>Pieris japonica</i>	lily-of-the-valley	C, F
		emerald green juniper	F		mountain fire lily-of-the-valley	
		wichita blue juniper	C, F		snowdrift lily-of-the-valley	C, F
		blue juniper	C, F		temple bells lily-of-the-valley	C, F
		blue star juniper	C, F		valley rose lily-of-the-valley	C, F
		parsonii juniper	C, F		valley valentine lily-of-the-valley	C, F
	<i>Kalmia latifolia</i>	mountain laurel	C, F	<i>Pieris x 'Forest Flame'</i>	forest flame lily-of-the-valley	C, F
	<i>Lagerstroemia indica</i>	crape myrtle	C, F			
<i>Leucophyllum frutescens</i>	Texas sage	C, F	<i>Pinus mugo var. mugo</i>	mugo pine	C, F	
<i>Leucophyllum laevigatum</i>	chihuahan sage	C, F	<i>Pittosporum tenuifolia</i>	golf ball pittosporum	C, F	
<i>Leucothoe axillaris</i>	coast leucothoe	C, F	'Golf Ball'			
<i>Leucothoe fontanesiana</i>	drooping leucothoe	C, F	<i>Pittosporum tobira</i>	green pittosporum	C, F	
<i>Ligustrum japonicum</i>	Japanese privet	C, F		wheeler's dwarf pittosporum	C, F	
	wax privet	C, F				
	yellow tip privet	C, F	<i>Plumbago ariculata</i>	blue cape plumbago	F	
<i>Ligustrum lucidum</i>	glossy privet	C, F	<i>Plumbago capensis</i>	plumbago	C, F	
<i>Ligustrum ovalifolium</i>	California privet	F	<i>Podocarpus macrophyllus</i>	yewpine	C, F	
<i>Ligustrum texanum</i>	Howard privet	C, F	<i>Polygala fruticosa</i>	sweet pea shrub	C, F	
	wax leaf privet	C, F	<i>Polystichum polyblepharum</i>	tassel fern	C, F	
<i>Ligustrum x vicaryi</i>	golden vicary privet	F	<i>Potentilla fragiformis</i>	cinquefoil	F	
<i>Ligustrum vulgare</i>	lodense common privet	C, F	<i>Potentilla fruticosa</i>	cinquefoil	C, F	
				floppy disc cinquefoil	C, F	
<i>Livistona chinensis</i>	Chinese fountain palm	F		gold drop potentilla	F	
<i>Lonicera fragrantissima</i>	winter honeysuckle	C, F		goldfinger potentilla	C, F	
<i>Lonicera periclymenum</i>	flowering woodbine	C, F		red ace potentilla	C, F	
	serotina woodbine	C, F		sunset potentilla	C, F	
<i>Lonicera sempervirens</i>	trumpet honeysuckle	C, F		tangerine potentilla	C, F	
<i>Loropetalum chinense</i>	sizzling pink fringe flower	C, F		cinquefoil		
<i>Loropetalum chinense var. rubrum 'Razzelebern'</i>	razzeleberri fringe flower	C, F	<i>Potentilla spp.</i>			
<i>Mahonia aquifolium</i>	dwarf Oregon hollygrape	C, F				
'Compactum'	mahonia					
<i>Mahonia bealei</i>	leather leaf mahonia	C, F				

Shrubs (Cont.)

Scientific Name	Common Name	Treatment Method ¹	Scientific Name	Common Name	Treatment Method ¹
<i>Potentilla verna</i>	spring cinquefoil	C, F	<i>Rhododendron indica</i>	formosa azalea	C, F
<i>Prunus glandulosa</i>	dwarf flowering almond	C, F	<i>Rhododendron kaempferi</i>	waucabusa azalea	C, F
<i>Prunus laurocerasus</i>	otto luykens English laurel	C, F	<i>Rhododendron kerume</i>	blue danube azalea	C, F
'Otto luykens'	laurel			coral bells azalea	C, F
<i>Psidium cattleianum</i>	strawberry guava	C, F		hino crimson azalea	C, F
<i>Pyracantha coccinea</i>	lalalandei firethorn	C, F		hino pink azalea	C, F
'Lalalandei'				Mildred azalea	C, F
<i>Pyracantha fortuneana</i>	lolendei monrovia	C, F	<i>Rhododendron maximum</i>	snow azalea	C, F
	pyracantha		<i>Rhododendron mucronulatum</i>	rhodie max (rosebay)	C, F
	monon pyracantha	F		rhododendron	F
	red elf hybrid pyracantha	F	<i>Rhododendron obtusum</i>	hino crimsom azalea	C, F
	rutgers hybrid	C, F	<i>Rhododendron ponticum</i>	chioniods	C, F
	pyracantha			rhododendron	
	Santa Cruz pyracantha	C, F		daphnoides	C, F
	victory pyracantha	F		rhododendron	
<i>Rhaphiolepis indica</i>	charisma-monruce	C, F	<i>Rhododendron racemosum</i>	dwarf scarlet wonder	C, F
	rhaphiolepis			rhododendron	
	enchantress-moness	F		tribly rhododendron	C, F
	rhaphiolepis			unique rhododendron	C, F
	rhaphiolepis (India hawthorn)	C, F	<i>Rhododendron sassthigiatim x carolinianum</i>	vulcan rhododendron	C, F
	springtime-Monme	F		ramapo rhododendron	C, F
	rhaphiolepis				
<i>Rhaphiolepis indica</i>	ballerina Indian hawthorn	C, F	<i>Rhododendron satuski</i>	gumpo pink azalea	C, F
'Ballerina'				higasa azalea	F
<i>Rhaphiolepis ovata</i>	roundleaf rhaphiolepis	C, F		reijn azalea	C, F
<i>Rhododendron calendulaceum</i>	cannon's double azalea	C, F	<i>Rhododendron spp. hybrids</i>	carror azalea	C, F
	flame azalea	F		fashion azalea	C, F
	golden flare azalea	C, F		gerard Christina azalea	F
	Klondike azalea	C, F		girard Roberta azalea	C, F
<i>Rhododendron campylocarpum</i>	butterfly rhododendron	F		golden flare exbury azalea	F
<i>Rhododendron carolinianum x daurium</i>	PJM rhododendron	C, F		helmut vogel azalea	F
<i>Rhododendron catawbiense</i>	catawba album	C, F		hershey red azalea	F
	rhododendron			hot shot azalea	C, F
	catawba rhododendron	C, F		hume azalea	F
	Lord Roberts	C, F		Inga azalea	F
	rhododendron			Irene Koster azalea	C, F
	rocket rhododendron	C, F	<i>Rhus lancea</i>	President Clay azalea	C, F
<i>Rhododendron caucasicum x ponticum</i>	cunningham white	C, F	<i>Rhus typhina</i>	traditional azalea	C, F
<i>Rhododendron exbury</i>	rhododendron		<i>Rosa x 'Flower carpet'</i>	African sumac	C, F
	cannon's double azalea	C, F	<i>Rosa rugosa</i>	staghorn sumac	C, F
	golden flare azalea	C, F	<i>Rosmarinus officinalis</i>	red groundcover rose	C, F
	Klondike azalea	C, F	<i>Senecio cineraria</i>	ramanas rose	C, F
<i>Rhododendron forrestii repens</i>	gomer waterer	C, F	<i>Skimmia japonica</i>	rosemary	F
	rhododendron		<i>Skimmia revesiana</i>	dusty miller	
<i>Rhododendron forrestii x griersonianum</i>	Elizabeth rhododendron	C, F	<i>Solanum rantonetii 'Royal purple'</i>	Japanese skimmia	C, F
<i>Rhododendron griffithianum</i>	Jean Marie	C, F	<i>Spiraea x bumalda 'Anthony Waterer'</i>	reeve's skimmia	C, F
	rhododendron			Paraguay nightshade	C, F
<i>Rhododendron hybrid spp.</i>	America rhododendron	C, F	<i>Spiraea x cinerea 'Grefsheim'</i>	Anthony Waterer spiraea	C, F
	English roseum	F		first snow spiraea	C, F
	rhododendron		<i>Spiraea japonica</i>	dolchia spiraea	C, F
	nova zembla	C, F		Japanese alpine spiraea	C, F
	rhododendron			magic carpet spiraea	C, F
	scintillation	C, F		neon flash spiraea	C, F
	rhododendron			shirobana spiraea	C, F
<i>Rhododendron impeditum</i>	rhododendron	C, F		vanhouttei spirea	C, F

Shrubs (Cont.)

Scientific Name	Common Name	Treatment Method ¹	Scientific Name	Common Name	Treatment Method ¹
<i>Streptosolen jamesonii</i>	marmalade bush	C, F	<i>Agapanthus africanus</i>	lily of the Nile queen anne lily of the Nile	C, F C, F
<i>Syringa rothomagensis</i>	Chinese lilac	C, F	<i>Agapanthus 'Peter pan'</i>	lily of the Nile	C, F
<i>Syringa vulgaris</i>	common lilac	F	<i>Ammophila breviflulata</i>	beechgrass	C, F
<i>Taxus cuspidata</i>	Japanese yew	F	<i>Apтения cordifolia</i>	red apple aptenia	C, F
<i>Tecomania capensis</i>	cape honeysuckle	C, F	<i>Aquilegia x 'Dragon fly'</i>	columbine	C, F
<i>Ternstroemia gymnanthera</i>	Japanese ternstroemia	C, F	<i>Arctotheca calendula</i>	cape weed	F
<i>Teucrium fruticans</i>	bush germander	C, F	<i>Argyranthemum frutescens</i>	butterfly argyranthemum	C, F
<i>Thevetia nerifolia</i>	yellow oleander	C, F	'Butterfly'		
<i>Thuja occidentalis</i>	emerald arborvitae	C, F	<i>Asparagus densiflorus</i>	pony tail fern	C, F
	globe-globe arborvitae	C, F	'Myers'		
	little giant-dwarf arborvitae	C, F	<i>Asparagus retrofractus</i>		C, F
	arborvitae		<i>Asparagus varieiegata</i>	tree fern	C, F
	nigra-dark American arborvitae	C, F	<i>Asparagus var. 'Meegers'</i>		C, F
	pyramidalis arborvitae	C, F	<i>Aspidistra elatior</i>	cast iron plant	C, F
	rheingold arborvitae	C, F	<i>Aster novae-angliae</i>	New England aster	C, F
	techny arborvitae	F	<i>Aster novi-belgii</i>	New York aster	C, F
	woodwardii arborvitae	C, F	<i>Aster novi-belgii</i>	Persian rose dwarf aster	C, F
<i>Thuja orientalis</i>	areus nana-dwarf golden arborvitae	C, F	'Persian rose'		
	minima glauca-dwarf arborvitae	C, F	<i>Begonia sepmerflorens</i>	white ambassador begonia	C, F
	princes flower	C, F	'Amb white'		
<i>Tibouchina urvilleana</i>	Christmas palm	F	<i>Bergenia cordifolia</i>	heartleaf bergenia	C, F
<i>Veitchia merrillii</i>	pink dawn viburnum	C, F	<i>Bidens ferulifolia 'Peters gold'</i>	Peter's gold bidens	C, F
<i>Viburnum bodnantense</i>	Koreanspice viburnum	C, F	<i>Brachycome x 'New amethyst'</i>	swan river daisy new amethyst	C, F
<i>Viburnum carlesii</i>	David viburnum	C, F	<i>Callistephus chinensis</i>	China aster	C, F
<i>Viburnum davidii</i>	Japanese viburnum	F	<i>Carex alba</i>	frosty curls sedge	C, F
<i>Viburnum japonicum</i>	judd viburnum	C, F	<i>Carex spp.</i>	variegated carex	C, F
<i>Viburnum judd (V. x juddii)</i>	wayfaringtree viburnum	F	<i>Carpobrotus edulis</i>	largeleaf ice plant	F
<i>Viburnum lantana</i>	common snowball viburnum	F	<i>Catharanthus roseus</i>	Madagascar periwinkle	C, F
<i>Viburnum opulus sterile</i>	doublefile viburnum	C, F	<i>Chasmanthium latifolium</i>	northern sea oats	C, F
<i>Viburnum plicatum var. tomentosum</i>	tea viburnum	F	<i>Chrysanthemum maximum</i>	shasta daisy	C, F
<i>Viburnum setigerum</i>	spring bouquet viburnum	F	<i>Chrysanthemum spp.</i>	chrysanthemum species	C, F
<i>Viburnum tinus 'Compactum'</i>	Am. cranberrybush viburnum	C, F	<i>Clivia miniata 'French hybrid'</i>	kafr lily	C, F
<i>Viburnum trilobum</i>	dwarf Am. cranberrybush viburnum	C, F	<i>Cordylone indyisa</i>	blue dracaena	C, F
<i>Viburnum trilobum 'Compactum'</i>	Prague viburnum	C, F	<i>Coreopsis verticillata</i>	threadleaf coreopsis	C, F
<i>Viburnum x pragense</i>	bristol Ruby weigela	C, F	<i>Coreopsis verticillata 'Moonbeam'</i>	moonbeam coreopsis	C, F
<i>Weigela florida</i>	java red weigela	C, F	<i>Cortaderia selloana</i>	pampas grass	C, F
	minuet weigela	C, F	<i>Cuphea hyssopifolia</i>	false or Mexican heather	C, F
<i>Xylosma congestum</i>	xylosma	F	<i>Cyperus albobstriatus</i>	dwarf umbrella grass	C, F
<i>Xylosma senticoso</i>	shiny xylosma	F	<i>Dahlia hybrid Dwarf</i>	dwarf dahlia	C, F
<i>Yucca filamentosa</i>	Adam's needle yucca	C, F	<i>Dahlia x 'Royal Dahlietta pink'</i>	dwarf dahlia wendy pink	C, F
¹ C=container grown, F=field grown			<i>Delosperma alba</i>	white iceplant	F
			<i>Delosperma cooperi</i>	ice plant	C, F
			<i>Delosperma nubigenum</i>	hardy ice plant	C, F
			<i>Descampsia caespitosa</i>	descampsia	C, F
			<i>Dianthus gratianopolitanus</i>	firewitch cheddar pink	C, F
			'Firewitch'		
			<i>Dianthus gratianopolitanus</i>	crimson treasure	C, F
			'Treasure'	cheddar pink	
			<i>Dianthus plumaris</i>	cottage pink	C, F
			<i>Dietes vegeta</i>	fortnight lily	C, F
			<i>Drosanthemum floribundum</i>	trailing rosea iceplant	F
			<i>Drosanthemum hispidum</i>	iceplant	C, F
			<i>Dryopteris erythrosora</i>	autumn fern	C, F
			<i>Dryopteris marginalis</i>	marginal wood fern	C, F

Groundcovers/Perennials

Scientific Name	Common Name	Treatment Method ¹
<i>Achillea</i> spp.	yarrow	C, F
<i>Achillea filipendulina</i>	moonshine-fern/leaf yarrow	C, F
<i>Achillea millefolium</i>	common yarrow	C, F
<i>Achillea millefolium 'Paprika'</i>	paprika yarrow	C, F
<i>Achillea tomentosa</i>	wooly yarrow	C, F

Groundcovers/Perennials (Cont.)

Scientific Name	Common Name	Treatment Method ¹	Scientific Name	Common Name	Treatment Method ¹
<i>Dryopteris x australis</i>	dixie wood fern	C, F	<i>Leptospermum scoparium</i>	broom teatree/manuka	C, F
<i>Dymondia margaritae</i>	diamond marguerite	C, F	<i>Liatris spicata</i> 'Floristan Violet'	floristan violet gay feather	C, F
<i>Ensete ventricosum</i>	abyssinian banana	C, F	<i>Limonium perezii</i>	statice	C, F
<i>Equisetum scirpoides</i>	dwarf horsetail	C, F	<i>Liriope gigantea</i>	white lily turf	C, F
<i>Erianthus ravennae</i>	hardy pampasgrass	C, F		giant lily turf	C, F
<i>Erigeron speciosus</i>	darkest of all fleabane	C, F	<i>Liriope muscari</i>	lilac beauty lily turf	C, F
'Darkest of all'				majestic lily turf	C, F
<i>Euryops pectinatus</i>	dwarf euryops	C, F		monroe white lily turf	C, F
'Munchkin'				silvery sunproof lily turf	C, F
<i>Eustoma grandiflorum</i>	pink lisianthus	C, F		variegated liriope lily turf	C, F
'Pink'				big blue lily turf	C, F
<i>Evolvulus nuttallianus</i>	blue daze	C, F	<i>Liriope spicata</i>	green/creeping lily turf	C, F
<i>Fatsyhedra japonica</i>	Japanese aralia	C, F		silver dragon lily turf	C, F
<i>Festuca ovina glauca</i>	blue fescue	C, F	<i>Lonicera japonica</i>	Japanese honeysuckle	C, F
<i>Gaillardia x grandiflora</i>	blanket flower	C, F	<i>Lysimachia mummularia</i>	moneywort	C, F
<i>Gaillardia x grandiflora</i>	goblin blanket flower	F	<i>Lysimachia punctata</i>	dotted loosestrife	C, F
'Goblin'			<i>Matteuccia struthiopteris</i>	ostrich fern	C, F
<i>Gazania spp.</i>	gazania	C, F	<i>Matthiola incana</i>	stock	C, F
<i>Geranium cinerium</i>	ballerina cranesbill	C, F	'Harmony'		
'Ballerina'			<i>Miscanthus sinensis</i>	eulalia grass	C, F
<i>Geranium sanguineum</i>	bloody cranesbill	C, F	<i>Miscanthus sinensis</i>	maiden grass	C, F
'Bloody cran'			'Gracillimus'		
<i>Geranium subcaulescens</i>	black eyed magenta cranesbill	C, F	<i>Moraea iridioides</i>	African iris	C, F
			<i>Oenothera missouriensis</i>	ozark sundrops	C, F
<i>Geum spp</i>	avens	C, F	<i>Oenothera speciosa</i>	siskiyou evening primrose	C, F
<i>Geum quellyon</i>	geum	C, F	'Siskiyou pink'		
<i>Gypsophila paniculata</i>	baby's breath	C, F	<i>Onoclea sensibilis</i>	sensitive fern	C, F
<i>Hakonechloa macroaureola</i>	golden hakonechloa	C, F	<i>Ophiopogon japonicus</i>	dwarf mondo grass mondo grass	C, F
<i>Hedera canariensis</i>	Algerian ivy	F	<i>Origanum libanoticum</i>	oregano	C, F
<i>Hedera helix</i>	English ivy	C, F	<i>Osmunda cinnamomea</i>	cinnamon fern	C, F
<i>Helichrysum petiolare</i>	white licorice	C, F	<i>Osmunda regalis</i>	royal fern	C, F
'White licorice'	helichrysum		<i>Osteospermum fruticosum</i>	trailing African daisy	F
<i>Heliotropium fragrans</i>	common heliotrope	C, F	<i>Pachysandra terminalis</i>	Japanese spurge	C, F
<i>Hemerocallis spp.</i>	daylily	C, F	<i>Pachysandra terminalis</i>	green sheen Japanese spurge	C, F
<i>Hesperaloe parvifolia</i>	red yucca	C, F	'Green sheen'		
<i>Heuchera x 'Bressingham'</i>	bressingham coral bells	C, F	<i>Parthenocissus quinquefolia</i>	Virginia creeper	C, F
<i>Heuchera micrantha</i>	coral bells	C, F			
<i>Hosta 'Francee'</i>	francee plantain lily	C, F	<i>Pelargonium x hortorum</i>	zonal geranium	C, F
<i>Hosta fortunei</i>	plantain lily	C, F	<i>Pelargonium peltatum</i>	ivy geranium	C, F
<i>Hosta lancifolia</i>	albo-marginata hosta	C, F	<i>Pennisetum alopecuroides</i>	fountain grass	C, F
	narrow leafed plantain lily	C	<i>Pennisetum setaceum</i>	chrimson fountaingrass	C, F
			<i>Penstemon x 'Apple blossom'</i>	apple blossom penstemon	C, F
<i>Hosta x 'Patriot'</i>	patriot plantain lily	C, F	<i>Pentas lanceolata</i>	star clusters	C, F
<i>Houttuynia cordata</i>	chameleon houttuynia	C, F	<i>Perovskia atriplicifolia</i>	Russian sage	C, F
'Chameleon'			<i>Petunia-hybrids</i>	garden petunias	C, F
<i>Hymenoxys acaulis</i>	angelita daisy	C, F	<i>Phalaris arundinacea picta</i>	ribbon grass	C, F
<i>Hypericum spp.</i>	St. Johnswort	C, F	<i>Phlox subulata</i>	moss pink	C, F
<i>Impatiens walleryana</i>	lipstick impatiens	C, F	<i>Phormium tenax</i>	Jack Spratt New Zealand flax	C, F
'Lipstick'			'Jack Spratt'		
<i>Imperata cylindrical</i>	Japanese blood grass	C, F	<i>Polystichum acrostichoides</i>	Christmas fern	C, F
'Rubra'					
<i>Ipomea acuminata</i>	blue dawn morning glory	C, F	<i>Polystichum polyblepharum</i>	tassel fern	C, F
'Blue dawn'					
<i>Iris pumila 'Yellow'</i>	yellow dwarf bearded iris	C, F	<i>Ratibida columnifera</i>	Mexican hat	C, F
<i>Iris siberica</i>	iris	C, F	<i>Rudbeckia fulgida</i>	blackeyed susan	C, F
<i>Jasminum nitidum</i>	angelwing jasmine	C, F	<i>Ruscus hypophyllum</i>	butcher's broom (Israeli ruscus)	C, F
<i>Jasminum polyanthum</i>	pink jasmine	C, F			
<i>Kniphofia uvaria</i>	flamenco red hot poker	C, F	<i>Salvia daghestanica</i>	platinum sage	C, F
'Flamenco'			<i>Salvia grahamii</i>	graham's sage	C, F
<i>Lampranthus spectabilis</i>	trailing iceplant	F			

Groundcovers/Perennials (Cont.)

Scientific Name	Common Name	Treatment Method ¹
<i>Sasa pygmaea</i>	dwarf bamboo	C, F
<i>Scutellaria resinosa</i>	skull cap	C, F
<i>Sedum</i> x 'Autumn joy'	autumn joy stonecrop	C, F
<i>Sedum</i> x 'Vera Jameson'	Vera Jameson stonecrop	C, F
<i>Tagetes patula</i> 'Little Hero'	little hero marigold	C, F
<i>Trachelospermum asiaticum</i>	Asian jasmine	C, F
<i>Tulbaghia violacea</i>	society garlic	C, F
<i>Verbena rigida</i>	veined verbena	C, F
<i>Vinca major</i>	bigleaf periwinkle	C, F
<i>Vinca minor</i>	dwarf periwinkle	F
<i>Vinca</i> spp.	periwinkle	F

¹C=container grown, F=field grown

Ornamental Bulbs in Landscape Settings

Gallery SC may be applied for control of susceptible annual weeds in ornamental bulbs such as bulbous iris, daffodil (narcissus), gladiolus, hyacinth, lilies, and tulip. Apply Gallery SC to the soil surface 2 to 4 weeks after planting but prior to the emergence of annual weeds. Gallery SC may also be applied following bulb emergence but prior to flowering and after flowering. For fall planted bulbs, apply Gallery SC in late winter or early spring to weed-free soil surfaces. For bulbs, make a single application within 30 days following planting and prior to bulb emergence. Do not exceed the 16 fl. oz of Gallery SC (0.5 lb. ai) per acre rate.

Specific Use Precautions and Restrictions:

- Do not use Gallery SC for weed control in ornamental bulbs grown for commercial production.
- Gallery SC is not for application to:
 - Tulip plants that have emerged to a height greater than 3/4 inch.
 - Gladiolus prior to emergence or if corms are less than one inch in diameter.
 - Bulbs while they are flowering.

Shadehouse Areas

Gallery SC may be applied in open shadehouse-type structures where the natural flow of air is unimpeded. Do not apply in enclosed greenhouses or in enclosed shadehouse-type structures. Do not apply within three weeks prior to enclosing greenhouses or poly-type structures.

Christmas Tree and Conifer Plantations

Gallery SC - Alone

Apply Gallery SC as a directed spray to the soil surface or as an over the top spray to established plantings of field grown Christmas tree and conifer species listed in this label. Follow all instructions provided in the Product Information section of this label.

Specific Use Precautions and Restrictions:

Do not apply Gallery SC to seedbeds or seedling transplant beds. Apply only to established plantings. Established plants are defined as those that have been transplanted into their final growing location for a sufficient period of time to allow the soil to be firmly settled around the roots from packing and rainfall or irrigation.

Gallery SC - Tank Mix

Tank mix combinations of Gallery SC plus other labeled herbicides may be used in established Christmas tree plantings. When applied according to use directions, these tank mixes will provide control of susceptible weed species listed on the respective product labels. Refer to tank mix product labels for specific use directions, precautions and limitations before use. Refer to tank mix instructions for Gallery SC in the Mixing Directions section.

Gallery SC plus Roundup: Apply tank mix combinations of Gallery SC plus Roundup as directed soil sprays only in Christmas tree plantings. When applied as directed, Gallery SC plus Roundup will provide postemergence control of susceptible weed species listed on the label for Roundup and residual preemergence control of susceptible weed species listed on the label for Gallery SC. Refer to the label for Roundup for specific use directions, precautions and limitations before use. Refer to tank mix instructions for Gallery SC in the Mixing Directions section.

Specific Use Precautions and Restrictions:

- Do not apply sprays containing Roundup over the top of Christmas tree plantings.
- Extreme care must be exercised to prevent contact of sprays containing Roundup with foliage or stems of Christmas trees or other desirable plants or severe plant damage or death may result.

Non-Cropland Areas

Use Gallery SC as a preemergence herbicide for control of listed broadleaf weeds in non-cropland areas including industrial sites, utility substations, parking lots, driveways, walking paths, highway guardrails, sign posts and delineators where maintenance of bare ground is desired.

Apply Gallery SC anytime prior to germination of target weeds. Areas to be treated should be free of established weeds or existing weeds should be controlled with postemergence herbicides.

Refer to the Product Information section prior to using this product on non-cropland areas.

Tank Mixing

Gallery SC is compatible and can be tank mixed with other herbicides registered for use on non-cropland areas. Applied as directed, tank mixes containing Gallery SC will provide control of susceptible weed species listed on the respective labels. All directions, precautions and limitations on the respective product labels apply to the tank mix use. Refer to tank mix instructions for Gallery SC in the Mixing Directions section.

Terms and Conditions of Use

If terms of the following Warranty Disclaimer, Inherent Risks of Use and Limitation of Remedies are not acceptable, return unopened package at once to the seller for a full refund of purchase price paid. To the extent permitted by law, otherwise, use by the buyer or any other user constitutes acceptance of the terms under Warranty Disclaimer, Inherent Risks of Use and Limitation of Remedies.

Warranty Disclaimer

Dow AgroSciences warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions, subject to the inherent risks set forth below. To the extent permitted by law, Dow AgroSciences MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.

Inherent Risks of Use

It is impossible to eliminate all risks associated with use of this product. Crop injury, lack of performance, or other unintended consequences may result because of such factors as use of the product contrary to label instructions (including conditions noted on the label, such as unfavorable temperatures, soil conditions, etc.), abnormal conditions (such as excessive rainfall, drought, tornadoes, hurricanes), presence of other materials, the

manner of application, or other factors, all of which are beyond the control of Dow AgroSciences or the seller. To the extent permitted by law, all such risks shall be assumed by buyer.

Limitation of Remedies

To the extent permitted by law, the exclusive remedy for losses or damages resulting from this product (including claims based on contract, negligence, strict liability, or other legal theories), shall be limited to, at Dow AgroSciences' election, one of the following:

1. Refund of purchase price paid by buyer or user for product bought, or
2. Replacement of amount of product used.

To the extent permitted by law, Dow AgroSciences shall not be liable for losses or damages resulting from handling or use of this product unless Dow AgroSciences is promptly notified of such loss or damage in writing. To the extent permitted by law, in no case shall Dow AgroSciences be liable for consequential or incidental damages or losses.

The terms of the Warranty Disclaimer, Inherent Risks of Use and this Limitation of Remedies cannot be varied by any written or verbal statements or agreements. No employee or sales agent of Dow AgroSciences or the seller is authorized to vary or exceed the terms of the Warranty Disclaimer or this Limitation of Remedies in any manner.

EPA accepted: 08/22/13

®Trademark of The Dow Chemical Company ("Dow") or an affiliated company of Dow



Dow AgroSciences

Gallery[®] SC

SPECIALTY HERBICIDE

A preemergence herbicide for control of certain broadleaf weeds in:

- Established Turfgrass
- Landscape Ornamentals
- Container Grown Ornamentals
- Field Grown Ornamentals
- Groundcovers/ Perennials
- Non-Cropland Ornamental Bulbs
- Christmas Tree/Conifer Plantations

Group	21	HERBICIDE
-------	----	-----------

Active Ingredient:

isoxaben: N-[3-(1-ethyl-1-methylpropyl)-5-isoxazolyl]-2,6-dimethoxybenzamide and isomers 45.45%

Other Ingredients 54.55%

Total 100%

Contains 4.16 lb active ingredient per gallon.

Keep Out of Reach of Children CAUTION

Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. Refer to the label booklet under "Agricultural Use Requirements" in the Directions for Use section for information about this standard.

For additional Precautionary Statements, First Aid, Storage and Disposal and other use information see inside this label.

Notice: Read the entire label. Use only according to label directions. **Before using this product, read Warranty Disclaimer, Inherent Risks of Use and Limitation of Remedies at end of label booklet. If terms are unacceptable, return at once unopened.**

In case of emergency endangering health or the environment involving this product, call 1-800-992-5994.

Agricultural Chemical: Do not ship or store with food, feeds, drugs or clothing.

Shake Well Before Use – Avoid Freezing

EPA Reg. No. 62719-658

900-021178 / 11079455

®Trademark of The Dow Chemical Company ("Dow") or an affiliated company of Dow

Produced for
Dow AgroSciences LLC
 9330 Zionsville Road
 Indianapolis, IN 46268

NET CONTENTS 2.5 GAL

PEEL FILM HERE



(01) 0 06 62974 71074 5

51 1/4"

1 7/16"

14 15/16"

9 15/16"

15 1/16"

9 7/8"

Do Not Cut To Open
Contents In Plastic

SHAKE WELL



BEFORE USE

4 7/8"



Gallery[®] SC

SPECIALTY HERBICIDE

A preemergence herbicide for control of certain broadleaf weeds in:

- Established Turfgrass
- Landscapes
- Ornamentals
- Container Grown Ornamentals
- Field Grown Ornamentals
- Groundcovers/Perennials
- Non-Cropland
- Ornamental Bulbs
- Christmas Tree/Conifer Plantations

Notice: Read the entire label. Use only according to label directions. **Before using this product, read Warranty Disclaimer, Inherent Risks of Use and Limitation of Remedies at end of label booklet. If terms are unacceptable, return at once unopened.**

In case of emergency endangering health or the environment involving this product, call 1-800-992-5994.

Agricultural Chemical: Do not ship or store with food, feeds, drugs or clothing.

Shake Well Before Use – Avoid Freezing

EPA Reg. No. 62719-658 EPA Est. 11773-IA-001

Group	21	HERBICIDE
Active Ingredient:		
isoxaben: N-[3-(1-ethyl-1-methylpropyl)-5-isoxazoly]-2,6-dimethoxybenzamide and isomers	45.45%	
Other Ingredients	54.55%	
Total	100%	

Contains 4.16 lb active ingredient per gallon.

Keep Out of Reach of Children

CAUTION


Agricultural Use Requirements
Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. Refer to the label booklet under "Agricultural Use Requirements" in the Directions for Use section for information about this standard.

Refer to inside of label booklet for additional precautionary information including Directions for Use.

®Trademark of The Dow Chemical Company ("Dow") or an affiliated company of Dow

Produced for
Dow AgroSciences LLC
9330 Zionsville Road
Indianapolis, IN 46268

11079206



11079206



25 1/16"
CORR.

15 5/16"



Gallery[®]

SC

SPECIALTY HERBICIDE

2 X 2.5 GAL



Gallery[®]

SC

SPECIALTY HERBICIDE

2 X 2.5 GAL



Gallery[®]

SC

SPECIALTY HERBICIDE

2 X 2.5 GAL

900-021179 / 11079456
BOX MFG. DATE:

4 7/8"