



TR-10

GRANULAR HERBICIDE

A selective herbicide for the preemergence control of annual grasses and broadleaf weeds in **Alfalfa, Almond, Apricot, Asparagus, Barley, Beans – All Dry and Fresh Beans/Peas, Broccoli, Brussels Sprouts, Cabbage, Carrot, Castor Oil Plant, Cauliflower, Celery, Chicory, Collard Greens, Corn, Cotton, Crambe, Cucurbits, Dry Peas, Durum, English Peas, Flax, Flaxseed, Grain Sorghum, Grapes, Grapefruit, Guar, Hops, Kale, Kenaf, Lemon, Lima Bean, Mungbean, Mustard Greens, Nectarine, Niger seed, Oil radish, Okra, Onions, Orange, Peach, Peanut, Pecan, Pepper, Plum, Potatoes, Prune, Radish, Radish, Raspberry, Snap Bean, Sesame Peas, Soybeans, Sugar Beets, Sugarcane, Sunflowers, Tangelo, Tangerine, Tomatoes, Turpin Greens, Walnut, and Wheat**.

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Wash and wash PPE separately from other laundry.

Engineering Controls
Wear eyewear with eye protection, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d)-(6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

Refer to inside of label booklet for additional precautionary information including Directions for Use.
EPA Reg. No. 62719-131 900-021310 / 00013988
*Trademark of The Dow Chemical Company ("Dow") or an affiliated company of Dow

Produced for Dow AgroSciences LLC 8330 Zionsville Road Indianapolis, IN 46268

User Safety Recommendations
Users should:
• Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
• Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
• Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

First Aid
If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-992-5994 for emergency medical treatment information.

Environmental Hazards
This pesticide is extremely toxic to freshwater marine, and estuarine fish and aquatic invertebrates including shrimp and oyster. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not apply in a manner which will directly expose canals, lakes, streams, ponds, marshes or estuaries to aerial drift. Do not contaminate water when disposing of equipment washwaters.

Agricultural Use Requirements
Use this product in accordance with the label and labeling with the Worker Protection Standard, 40 CFR Part 170. Refer to label booklet under "Agricultural Use Requirements" in the Directions for Use section for information about this standard.

Directions for Use
It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Read all Directions for Use carefully before applying. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. Read all requirements specific to your state or tribe, consult the agency responsible for pesticide registration.

Agricultural Use Requirements (Cont.)
Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains

Precautionary Statements
Hazards to Humans and Domestic Animals
CAUTION
Causes Eye Irritation • Prolonged or Frequently Repeated Skin Contact May Cause Allergic Reactions In Some Individuals

Personal Protective Equipment (PPE)
Applicators and other handlers must wear:
• Long-sleeved shirt and long pants
• Chemical-resistant gloves, such as butyl rubber ≥1 mils, or nitrile rubber ≥14 mils, or neoprene rubber ≥14 mils; or nitrile rubber ≥14 mils
• Shoes plus socks
Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Wash and wash PPE separately from other laundry.

Storage and Disposal
Do not contaminate water, food or feed by storage or disposal.
Pesticide Storage: Store in original container only. In case of spill, contain material and dispose as waste.
Pesticide Disposal: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.
Nonrefillable rigid containers 5 gallons or less:
Container Handling: Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures allowed by state and local authorities.
Triple rinse or pressure rinse container (or equivalent) promptly after emptying. **Triple rinse** as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. **Pressure rinse** as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 psi for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

Nonrefillable rigid containers larger than 5 gal:
Container Handling: Refillable container. Refill this container with pesticide only. Do not reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Clean before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from this container into application equipment or a mix tank. Fill the container about 10% full with water and, if possible, add sides while adding water. If practical, agitate vigorously or recirculate water with the pump for two minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times.

Container Handling: Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures allowed by state and local authorities.
Triple rinse or pressure rinse container (or equivalent) promptly after emptying. **Triple rinse** as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. **Pressure rinse** as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 psi for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

Application
Apply Treflan TR-10 to soils that are wet or are subject to prolonged periods of flooding as poor weed control may result.
Do not use Treflan TR-10 on any crop grown in Pecos county or Reeves county, Texas.
In Montana, uses of Treflan TR-10 are limited to those described in supplemental labeling. Refer to supplemental labeling for crops and specific use directions.

Application
Apply Treflan TR-10 to soils that are wet or are subject to prolonged periods of flooding as poor weed control may result.
Do not use Treflan TR-10 on any crop grown in Pecos county or Reeves county, Texas.
In Montana, uses of Treflan TR-10 are limited to those described in supplemental labeling. Refer to supplemental labeling for crops and specific use directions.

Application
Apply Treflan TR-10 to soils that are wet or are subject to prolonged periods of flooding as poor weed control may result.
Do not use Treflan TR-10 on any crop grown in Pecos county or Reeves county, Texas.
In Montana, uses of Treflan TR-10 are limited to those described in supplemental labeling. Refer to supplemental labeling for crops and specific use directions.

Application
Apply Treflan TR-10 to soils that are wet or are subject to prolonged periods of flooding as poor weed control may result.
Do not use Treflan TR-10 on any crop grown in Pecos county or Reeves county, Texas.
In Montana, uses of Treflan TR-10 are limited to those described in supplemental labeling. Refer to supplemental labeling for crops and specific use directions.

Application
Apply Treflan TR-10 to soils that are wet or are subject to prolonged periods of flooding as poor weed control may result.
Do not use Treflan TR-10 on any crop grown in Pecos county or Reeves county, Texas.
In Montana, uses of Treflan TR-10 are limited to those described in supplemental labeling. Refer to supplemental labeling for crops and specific use directions.

Application
Apply Treflan TR-10 to soils that are wet or are subject to prolonged periods of flooding as poor weed control may result.
Do not use Treflan TR-10 on any crop grown in Pecos county or Reeves county, Texas.
In Montana, uses of Treflan TR-10 are limited to those described in supplemental labeling. Refer to supplemental labeling for crops and specific use directions.

Storage and Disposal (Cont.)
the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring all fresh contents are completely incorporated. Empty remaining contents into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. **Pressure** rise as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 psi for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

Application
Apply Treflan TR-10 to soils that are wet or are subject to prolonged periods of flooding as poor weed control may result.
Do not use Treflan TR-10 on any crop grown in Pecos county or Reeves county, Texas.
In Montana, uses of Treflan TR-10 are limited to those described in supplemental labeling. Refer to supplemental labeling for crops and specific use directions.

Application
Apply Treflan TR-10 to soils that are wet or are subject to prolonged periods of flooding as poor weed control may result.
Do not use Treflan TR-10 on any crop grown in Pecos county or Reeves county, Texas.
In Montana, uses of Treflan TR-10 are limited to those described in supplemental labeling. Refer to supplemental labeling for crops and specific use directions.

Application
Apply Treflan TR-10 to soils that are wet or are subject to prolonged periods of flooding as poor weed control may result.
Do not use Treflan TR-10 on any crop grown in Pecos county or Reeves county, Texas.
In Montana, uses of Treflan TR-10 are limited to those described in supplemental labeling. Refer to supplemental labeling for crops and specific use directions.

Application
Apply Treflan TR-10 to soils that are wet or are subject to prolonged periods of flooding as poor weed control may result.
Do not use Treflan TR-10 on any crop grown in Pecos county or Reeves county, Texas.
In Montana, uses of Treflan TR-10 are limited to those described in supplemental labeling. Refer to supplemental labeling for crops and specific use directions.

Application
Apply Treflan TR-10 to soils that are wet or are subject to prolonged periods of flooding as poor weed control may result.
Do not use Treflan TR-10 on any crop grown in Pecos county or Reeves county, Texas.
In Montana, uses of Treflan TR-10 are limited to those described in supplemental labeling. Refer to supplemental labeling for crops and specific use directions.

Application
Apply Treflan TR-10 to soils that are wet or are subject to prolonged periods of flooding as poor weed control may result.
Do not use Treflan TR-10 on any crop grown in Pecos county or Reeves county, Texas.
In Montana, uses of Treflan TR-10 are limited to those described in supplemental labeling. Refer to supplemental labeling for crops and specific use directions.

Application
Apply Treflan TR-10 to soils that are wet or are subject to prolonged periods of flooding as poor weed control may result.
Do not use Treflan TR-10 on any crop grown in Pecos county or Reeves county, Texas.
In Montana, uses of Treflan TR-10 are limited to those described in supplemental labeling. Refer to supplemental labeling for crops and specific use directions.

Application
Apply Treflan TR-10 to soils that are wet or are subject to prolonged periods of flooding as poor weed control may result.
Do not use Treflan TR-10 on any crop grown in Pecos county or Reeves county, Texas.
In Montana, uses of Treflan TR-10 are limited to those described in supplemental labeling. Refer to supplemental labeling for crops and specific use directions.

Application
Apply Treflan TR-10 to soils that are wet or are subject to prolonged periods of flooding as poor weed control may result.
Do not use Treflan TR-10 on any crop grown in Pecos county or Reeves county, Texas.
In Montana, uses of Treflan TR-10 are limited to those described in supplemental labeling. Refer to supplemental labeling for crops and specific use directions.

Application
Apply Treflan TR-10 to soils that are wet or are subject to prolonged periods of flooding as poor weed control may result.
Do not use Treflan TR-10 on any crop grown in Pecos county or Reeves county, Texas.
In Montana, uses of Treflan TR-10 are limited to those described in supplemental labeling. Refer to supplemental labeling for crops and specific use directions.

Application
Apply Treflan TR-10 to soils that are wet or are subject to prolonged periods of flooding as poor weed control may result.
Do not use Treflan TR-10 on any crop grown in Pecos county or Reeves county, Texas.
In Montana, uses of Treflan TR-10 are limited to those described in supplemental labeling. Refer to supplemental labeling for crops and specific use directions.

Application
Apply Treflan TR-10 to soils that are wet or are subject to prolonged periods of flooding as poor weed control may result.
Do not use Treflan TR-10 on any crop grown in Pecos county or Reeves county, Texas.
In Montana, uses of Treflan TR-10 are limited to those described in supplemental labeling. Refer to supplemental labeling for crops and specific use directions.

Application
Apply Treflan TR-10 to soils that are wet or are subject to prolonged periods of flooding as poor weed control may result.
Do not use Treflan TR-10 on any crop grown in Pecos county or Reeves county, Texas.
In Montana, uses of Treflan TR-10 are limited to those described in supplemental labeling. Refer to supplemental labeling for crops and specific use directions.

Application
Apply Treflan TR-10 to soils that are wet or are subject to prolonged periods of flooding as poor weed control may result.
Do not use Treflan TR-10 on any crop grown in Pecos county or Reeves county, Texas.
In Montana, uses of Treflan TR-10 are limited to those described in supplemental labeling. Refer to supplemental labeling for crops and specific use directions.

Application
Apply Treflan TR-10 to soils that are wet or are subject to prolonged periods of flooding as poor weed control may result.
Do not use Treflan TR-10 on any crop grown in Pecos county or Reeves county, Texas.
In Montana, uses of Treflan TR-10 are limited to those described in supplemental labeling. Refer to supplemental labeling for crops and specific use directions.

Application
Apply Treflan TR-10 to soils that are wet or are subject to prolonged periods of flooding as poor weed control may result.
Do not use Treflan TR-10 on any crop grown in Pecos county or Reeves county, Texas.
In Montana, uses of Treflan TR-10 are limited to those described in supplemental labeling. Refer to supplemental labeling for crops and specific use directions.

Application
Apply Treflan TR-10 to soils that are wet or are subject to prolonged periods of flooding as poor weed control may result.
Do not use Treflan TR-10 on any crop grown in Pecos county or Reeves county, Texas.
In Montana, uses of Treflan TR-10 are limited to those described in supplemental labeling. Refer to supplemental labeling for crops and specific use directions.

Application
Apply Treflan TR-10 to soils that are wet or are subject to prolonged periods of flooding as poor weed control may result.
Do not use Treflan TR-10 on any crop grown in Pecos county or Reeves county, Texas.
In Montana, uses of Treflan TR-10 are limited to those described in supplemental labeling. Refer to supplemental labeling for crops and specific use directions.

Application
Apply Treflan TR-10 to soils that are wet or are subject to prolonged periods of flooding as poor weed control may result.
Do not use Treflan TR-10 on any crop grown in Pecos county or Reeves county, Texas.
In Montana, uses of Treflan TR-10 are limited to those described in supplemental labeling. Refer to supplemental labeling for crops and specific use directions.

Application
Apply Treflan TR-10 to soils that are wet or are subject to prolonged periods of flooding as poor weed control may result.
Do not use Treflan TR-10 on any crop grown in Pecos county or Reeves county, Texas.
In Montana, uses of Treflan TR-10 are limited to those described in supplemental labeling. Refer to supplemental labeling for crops and specific use directions.

Application
Apply Treflan TR-10 to soils that are wet or are subject to prolonged periods of flooding as poor weed control may result.
Do not use Treflan TR-10 on any crop grown in Pecos county or Reeves county, Texas.
In Montana, uses of Treflan TR-10 are limited to those described in supplemental labeling. Refer to supplemental labeling for crops and specific use directions.

Application
Apply Treflan TR-10 to soils that are wet or are subject to prolonged periods of flooding as poor weed control may result.
Do not use Treflan TR-10 on any crop grown in Pecos county or Reeves county, Texas.
In Montana, uses of Treflan TR-10 are limited to those described in supplemental labeling. Refer to supplemental labeling for crops and specific use directions.

Application
Apply Treflan TR-10 to soils that are wet or are subject to prolonged periods of flooding as poor weed control may result.
Do not use Treflan TR-10 on any crop grown in Pecos county or Reeves county, Texas.
In Montana, uses of Treflan TR-10 are limited to those described in supplemental labeling. Refer to supplemental labeling for crops and specific use directions.

Application
Apply Treflan TR-10 to soils that are wet or are subject to prolonged periods of flooding as poor weed control may result.
Do not use Treflan TR-10 on any crop grown in Pecos county or Reeves county, Texas.
In Montana, uses of Treflan TR-10 are limited to those described in supplemental labeling. Refer to supplemental labeling for crops and specific use directions.

Application
Apply Treflan TR-10 to soils that are wet or are subject to prolonged periods of flooding as poor weed control may result.
Do not use Treflan TR-10 on any crop grown in Pecos county or Reeves county, Texas.
In Montana, uses of Treflan TR-10 are limited to those described in supplemental labeling. Refer to supplemental labeling for crops and specific use directions.

Application
Apply Treflan TR-10 to soils that are wet or are subject to prolonged periods of flooding as poor weed control may result.
Do not use Treflan TR-10 on any crop grown in Pecos county or Reeves county, Texas.
In Montana, uses of Treflan TR-10 are limited to those described in supplemental labeling. Refer to supplemental labeling for crops and specific use directions.

Application
Apply Treflan TR-10 to soils that are wet or are subject to prolonged periods of flooding as poor weed control may result.
Do not use Treflan TR-10 on any crop grown in Pecos county or Reeves county, Texas.
In Montana, uses of Treflan TR-10 are limited to those described in supplemental labeling. Refer to supplemental labeling for crops and specific use directions.

Application
Apply Treflan TR-10 to soils that are wet or are subject to prolonged periods of flooding as poor weed control may result.
Do not use Treflan TR-10 on any crop grown in Pecos county or Reeves county, Texas.
In Montana, uses of Treflan TR-10 are limited to those described in supplemental labeling. Refer to supplemental labeling for crops and specific use directions.

Application
Apply Treflan TR-10 to soils that are wet or are subject to prolonged periods of flooding as poor weed control may result.
Do not use Treflan TR-10 on any crop grown in Pecos county or Reeves county, Texas.
In Montana, uses of Treflan TR-10 are limited to those described in supplemental labeling. Refer to supplemental labeling for crops and specific use directions.

Application
Apply Treflan TR-10 to soils that are wet or are subject to prolonged periods of flooding as poor weed control may result.
Do not use Treflan TR-10 on any crop grown in Pecos county or Reeves county, Texas.
In Montana, uses of Treflan TR-10 are limited to those described in supplemental labeling. Refer to supplemental labeling for crops and specific use directions.

Application
Apply Treflan TR-10 to soils that are wet or are subject to prolonged periods of flooding as poor weed control may result.
Do not use Treflan TR-10 on any crop grown in Pecos county or Reeves county, Texas.
In Montana, uses of Treflan TR-10 are limited to those described in supplemental labeling. Refer to supplemental labeling for crops and specific use directions.

Application
Apply Treflan TR-10 to soils that are wet or are subject to prolonged periods of flooding as poor weed control may result.
Do not use Treflan TR-10 on any crop grown in Pecos county or Reeves county, Texas.
In Montana, uses of Treflan TR-10 are limited to those described in supplemental labeling. Refer to supplemental labeling for crops and specific use directions.

Application
Apply Treflan TR-10 to soils that are wet or are subject to prolonged periods of flooding as poor weed control may result.
Do not use Treflan TR-10 on any crop grown in Pecos county or Reeves county, Texas.
In Montana, uses of Treflan TR-10 are limited to those described in supplemental labeling. Refer to supplemental labeling for crops and specific use directions.

Application
Apply Treflan TR-10 to soils that are wet or are subject to prolonged periods of flooding as poor weed control may result.
Do not use Treflan TR-10 on any crop grown in Pecos county or Reeves county, Texas.
In Montana, uses of Treflan TR-10 are limited to those described in supplemental labeling. Refer to supplemental labeling for crops and specific use directions.

Application
Apply Treflan TR-10 to soils that are wet or are subject to prolonged periods of flooding as poor weed control may result.
Do not use Treflan TR-10 on any crop grown in Pecos county or Reeves county, Texas.
In Montana, uses of Treflan TR-10 are limited to those described in supplemental labeling. Refer to supplemental labeling for crops and specific use directions.

Application
Apply Treflan TR-10 to soils that are wet or are subject to prolonged periods of flooding as poor weed control may result.
Do not use Treflan TR-10 on any crop grown in Pecos county or Reeves county, Texas.
In Montana, uses of Treflan TR-10 are limited to those described in supplemental labeling. Refer to supplemental labeling for crops and specific use directions.

Application
Apply Treflan TR-10 to soils that are wet or are subject to prolonged periods of flooding as poor weed control may result.
Do not use Treflan TR-10 on any crop grown in Pecos county or Reeves county, Texas.
In Montana, uses of Treflan TR-10 are limited to those described in supplemental labeling. Refer to supplemental labeling for crops and specific use directions.

Application
Apply Treflan TR-10 to soils that are wet or are subject to prolonged periods of flooding as poor weed control may result.
Do not use Treflan TR-10 on any crop grown in Pecos county or Reeves county, Texas.
In Montana, uses of Treflan TR-10 are limited to those described in supplemental labeling. Refer to supplemental labeling for crops and specific use directions.

Application
Apply Treflan TR-10 to soils that are wet or are subject to prolonged periods of flooding as poor weed control may result.
Do not use Treflan TR-10 on any crop grown in Pecos county or Reeves county, Texas.
In Montana, uses of Treflan TR-10 are limited to those described in supplemental labeling. Refer to supplemental labeling for crops and specific use directions.

Application
Apply Treflan TR-10 to soils that are wet or are subject to prolonged periods of flooding as poor weed control may result.
Do not use Treflan TR-10 on any crop grown in Pecos county or Reeves county, Texas.
In Montana, uses of Treflan TR-10 are limited to those described in supplemental labeling. Refer to supplemental labeling for crops and specific use directions.

Application
Apply Treflan TR-10 to soils that are wet or are subject to prolonged periods of flooding as poor weed control may result.
Do not use Treflan TR-10 on any crop grown in Pecos county or Reeves county, Texas.
In Montana, uses of Treflan TR-10 are limited to those described in supplemental labeling. Refer to supplemental labeling for crops and specific use directions.

Application
Apply Treflan TR-10 to soils that are wet or are subject to prolonged periods of flooding as poor weed control may result.
Do not use Treflan TR-10 on any crop grown in Pecos county or Reeves county, Texas.
In Montana, uses of Treflan TR-10 are limited to those described in supplemental labeling. Refer to supplemental labeling for crops and specific use directions.

Application
Apply Treflan TR-10 to soils that are wet or are subject to prolonged periods of flooding as poor weed control may result.
Do not use Treflan TR-10 on any crop grown in Pecos county or Reeves county, Texas.
In Montana, uses of Treflan TR-10 are limited to those described in supplemental labeling. Refer to supplemental labeling for crops and specific use directions.

Application
Apply Treflan TR-10 to soils that are wet or are subject to prolonged periods of flooding as poor weed control may result.
Do not use Treflan TR-10 on any crop grown in Pecos county or Reeves county, Texas.
In Montana, uses of Treflan TR-10 are limited to those described in supplemental labeling. Refer to supplemental labeling for crops and specific use directions.

Application
Apply Treflan TR-10 to soils that are wet or are subject to prolonged periods of flooding as poor weed control may result.
Do not use Treflan TR-10 on any crop grown in Pecos county or Reeves county, Texas.
In Montana, uses of Treflan TR-10 are limited to those described in supplemental labeling. Refer to supplemental labeling for crops and specific use directions.

Application
Apply Treflan TR-10 to soils that are wet or are subject to prolonged periods of flooding as poor weed control may result.
Do not use Treflan TR-10 on any crop grown in Pecos county or Reeves county, Texas.
In Montana, uses of Treflan TR-10 are limited to those described in supplemental labeling. Refer to supplemental labeling for crops and specific use directions.

Application
Apply Treflan TR-10 to soils that are wet or are subject to prolonged periods of flooding as poor weed control may result.
Do not use Treflan TR-10 on any crop grown in Pecos county or Reeves county, Texas.
In Montana, uses of Treflan TR-10 are limited to those described in supplemental labeling. Refer to supplemental labeling for crops and specific use directions.

Application
Apply Treflan TR-10 to soils that are wet or are subject to prolonged periods of flooding as poor weed control may result.
Do not use Treflan TR-10 on any crop grown in Pecos county or Reeves county, Texas.
In Montana, uses of Treflan TR-10 are limited to those described in supplemental labeling. Refer to supplemental labeling for crops and specific use directions.

Application
Apply Treflan TR-10 to soils that are wet or are subject to prolonged periods of flooding as poor weed control may result.
Do not use Treflan TR-10 on any crop grown in Pecos county or Reeves county, Texas.
In Montana, uses of Treflan TR-10 are limited to those described in supplemental labeling. Refer to supplemental labeling for crops and specific use directions.

Grain Sorghum (Milo)

Treflan TR-10 may be applied as a postemergence treatment following the use of a preemergence herbicide. Uniformly apply to the soil surface when grain sorghum is 8 to 24 inches tall. Treflan TR-10 does not control established weeds.

Soil Preparation

Cultivate soil before application to remove established weeds and cover the base of plants with soil. Set cultivation equipment to add approximately one inch of soil to the base of sorghum plants.

Activation and Incorporation Directions

Treflan TR-10 should be incorporated within 24 hours after application. Incorporation may be accomplished with one pass of a sweep-type cultivator or properly adjusted rolling cultivator. The sweep-type cultivator should have 3 to 5 sweeps per row in the middle and be operated at a speed sufficient to provide vigorous soil mixing. Set middle sweeps so as to avoid exposing untreated soil. Adjust incorporation equipment so as to avoid mechanical injury to the crop.

Broadcast Application Rates/Acre:

Soil Texture	Treflan TR-10 (lb)
coarse	4 - 5
medium	5 - 7.5
fine	7.5 - 10

Apply lower rate in rate range in areas receiving less than 20 inches total annual rainfall and irrigation.

Restriction: Do not apply after grain sorghum is 24 inches tall. Do not apply to grain sorghum as a preplant or preemergence treatment or crop injury will occur.

Greens: Collard, Kale, Mustard and Turnip (Fresh, for Processing, or Grown for Seed)

Apply to greens as a preplant incorporated treatment.

Broadcast Application Rates/Acre:

Soil Texture	Treflan TR-10 (lb)
coarse	5
medium	7.5
fine	7.5

Hops

Apply and incorporate Treflan TR-10 to established hops during dormancy. Incorporate once using incorporation equipment that will insure thorough soil mixing with minimal damage to crop stand.

Broadcast Application Rates/Acre:

Soil Texture	Treflan TR-10	
	Spring ¹ Application (lb)	Fall ² Application (lb)
coarse	5	7.5
medium	6.25 - 7.5	10
fine	7.5 - 10	12.5

¹Spring application:
• Coarse and medium soils with 2 to 5% organic matter - 7.5 lb product per acre
• Fine soils with 2 to 5% organic matter - 10 lb product per acre

²Fall application:
• Coarse and medium soils with 2 to 5% organic matter - 7.5 lb product per acre
• Fine soils with 2 to 5% organic matter - 10 lb product per acre

• Use lower rate in rate range in areas receiving less than 20 inches total annual rainfall and irrigation.

Kenaf

Apply as a preplant soil incorporated treatment.

Broadcast Application Rates/Acre:

Soil Texture	Treflan TR-10 (lb)
coarse	5.0
medium	5.0 - 7.5
fine	7.5

• Coarse soils with 2 to 5% organic matter - 7.5 lb product per acre
• Use higher rate in rate range where high weed populations are anticipated.

Restriction: Do not graze or harvest treated crop for livestock forage.

Oilseed - Crab, Rapeseed (Canola Varieties Only)

Apply as a soil incorporated treatment in the spring before planting or in late summer or early fall before a fall planting. Follow soil preparation, application and incorporation directions for Treflan TR-10.

Broadcast Application Rates/Acre:

Soil Texture	Treflan TR-10 (lb)
coarse	5.0
medium	7.5
fine	10.0

Restrictions:
• Do not apply to rapeseed (canola) grown in the state of Alaska.
• Do not graze or harvest crabs for livestock forage.

Precaution:

• Where applications are made in late summer or fall, plant as rotation crops in the season following application only those crops to which Treflan TR-10 may be applied as a preplant incorporated treatment or crop injury may occur.

Oilseed -Castor oil plant, Sunflower

Apply and incorporate Treflan TR-10 in the spring or in the fall between September 15 and December 31 in California, Minnesota, North Dakota and South Dakota, and between October 15 and December 31 in other states.

Broadcast Application Rates/Acre:

Soil Texture	Treflan TR-10 (lb)
coarse	5
medium	6.25 - 7.5
fine	7.5 - 10

• Coarse and medium soils with 2 to 5% organic matter - 7.5 lb product per acre
• All soils with 5 to 10% organic matter - 10 lb product per acre
• Use lower rate in rate range in areas receiving less than 20 inches total annual rainfall and irrigation.

Oilseed - Safflower

Apply and incorporate Treflan TR-10 in the spring before planting, or in the fall. See instructions for fall application under Application Timing in the Product Information section of this label.

Broadcast Application Rates/Acre:

Soil Texture	Treflan TR-10	
	Spring ¹ Application (lb)	Fall ² Application (lb)
coarse	5	7.5
medium	6.25 - 7.5	10
fine	7.5 - 10	12.5

¹Spring application:
• Coarse and medium soils with 2 to 5% organic matter - 7.5 lb product per acre
• Fine soils with 2 to 5% organic matter - 10 lb product per acre

²Fall application:
• Coarse and medium soils with 2 to 5% organic matter - 7.5 lb product per acre
• Fine soils with 2 to 5% organic matter - 10 lb product per acre

• Use lower rate in rate range in areas receiving less than 20 inches total annual rainfall and irrigation.

Okra

Apply and incorporate Treflan TR-10 before planting.

Broadcast Application Rates/Acre:

Soil Texture	Treflan TR-10 (lb)
coarse	5
medium	6.25 - 7.5
fine	7.5 - 10

• Coarse soils with 2 to 5% organic matter - 7.5 lb product per acre
• Use higher rate in rate range where high weed populations are anticipated.

• Coarse and medium soils with 2 to 5% organic matter - 7.5 lb product per acre
• Fine soils with 2 to 5% organic matter - 10 lb product per acre
• All soils with 5 to 10% organic matter - 10 lb product per acre
• Use lower rate in rate range in areas receiving less than 20 inches total annual rainfall and irrigation.

Onion (Dry Bulbs Only)

Postemergence Lay-by Application

Apply at lay-by to the soil between onion rows. Avoid applying directly to the tops or exposed bulbs of onion plants. Emerged weeds should be removed prior to application of Treflan TR-10. Treflan TR-10 will not control established weeds.

Broadcast Application Rates/Acre:

Soil Texture	Treflan TR-10 (lb)
coarse	3.75 - 5.0
medium	5.0 - 6.25

• Apply only to soils containing 3.5% or less organic matter
• **Note:** Use the lower rate in rate range where light weed pressure is anticipated.

Incorporation Directions

Treflan TR-10 should be uniformly incorporated into the soil between the onion rows. Incorporation may be accomplished by operating a sweep-type or rolling cultivator 2 to 3 inches deep at 4 to 6 mph. Two incorporation passes are required with the first occurring within 24 hours after application or erratic weed control may result. Delay the second incorporation for a minimum of 5 days after the first. Avoid covering onions with treated soil during incorporation as injury to the crop may occur. Care should be taken to avoid mechanical injury to onion roots during incorporation.

Broadcast Application Rates/Acre:

Soil Texture	Treflan TR-10 (lb)
coarse	5
medium	6.25 - 7.5
fine	7.5 - 10

• Coarse and medium soils with 2 to 5% organic matter - 7.5 lb product per acre
• All soils with 5 to 10% organic matter - 10 lb product per acre
• Use lower rate in rate range in areas receiving less than 20 inches total annual rainfall and irrigation.

Peas - English Peas

Apply and incorporate Treflan[®] TR-10[®] granular herbicide in the spring before planting or in the fall. Refer to instructions for fall application under Application Timing in the Product Information section of this label.

Broadcast Application Rates/Acre:

Soil Texture	Treflan TR-10	
	Spring Application (lb)	Fall ² Application (lb)
coarse	5	5
medium	6.25 - 7.5	10
fine	7.5 - 10	7.5

¹Treflan TR-10 may be fall applied to dry and English peas in the states of Idaho, Oregon and Washington.

²Medium soils with 3% or greater organic matter - 7.5 lb per acre.

• Use the lower rate in areas receiving less than 20 inches total annual rainfall and irrigation.

Peas - Southern Peas

Apply and incorporate Treflan TR-10 before planting.

Broadcast Application Rates/Acre:

Soil Texture	Treflan TR-10 (lb)
coarse	5
medium	6.25 - 7.5
fine	7.5 - 10

• Coarse and medium soils with 2 to 5% organic matter - 7.5 lb product per acre
• Fine soils with 2 to 5% organic matter - 10 lb product per acre
• All soils with 5 to 10% organic matter - 10 lb product per acre
• Use lower rate in rate range in areas receiving less than 20 inches total annual rainfall and irrigation.

Peanuts

(For Use in Texas, Oklahoma and New Mexico Only)

Apply and incorporate Treflan TR-10 before planting, at planting or immediately after planting. When incorporating after planting, adjust equipment to avoid disturbing planted seeds.

Broadcast Application Rates/Acre:

Soil Texture	Treflan TR-10 (lb)
coarse	5
medium	7.5

Pepper (Transplant Only)

Apply and incorporate Treflan TR-10 prior to transplanting. Do not apply after transplanting.

Broadcast Application Rates/Acre:

Soil Texture	Treflan TR-10 (lb)
coarse	5
medium	6.25 - 7.5
fine	7.5 - 10

• Coarse and medium soils with 2 to 5% organic matter - 7.5 lb product per acre
• Fine soils with 2 to 5% organic matter - 10 lb product per acre
• All soils with 5 to 10% organic matter - 10 lb product per acre
• Use lower rate in rate range in areas receiving less than 20 inches total annual rainfall and irrigation.

Potatoes

(Not for Use in the State of Maine)

Apply and incorporate Treflan TR-10 after planting prior to crop emergence, immediately following dragoff, or after potato plants have fully emerged.

Incorporation Directions

Set incorporation equipment so that the bed and furrow are uniformly covered with a layer of treated soil. If the layer of treated soil is not uniform and the herbicide is concentrated over the bed, potato emergence may be retarded and stem brittleness may occur. When applying and incorporating Treflan TR-10 after potato plants have fully emerged, do not completely cover the plants with treated soil. Likewise, do not completely cover plants during subsequent cultivation. Be careful that incorporation equipment does not damage potato seed pieces or elongating sprouts.

Soil Texture	Treflan TR-10 (lb)	
	Spring Application (lb)	Fall ² Application (lb)
coarse	5	5
medium	5 - 7.5 ²	6.25 - 7.5
fine	7.5	7.5

¹Treflan TR-10 may be fall applied to dry and English peas in the states of Idaho, Oregon and Washington.

²Medium soils with 3% or greater organic matter - 7.5 lb per acre.

• Use the lower rate in areas receiving less than 20 inches total annual rainfall and irrigation.

Small Grains - Barley, Durum and Wheat

Special Precautions for Use of Treflan TR-10 on Wheat

Carefully follow directions for use of Treflan[®] TR-10[®] granular herbicide on small grains to minimize potential crop stress. Under certain conditions, delayed crop emergence and/or stand reduction may occur when Treflan TR-10 is applied to barley, durum or wheat. The combined effect of certain cultural practices and unfavorable soil or environmental conditions may cause excessive crop

seedling stress resulting in retarded crop growth, stand reduction and possibly reduced yield. **For best results, observe the following cultural practices or precautions:**

Use tillage methods that provide a uniformly firm seedbed and time tillage operations to conserve moisture.
Irrigate prior to planting or after germination and emergence. Moisture received between planting and emergence may cause crusting, especially on loess seedbeds.

Do not exceed soil-specific application rates for Treflan TR-10. This is particularly important on coarse textured or low organic matter soils.

Carefully follow incorporation directions. When applying preplant incorporated treatments, operate equipment at recommended depth and speed to place Treflan TR-10 into the upper 1 to 1.5 inches of soil. If applied after planting, set equipment so as to avoid disturbance of planted seed.

Set drills to place seed at the depth specified in use directions. A planting depth greater than 2.5 inches for spring wheat or durum will result in increased seedling stress and decreased emergence.

Use only high quality seed where Treflan TR-10 is to be applied (avoid use of small seed with low starch reserves).

If seed treatments are used, apply at the correct rate and uniformly across all seeds. Misapplication may result in reduced germination and/or seedling vigor.

Avoid use of seed varieties known to have poor seedling (emergence) vigor.

Soil characteristics and environmental conditions which may contribute to reduced emergence rates that may be accentuated by use of Treflan TR-10 include:
Soil related: High salinity, eroded knolls/hilltops, loose dry soils and compaction.

Weather related: Cold and/or wet soils, excessively hot soils, excessive moisture, drought, and soil crusting from heavy rainfall.

Restriction: Do not apply Treflan TR-10 on small grains where a dinitroaniline herbicide such as Treflan (trifluralin) or Sonalan[®] herbicide (ethalfuralin) was applied at a rate greater than 0.5 lb active ingredient per acre the previous growing season.

Incorporation Directions and Equipment for Small Grains

Apply and incorporate Treflan TR-10 in the manner described below unless otherwise specified in use directions. Any of the following implements listed below may be used for the first incorporation of Treflan TR-10. Use only the disc or field cultivator for the second incorporation pass and incorporate in a different direction.

Poor weed control may result if untreated soil is moved to the surface during the second incorporation pass. To avoid this problem, the second incorporation should not be deeper than the first.

Chisel Plow: May be used for the first incorporation pass only. Operate at 4 to 5 inches deep and at 4 to 6 mph. A chisel plow is defined as having 3 rows of 14- to 18-inch sweeps on no greater than 12-inch centers. Stagger sweeps so that no soil is left untreated. Use of chisel points may result in inadequate soil mixing.

Tandem Disc: Operate at 3 to 4 inches deep and at 4 to 6 mph.

Field Cultivator: Operate at 3 to 4 inches deep and at 5 mph or more. A field cultivator is defined as having 3 to 4 rows of sweeps with "C" or "S" shaped shanks on no greater than 7-inch centers. Stagger sweeps so that no soil is left untreated.

Application Directions for Small Grains

Barley, Spring Seeded -- Fall Application for Weed Control During the Following Season (for Use in Minnesota, North Dakota and South Dakota)

For weed control (except for special rate or use programs) in Minnesota, North Dakota and South Dakota.

Apply in the fall for weed control during the following growing season. Incorporate one time within 24 hours. Incorporate a second time before planting to destroy existing weeds and insure uniform distribution of Treflan TR-10 in treated soil. Delay the second incorporation a minimum of 5 days after the first.

Note: See information on incorporation and incorporation equipment at the beginning of the Small Grains section.

Planting Directions: Set planting equipment to place seed approximately 1.5 inches deep.

Broadcast Application Rate: Set planting equipment to place seed approximately 1.5 inches deep.

Precautions: Use of this practice may result in a slight stand reduction. Follow the most severe grazing restrictions imposed by either the USDA for Treflan TR-10 or the USDA Acresage Conservation Reserve Program, whichever is longest. Consult the local ASCS office or other state agency to determine the period of the USDA grazing restriction.

Spring Seeded Wheat or Durum - Spring Application Preplant Incorporated for Suppression of Foxtail (Pigeongrass) (for Use in North Dakota West of ND Highway Number 3 and in South Dakota)

Apply Treflan TR-10 in the spring as a preplant incorporated treatment for suppression of green and yellow foxtail (pigeongrass) in spring seeded wheat or durum.

Broadcast Application Rate: Set planting equipment to place seed approximately 1.5 inches deep.

Broadcast Application Rates/Acre:

Soil Texture	Treflan TR-10 (lb)
coarse	5
medium	7.5
fine	7.5

Treflan TR-10 applied at 7.5 lb per acre will provide partial control or suppression of Kochia and Russian thistle.

Precaution: While use of this practice may result in a stand reduction, slight stand reductions do not normally affect yield.

Barley, Spring Seeded -- Spring Application for Foxtail (Pigeongrass) Control (for Use in Minnesota, North Dakota and South Dakota)

Apply in the spring as a preplant incorporated treatment for foxtail (pigeongrass) control in spring seeded barley.

Incorporate one time within 24 hours. Incorporate a second time before planting to destroy existing weeds and insure uniform distribution of Treflan TR-10 in treated soil. Delay the second incorporation a minimum of 5 days after the first.

Note: See information on incorporation and incorporation equipment at the beginning of the Small Grains section.

Planting Directions: Set planting equipment to place seed approximately 1.5 inches deep.

Broadcast Application Rates/Acre:

Soil Texture	Treflan TR-10 (lb)
coarse	5
medium	5
fine	5

Precaution: While use of this practice may result in a stand reduction, slight stand reductions do not normally affect yield.

Spring Seeded Durum or Spring Wheat - Fall Applied Preplant Incorporated for Suppression of Foxtail (Pigeongrass) (for Use in North Dakota West of the Red River Valley and in South Dakota)

Apply Treflan TR-10 in the fall as a preplant incorporated treatment for suppression of green and yellow foxtail (pigeongrass) in spring seeded wheat.

Broadcast Application Rates/Acre:

Soil Texture	Treflan TR-10 (lb)
coarse	5
medium	3.5 - 4.0
fine	5.0

Use the lower rate in the rate range where light weed pressure is anticipated.

Incorporation Directions
The first incorporation should occur within 24 hours after application. Incorporate a second time before planting to destroy existing weeds and insure uniform distribution of Treflan TR-10 in treated soil. Delay the second incorporation a minimum of 5 days after the first.

Note: See information on incorporation and incorporation equipment at the beginning of the Small Grains section.

Broadcast Application Rates/Acre:

Soil Texture	Treflan TR-10 (lb)
coarse	5
medium	3.5 - 4.0
fine	5.0

Precautions: Use of this practice may result in a slight stand reduction. Follow the most severe grazing restrictions imposed by either the USDA for Treflan TR-10 or the USDA Acresage Conservation Reserve Program, whichever is longest. Consult the local ASCS office or other state agency to determine the period of the USDA grazing restriction.

Spring Seeded Wheat or Durum - Spring Application Preplant Incorporated for Suppression of Foxtail (Pigeongrass) (for Use in North Dakota West of ND Highway Number 3 and in South Dakota)

Apply Treflan TR-10 in the spring as a preplant incorporated treatment for suppression of green and yellow foxtail (pigeongrass) in spring seeded wheat or durum.

Broadcast Application Rate: Set planting equipment to place seed approximately 1.5 inches deep.

Precautions: Use of this practice may result in a slight stand reduction. Follow the most severe grazing restrictions imposed by either the USDA for Treflan TR-10 or the USDA Acresage Conservation Reserve Program, whichever is longest. Consult the local ASCS office or other state agency to determine the period of the USDA grazing restriction.

Winter Wheat -- Preplant Incorporated for Control of Cheatgrass and Other Annual Grasses (for Use in Idaho, Oregon and Washington)

Apply Treflan TR-10 as a preplant incorporated treatment up to 3 weeks before planting.

Broadcast Application Rates/Acre:

Soil Texture	Treflan TR-10 (lb)
coarse	7.5
medium	10
fine	7.5

Incorporate Treflan TR-10 with a flexible tine-tooth harrow (Flexline or Melroe) set to cut 1 to 2 inches deep and operated at 3 to 6 mph. Thorough incorporation requires two incorporation passes over the field in different directions. Incorporate one time within 24 hours after application. Make a second incorporation pass prior to planting and a minimum of 5 days after the first. Do not till the soil with a disc after Treflan TR-10 has been incorporated with a flexible tine harrow.

Planting Directions: Set equipment to place seed approximately 1.5 inches deep.

Precaution: While use of this practice may result in a stand reduction, slight stand reductions do not normally affect yield.

Winter Wheat -- Fall Application for Annual Grass Control Prior to Planting (for Use in Idaho, Oregon and Washington)

Apply and shallowly incorporate Treflan TR-10 up to 4 months before planting to control cheatgrass and certain other annual grasses and broadleaf weeds during the fallow period and during the following growing season. Apply any time from May to September prior to fall planting of winter wheat.

Precaution: While use of this practice may result in a stand reduction, slight stand reductions do not normally affect yield.

Planting Directions: Set planting equipment to place seed approximately 1.5 inches deep.

Broadcast Application Rates/Acre:

Soil Texture	Treflan TR-10 (lb)
coarse	5
medium	5
fine	7.5

Precaution: While use of this practice may result in a stand reduction, slight stand reductions do not normally affect yield.

Spring Seeded Durum or Spring Wheat - Fall Applied Preplant Incorporated for Suppression of Foxtail (Pigeongrass) (for Use in North Dakota West of the Red River Valley and in South Dakota)

Apply Treflan TR-10 in the fall as a preplant incorporated treatment for suppression of green and yellow foxtail (pigeongrass) in spring seeded wheat.

Broadcast Application Rates/Acre:

Soil Texture	Treflan TR-
--------------	-------------

900-011166
PBOM 16 x 4.75 x 32

A-000-000-C Yellow Magenta Cyan Black Black Black Black

TOP
Live Copy Area



Dow AgroSciences

Treflan®

TR-10®

GRANULAR HERBICIDE

A selective herbicide for the preemergence control of annual grasses and broadleaf weeds in Alfalfa, Almond, Apricot, Asparagus, Barley, Beans – All Dry and Fresh Beans/Peas, Broccoli, Brussels Sprouts, Cabbage, Carrot, Castor Oil Plant, Cauliflower, Celery, Chicory, Collard Greens, Corn, Cotton, Crambe, Cucurbits, Dry Peas, Durum, English Peas, Flax, Flaxseed, Grain Sorghum, Grapes, Grapefruit, Guar, Hops, Kale, Kenaf, Lemon, Lima Bean, Mungbean, Mustard Greens, Nectarine, Niger seed, Oil radish, Okra, Onions, Orange, Peach, Peanuts, Pecan, Pepper, Plum, Potatoes, Prune, Radish, Rapeseed, Safflower, Snap Bean, Southern Peas, Soybeans, Sugar Beets, Sugarcane, Sunflowers, Tangelo, Tangerine, Tomatoes, Turnip Greens, Walnut, and Wheat

Active Ingredient:
trifluralin: α, α, α -trifluoro-2,6-dinitro-*N,N*-dipropyl-*p*-toluidine 10%
Other Ingredients..... 90%
Total..... 100%

Contains 5 lb active ingredient per 50 lb bag.

**Keep Out of Reach of Children
CAUTION PRECAUCION**

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. Refer to label booklet under "Agricultural Use Requirements" in the Directions for Use section for information about this standard.

Notice: Read the entire label. Use only according to label directions. **Before using this product, read Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies at end of label booklet. If terms are unacceptable, return at once unopened.**

In case of emergency endangering health or the environment involving this product, call 1-800-992-5994.

Agricultural Chemical: Do not ship or store with food, feeds, drugs or clothing.

EPA Reg. No. 62719-131

EPA Est. 5905-IA-01

®Trademark of The Dow Chemical Company ("Dow") or an affiliated company of Dow

Produced for
Dow AgroSciences LLC
9330 Zionsville Road
Indianapolis, IN 46268

NET WEIGHT 50 LB



(01) 0 06 62974 32543 7

00032543

Treflan® TR-10®

GRANULAR HERBICIDE

900-021311 / 00013774

Dow AgroSciences



Dow AgroSciences

Treflan® TR-10®

GRANULAR HERBICIDE

900-021311 / 00013774

00032543

Precautionary Statements
Hazards to Humans and Domestic Animals
Causes Eye Irritation • Prolonged Or Frequently Repeated Skin Contact May Cause Allergic Reactions In Some Individuals
Avoid contact with eyes or clothing.
Personal Protective Equipment (PPE)
Applicators and other handlers must wear:
• Long-sleeved shirt and long pants
• Chemical-resistant gloves, such as butyl rubber \geq 11 mils; or natural rubber \geq 14 mils; or neoprene rubber \geq 14 mils; or nitrile rubber \geq 14 mils
• Shoes plus socks
Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.
Engineering Controls
When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d) (4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.
User Safety Recommendations
Users should:
• Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
• Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
• Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.
First Aid
If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-992-5994 for emergency medical treatment information.
Environmental Hazards
This pesticide is extremely toxic to freshwater marine, and estuarine fish and aquatic invertebrates including shrimp and oyster. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not apply in a manner which will directly expose canals, lakes, streams, ponds, marshes or estuaries to aerial drift. Do not contaminate water when disposing of equipment washwaters.
Storage and Disposal
Do not contaminate water, food, or feed by storage and disposal.
Pesticide Storage: Store in original container only. In case of spill, contain material and dispose as waste.
Pesticide Disposal: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.
Container Handling: Nonrefillable container. Do not reuse or refill this container. Completely empty bag into application equipment. Then offer for recycling if available, or dispose in a sanitary landfill, or by incineration, or by other procedures allowed by state and local authorities.
Refer to label booklet for Directions for Use.
DIRECTIONS ENCLOSED

BUTT
Live Copy Area