



ACCEPTED

LICENSE NO. **9805.107**

Change Up™

Selective Herbicide

**FOR SELECTIVE BROADLEAF WEED CONTROL
IN ORNAMENTAL LAWNS AND TURF GRASSES.**

CONTAINS MCPA, FLUROXYPYR AND DICAMBA

CONTROLS: Dandelion, Chickweed, Black medic, Knotweed, Plantain, Oxalis, Clover, Cocklebur, Thistle and many other species of broadleaf weeds; some of which are listed on this label.

ACTIVE INGREDIENTS:

Dimethylamine Salt of 2-Methyl-4-Chlorophenoxyacetic Acid*	51.05%
1-Methylheptyl Ester of Fluroxypyr: [(4-amino-3-5-dichloro-6-fluoro-2-pyridinyl)oxy]acetic Acid, 1-methylheptyl ester**	6.00%
Dicamba (3,6-Dichloro-o-Anisic Acid)***	4.17%


OTHER INGREDIENTS 38.78%

TOTAL: 100.00%

Isomer Specific Method, Equivalent to:

*2-Methyl-4-Chlorophenoxyacetic Acid	41.68%, 4.0 lbs./gal.
**[(4-amino-3-5-dichloro-6-fluoro-2-pyridinyl)oxy]acetic acid	4.17%, 0.4 lbs./gal.
***3,6-Dichloro-o-Anisic Acid	4.17%, 0.4 lbs./gal.

DO NOT SELL, DISTRIBUTE OR USE THE PRODUCT IN NASSAU AND SUFFOLK COUNTY IN NEW YORK.



**KEEP OUT OF REACH OF CHILDREN
WARNING / AVISO**

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

See inside booklet for FIRST AID and
ADDITIONAL PRECAUTIONARY STATEMENTS

For Chemical Spill, Leak, Fire,
or Exposure, Call CHEMTREC
(800) 424-9300
For Medical Emergencies Only,
Call (877) 325-1840

EPA Reg. No. 228-445

EPA Est. No. 228-IL-1

Net Contents
2.5 Gal.
(9.46 L)

Manufactured for
Nufarm Americas Inc.
11901 S. Austin Ave.,
Alsip, IL 60803



PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS WARNING / AVISO

Causes substantial but temporary eye injury. Harmful if swallowed or absorbed through skin. Do not get in eyes, on skin or on clothing.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Some materials that are chemical-resistant to this product are made of waterproof material. If you want more options, follow the instructions for category A on an EPA chemical-resistant selection chart.

Mixers, loaders, applicators, and other handlers must wear:

- long-sleeved shirt and long pants
- shoes plus socks
- chemical-resistant gloves when mixing, loading, or using any hand-held equipment
- protective eyewear (goggles, face shield, or safety glasses)

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS

Users Should:

- Wash hands, face and arms with soap and water before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

FIRST AID

IF IN EYES	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
IF SWALLOWED	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by the poison control center or doctor. • Do not give anything by mouth to an unconscious person.
IF ON SKIN OR CLOTHING	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15 to 20 minutes. • Call a poison control center or doctor for treatment advice.

HOT LINE NUMBER

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-877-325-1840 for emergency medical treatment information.

ENVIRONMENTAL HAZARDS

Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwater or rinsate. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas. Runoff of this product will be reduced by avoiding applications when rainfall is forecasted to occur within 48 hours.

This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. READ ENTIRE LABEL BEFORE USING THIS PRODUCT. USE STRICTLY IN ACCORDANCE WITH LABEL PRECAUTIONARY STATEMENTS AND DIRECTIONS.

This pesticide should be used strictly in accordance with the drift and run-off precautions on this label in order to minimize off-site exposure. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

Under some conditions this product may have a potential to run-off to surface water or adjacent land. Where possible, use methods which reduce soil erosion, such as no till, limited till and contour plowing; these methods also reduce pesticide run-off. Where feasible, use application techniques such as T-banding and in-furrow techniques which incorporate the pesticide into the soil. Use of vegetation filter strips along rivers, creeks, streams, wetlands, etc. or on the downhill side of fields where run-off could occur will minimize water run-off is recommended.

Low humidity and high temperatures increase the evaporation rate of spray droplets and therefore, the likelihood of increased spray drift. Avoid spraying during conditions of low humidity and/or high temperatures.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about Personal Protective Equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 48 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- coveralls worn over short-sleeved shirt and short pants
- chemical-resistant footwear plus socks
- chemical-resistant gloves made of any waterproof material
- chemical-resistant headgear for overhead exposure, and
- protective eyewear

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses. For Turf use, the maximum number of broadcast applications per treatment site is 2 per year.

Do not allow people (other than applicator) or pets on treatment area during application. Do not enter or allow others to enter the treated area until sprays have dried.

This product is for use on Ornamental Turf Lawns (Residential, Industrial and Institutional), Parks, Cemeteries, Athletic Fields and Golf Courses (Fairways, Aprons, Tees* and Roughs); also for use on Sod Farms.

*Excluding Bentgrass Tees.

SPRAY DRIFT MANAGEMENT

Avoiding spray drift at the application site is the responsibility of the applicator. The interaction of many equipment and weather related factors determine the potential for spray drift. The applicator and the grower are responsible for considering all these factors when making decisions.

Apply only as a medium or coarser spray (ASAE standard 572) or a volume mean diameter of 300 microns or greater for spinning atomizer nozzles.

Apply only when the wind speed is 2–10 mph at the application site.

APPLICATION RESTRICTIONS

Do not apply this product in a way that will contact workers any person, or pet, either directly or through drift. Keep people and pets out of the area during application.

Avoid drift of spray mist to vegetables, flowers, ornamental plants, shrubs, trees and other desirable plants. Do not use on Dichondra, nor on lawns or turf where desirable clovers are present. Avoid fine mists. Use lawn type sprayer with coarse spray as wind drift is less likely. Avoid contact with exposed feeder roots of ornamentals and trees. Maximum control of weeds will be obtained from spring or early fall applications when weeds are actively growing. The degree of weed control and duration of effect will vary with weed size and density, spray rate and coverage, and growing conditions before, during, and after the time of treatment. Use the higher rate for hard-to-control weeds. Do not exceed specified dosages for any area; be particularly careful within the dripline of tree and other ornamental species. Do not apply to newly seeded grasses until well established. Avoid broadcast applications when air temperature exceeds 90°F. When using in small, spot treatment applications in temperatures over 90°F, turf injury may occur.

Aerial application is prohibited.

When treating Carpetgrass and St. Augustine grass, avoid broadcast applications when air temperature exceeds 80°F. When air temperature exceeds 80°F limit application to spot treatment only.

The suitable use of this product on non-recommended turf species may be determined by treating a small area at any rate/acre which does not exceed 3 pints/acre. The treated area should be observed for any sign of turf injury for a period of 30 days of normal growing conditions to determine the phytotoxicity and efficacy to the treated area.

For optimum results: (1) avoid applying during excessively dry or hot periods unless irrigation is used; (2) turf should not be mowed 1 to 2 days before and following application; (3) reseed no sooner than 3 to 4 weeks after application of this product. Adding oil, wetting agent, or other surfactant to the spray may be used to increase effectiveness on weeds but doing so may reduce selectivity to turf resulting in turf damage. Clean and rinse spray equipment using soap or detergent and water, and rinse thoroughly before reuse for other sprays.

WEEDS CONTROLLED

This product is designed to control many broadleaf weeds such as:

Alder	Common mullein	Indiana mallow	Prostrate vervain	Toadflax
Annual yellow	Common	Ironweed	Puncture vine	Trailing Crownvetch
sweet clover	sowthistle	Jewelweed	Purslane	Tumble mustard
Artichoke	Corn Chamomile	Jimsonweed	Ragweed	Tumble pigweed
Aster	Creeping jenny	Kochia	Red clover	Tumbleweed
Austrian fieldcress	Crimson clover	Knawel	Redroot pigweed	Velvet leaf
Bedstraw	Croton	Knotweed	Red sorrel	Venice mallow
Beggartick	Cudweed	Lambsquarter	Redstem filaree	Veronica
Biden	Curly dock	Lespedeza	Rough cinquefoil	Vervain
Bindweed	Curly indigo	Locoweed	Rough fleabane	Vetch
Bird vetch	Dandelion	Lupine	Roundleaved marigold	Virginia buttonweed
Bitterweed	Dead nettle	Mallow	Rush	Virginia creeper
Bitter wintercress	Dock	Marshelder	Russian pigweed	Virginia pepperweed
Black-eyed Susan	Dogbane	Matchweed	St. Johnswort	Wavyleaf bullthistle
Black medic	Dogfennel	Mexicanweed	Scarlet pimpernel	Western clematis
Black mustard	Elderberry	Milk vetch	Scotch thistle	Western salsify
Black-seed plantain	English daisy	Milkweed bloodflower	Sheep sorrel	White clover
Blessed thistle	Fall dandelion	Mugwort	Shepherdspurse	White mustard
Blue lettuce	False dandelion	Morningglory	Slender plantain	Wild mustard
Blue vervain	False flax	Mousear chickweed	Smallflower galinsoga	Wild aster
Box elder	False sunflower	Musk thistle	Smartweed	Wild buckwheat
Bracted plantain	Fiddleneck	Mustard	Smooth dock	Wild carrot
Brassbuttons	Field bindweed	Narrowleaf plantain	Smooth pigweed	Wild four-o'-clock
Bristly oxtongue	Field pansy	Narrowleaf vetch	Sneezeweed	Wild garlic
Broadleaf dock	Fleabane (daisy)	Nettle	Southern wild rose	Wild geranium
Broadleaf plantain	Flixweed	Nutgrass	Sowthistle	Wild lettuce
Broomweed	Florida betony	Orange hawkweed	Spanishneedle	Wild marigold
Buckhorn	Florida pusley	Oxalis	Spatterdock	Wild onion
Buckhorn plantain	Frenchweed	Oxeye daisy	Speedwell	Wild parsnip
Bulbous buttercup	Galinsoga	Parsley-piert	Spiny Amaranth	Wild radish
Bull nettle	Garlic mustard	Parsnip	Spiny cocklebur	Wild rape
Bull thistle	Goathead	Pearlwort	Spotted catsear	Wild strawberry
Burdock	Goatsbeard	Pennycress	Spotted knapweed	Wild sweet potato
Burning nettle	Goldenrod	Pennywort	Spurred spurge	Wild vetch
Bur ragweed	Ground ivy	Peppergrass	Spurge	Willow
Burweed	Gumweed	Pepperweed	Spurweed	Witchweed
Buttercup	Hairy bittercress	Pigweed	Stinging nettle	Woolly morningglory
Canada thistle	Hairy fleabane	Pineywoods bedstraw	Stinkweed	Woodsorrel
Carolina geranium	Hawkweed	Plains coreopsis	Stitchwort	Woolly croton
Carpetweed	Healall	Plantain	Strawberry clover	Woolly plantain
Catchweed bedstraw	Heartleaf drymary	Poison hemlock	Sumac	Wormseed
Catsear	Hedge bindweed	Poison ivy	Sunflower	Yarrow
Catnip	Hedge mustard	Poison oak	Sweet clover	Yellow rocket
Chickweed	Hemp	Pokeweed	Tall nettle	Yellowflower
Chicory	Henbit	Poorjoe	Tall vervain	pepperweed
Cinquefoil	Hoary cress	Povertyweed	Tansy mustard	and other broadleaf
Clover	Hoary plantain	Prickly lettuce	Prickly sida	weeds.
Cockle	Hoary vervain	Primrose	Prostrate knotweed	
Cocklebur	Honeysuckle	Prostrate pigweed	Prostrate spurge	
Coffeebean	Hop clover			
Coffeeweed	Horsenettle			
Common chickweed	Horsetail			

RECOMMENDED MIXING INSTRUCTIONS

Fill spray tank with water. Start agitation and slowly add the recommended amount of concentrate. Maintain continuous agitation after mixing and during application. If this product is allowed to stand in tank for extended periods of time, some separation may occur. Reagitate before use.

ORNAMENTAL LAWNS AND TURF

For residential turf and golf course uses, do not apply more than 1.5 lb ae/acre (3 pints product) per year. Do not apply more than 2 applications per year with a minimum retreatment interval of 21 days.

Apply this product at the rate of 2 to 3 pints in 20 to 240 gallons of water per acre (0.73 to 1.10 fl. oz. in 0.5 to 5.5 gallons of water per 1,000 square feet) to control weeds growing in turf grasses planted to Bahia grass, Bent (excluding golf course greens and tees), Bermudagrass, Bluegrass, Carpetgrass, Centipede grass, Fescue, Ryegrass, St. Augustine grass, and Zoysiagrass.

Lower Volume Equipment: Use as little as 5 gallons of water per acre. Use only application equipment that is capable of spreading a uniform droplet, wetting each weed surface.

NOTE: For all grasses (1) avoid double coverage resulting from overlap unless temporary turf injury can be tolerated and (2) use reduced rates if grass is stressed from heat, drought, etc.

SOD FARMS

For sod farm and grass grown for seed use, do not apply more than 1.5 lb ae/acre (3 pints product) per year. Do not apply more than 2 applications per year with a minimum retreatment interval of 21 days.

This product is also intended for use on sod farms. This herbicide provides selective broadleaf control in warm season and cool season turfgrass established for commercial sod production.

Timing

Apply this herbicide to broadleaf weeds that are actively growing. Follow-up applications may be required for dense infestations of perennial and biennial weeds. Do not apply this product to Floratam St. Augustinegrass.

The application of this herbicide to grass seedlings is recommended after the second mowing.

The application of this product to newly sodded, sprigged, or plugged grasses should be delayed until 3 to 4 weeks after the sodding, sprigging, or plugging operations. Also, applications to dormant bermudagrass, dormant zoysiagrass and dormant bahiagrass are suggested.

Recommended Rates for Sod Farms

For Bluegrass, Ryegrass and Fescue, apply 2 to 3 pints per acre. For Creeping Bentgrass use 1.25 pints per acre. For Common and Hybrid Bermudagrass, Bahia, Zoysia and St. Augustine grass apply 1.5 to 1.8 pints per acre. Spray volume for Sod Farm application is 5 to 175 gallons per acre.

Generally, the lower application rates will provide satisfactory control of sensitive weeds. The higher application rates will be required for dense infestations of perennial weeds, for adverse/extreme environmental conditions, or for weeds beyond the appropriate growth stages.

OTHER NON-CROP AREAS

(fencerows, hedgerows, roadsides (including aprons and guard rails), ditches, rights-of-way, utility, pipelines, powerlines, railroads, airports, commercial plants, storage and lumber yards, barrier strips and firebreaks, equipment areas, nurseries and ornamental plantings, fuel tank farms, pumping stations and other industrial sites)

For Noncropland Rights-of-Way use, do not apply more than 1.5 lb ae/acre (3 pints product) per year. Do not apply more than 2 applications per year with a minimum retreatment interval of 21 days.

For spot treatment Noncropland Rights-of-Way use, do not apply more than 3.0 lb ae/acre per year.

Roadsides (including aprons and guardrails), rights-of-way: For control of broadleaf weeds, mix at a rate of (4 to 5 pints) 0.25 to 0.625 gallons of this product per 50 to 300 gallons of water. This mixture will cover 1 acre. Thoroughly saturate all weeds with spray mixture. Apply any time between the time when plants come into full leaf (Spring) to when the plants begin to go dormant. Best results are obtained when weeds are young and actively growing. Do not cut weeds until herbicide has translocated throughout the plant causing root death. For small broadleaf weeds, use the lower rate. Heavy, dense stands require the higher rate with high water volume. For small (spot) applications with small tank sprayers, apply at the rate of 0.75 to 2.25 ounces of this product per gallon of water and spray to thoroughly wet all foliage.

For best results do not cut weeds. For control of woody plants, apply to both stems and foliage any time from the time foliage is completely matured until the time plants start to go dormant. All leaves, stems and suckers must be completely wet to the ground line for effective control. Regrowth may be anticipated on the more resistant species of this product. Add 4 to 5 pints of this product to 100 gallons of water applying 200 to 600 gallons of spray mixture per acre, depending upon the height and thickness of the brush. Mix thoroughly before spraying.

COMPATIBILITY

This product can be mixed with some liquid fertilizers or liquid iron materials. Because liquid fertilizers and liquid iron differ in pH, free ammonia content, density, salt concentration and percentage of water, a compatibility test (given below) is recommended prior to mixing in the application equipment. All regulations, either State or Federal, relating to the application of liquid fertilizers or liquid iron and this product must be strictly followed. The following compatibility test should always be performed prior to full-scale mixing.

1. Pour 18 ounces of water into a quart jar.
2. Add 1 ounce of either the liquid fertilizer or liquid iron to be used.
3. Add 1 ounce of this product.
4. Close jar and shake well.
5. Watch the mixture for several seconds after shaking and check again after 30 minutes.
6. If the mixture does not show signs of separating, the combination may be used. If the mixture foams excessively, gels, separates or gets very thick, do not combine for field application.
7. Compatibility may be improved by the use of a compatibility agent. [Optional Language: Some suggested compatibility agents to try are Kalo Laboratories Complex, Farm Chemicals Inc.'s Compat, Harcros Chemicals' T-Mulz 734-2, Rigo Company's Rigo Compatibility Agent, Witco Chemical's Sponto 168D, Amoco Oil's Amoco Spray Mate and Universal Coop.'s Chem-Link.] These agents are all used in the same manner. Follow the previously outlined test procedures and add 1/6 ounce of the compatibility agent between steps (the compatibility agent must be added to the fertilizer or iron before adding this product.)
8. If the mixture does not separate, gel, foam or get very thick, it may be used for field application. Mix only the amount to be sprayed. Do not allow to stand overnight.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE: Always use original container to store pesticides in a secured warehouse or storage building. Store at temperatures above 25°F. Protect product from freezing. If allowed to freeze, remix well before using. This does not alter this product. Containers should be opened in well ventilated areas. Keep container tightly sealed when not in use. Do not stack cardboard cases more than two pallets high. Do not store near open containers of fertilizer, seed or other pesticides.

PESTICIDE DISPOSAL: Pesticide wastes are toxic. If container is damaged or if pesticide has leaked, contain all spillage. Absorb and clean up all spilled material with granules or sand. Place in a closed labeled container for proper disposal. Improper disposal of excess pesticide, spray mixtures, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL:

[Nonrefillable Containers 5 Gallons or Less:] Nonrefillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by State and local authorities. Plastic containers are also disposable by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

[Nonrefillable Containers Larger than 5 Gallons:] Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available. Triple rinse or pressure rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Pressure rinse as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 psi for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

[Refillable Container Larger than 5 Gallons:] Refillable container. Refill this container with pesticide only. Do not reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from this container into application equipment or a mix tank. Fill the container about 10% full with water and, if possible, spray all sides while adding water. If practical, agitate vigorously or recirculate water with the pump for two minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times.

WARRANTY DISCLAIMER

The directions for use of this product must be followed carefully. **TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, (1) THE GOODS DELIVERED TO YOU ARE FURNISHED "AS IS" BY MANUFACTURER OR SELLER AND (2) MANUFACTURER AND SELLER MAKE NO WARRANTIES, GUARANTEES, OR REPRESENTATIONS OF ANY KIND TO BUYER OR USER, EITHER EXPRESS OR IMPLIED, OR BY USAGE OF TRADE, STATUTORY OR OTHERWISE, WITH REGARD TO THE PRODUCT SOLD, INCLUDING, BUT NOT LIMITED TO MERCHANTABILITY, FITNESS FOR A PARTICULAR PURPOSE, USE, OR ELIGIBILITY OF THE PRODUCT FOR ANY PARTICULAR TRADE USAGE.** UNINTENDED CONSEQUENCES, INCLUDING BUT NOT LIMITED TO INEFFECTIVENESS, MAY RESULT BECAUSE OF SUCH FACTORS AS THE PRESENCE OR ABSENCE OF OTHER MATERIALS USED IN COMBINATION WITH THE GOODS, OR THE MANNER OF USE OR APPLICATION, INCLUDING WEATHER, ALL OF WHICH ARE BEYOND THE CONTROL OF MANUFACTURER OR SELLER AND ASSUMED BY BUYER OR USER. THIS WRITING CONTAINS ALL OF THE REPRESENTATIONS AND AGREEMENTS BETWEEN BUYER, MANUFACTURER AND SELLER, AND NO PERSON OR AGENT OF MANUFACTURER OR SELLER HAS ANY AUTHORITY TO MAKE ANY REPRESENTATION OR WARRANTY OR AGREEMENT RELATING IN ANY WAY TO THESE GOODS.

LIMITATION OF LIABILITY

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO EVENT SHALL MANUFACTURER OR SELLER BE LIABLE FOR SPECIAL, INCIDENTAL, OR CONSEQUENTIAL DAMAGES, OR FOR DAMAGES IN THEIR NATURE OF PENALTIES RELATING TO THE GOODS SOLD, INCLUDING USE, APPLICATION, HANDLING, AND DISPOSAL. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, MANUFACTURER OR SELLER SHALL NOT BE LIABLE TO BUYER OR USER BY WAY OF INDEMNIFICATION TO BUYER OR TO CUSTOMERS OF BUYER, IF ANY, OR FOR ANY DAMAGES OR SUMS OF MONEY, CLAIMS OR DEMANDS WHATSOEVER, RESULTING FROM OR BY REASON OF, OR ARISING OUT OF THE MISUSE, OR FAILURE TO FOLLOW LABEL WARNINGS OR INSTRUCTIONS FOR USE, OF THE GOODS SOLD BY MANUFACTURER OR SELLER TO BUYER. ALL SUCH RISKS SHALL BE ASSUMED BY THE BUYER, USER, OR ITS CUSTOMERS. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BUYER'S OR USER'S EXCLUSIVE REMEDY, AND MANUFACTURER'S OR SELLER'S TOTAL LIABILITY SHALL BE FOR DAMAGES NOT EXCEEDING THE COST OF THE PRODUCT.

If you do not agree with or do not accept any of the directions for use, the warranty disclaimers, or limitations on liability, do not use the product, and return it unopened to the Seller, and the purchase price will be refunded.

(RV071713)

Change Up is a Trademark of Nufarm Americas Inc.
All other trademarks are the property of their respective owners.

— NOTES —

CHANGE UP™

SELECTIVE HERBICIDE

ACTIVE INGREDIENTS:

Dimethylamine Salt of 2-Methyl-4-Chlorophenoxyacetic Acid*	51.05%
1-Methylheptyl Ester of Fluroxypyr:([4-amino-3-5-dichloro-6-fluoro-2-pyridinyl)oxy]acetic Acid, 1-methylheptyl ester**	6.00%
Dicamba (3,6-Dichloro-o-Anisic Acid)***	4.17%

OTHER INGREDIENTS

38.78%

TOTAL:

100.00%

Isomer Specific Method, Equivalent to:	**2-Methyl-4-Chlorophenoxyacetic Acid	41.68%, 4.0 lbs./gal.
	**[(4-amino-3-5-dichloro-6-fluoro-2-pyridinyl)oxy]acetic acid	4.17%, 0.4 lbs./gal.
	***3,6-Dichloro-o-Anisic Acid	4.17%, 0.4 lbs./gal.

KEEP OUT OF REACH OF CHILDREN

WARNING / AVISO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.
(If you do not understand the label, find someone to explain it to you in detail.)

See inside booklet for ADDITIONAL PRECAUTIONARY STATEMENTS and DIRECTIONS FOR USE

For Chemical Spill, Leak, Fire, or Exposure, Call CHEMTREC (800) 424-9300.
For Medical Emergencies Only, Call (877) 325-1840.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

WARNING / AVISO

Causes substantial but temporary eye injury. Harmful if swallowed or absorbed through skin. Do not get in eyes, on skin or on clothing.

FIRST AID	
IF IN EYES	<ul style="list-style-type: none"> Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.
IF SWALLOWED	<ul style="list-style-type: none"> Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor. Do not give anything by mouth to an unconscious person.
IF ON SKIN OR CLOTHING	<ul style="list-style-type: none"> Take off contaminated clothing. Rinse skin immediately with plenty of water for 15 to 20 minutes. Call a poison control center or doctor for treatment advice.
HOT LINE NUMBER	
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-877-325-1840 for emergency medical treatment information.	

ENVIRONMENTAL HAZARDS

Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwater or rinsate. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas. Runoff of this product will be reduced by avoiding applications when rainfall is forecasted to occur within 48 hours.

This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE: Always use original container to store pesticides in a secured warehouse or storage building. Store at temperatures above 25°F. Protect product from freezing. If allowed to freeze, remix well before using. This does not alter this product. Containers should be opened in well ventilated areas. Keep container tightly sealed when not in use. Do not stack cardboard cases more than two pallets high. Do not store near open containers of fertilizer, seed or other pesticides.

PESTICIDE DISPOSAL: Pesticide wastes are toxic. If container is damaged or if pesticide has leaked, contain all spillage. Absorb and clean up all spilled material with granules or sand. Place in a closed labeled container for proper disposal. Improper disposal of excess pesticide, spray mixtures, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL:

[Nonrefillable Containers 5 Gallons or Less:] Nonrefillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by State and local authorities. Plastic containers are also disposable by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

EPA Reg. No. 228-445

EPA Est. No. 228-IL-1

Manufactured by
NUFARM AMERICAS INC.
11901 S. Austin Ave.
Alsip, IL 60803

PULL HERE TO OPEN

14679000

(RV071713)