

STATE OF HAWAII
Department of Agriculture

ACCEPTED

LICENSE NO. **9805.112**

RESTRICTED USE PESTICIDE

TOXIC TO FISH, MAMMALS, AND AQUATIC ORGANISMS
FOR RETAIL SALE TO AND USE ONLY BY CERTIFIED APPLICATORS OR PERSONS UNDER THEIR DIRECT SUPERVISION, AND ONLY FOR THOSE USES COVERED BY THE CERTIFIED APPLICATOR'S CERTIFICATION.

GROUP **6** INSECTICIDE



Abamex™
Miticide/Insecticide

ACTIVE INGREDIENT:

Abamectin (CAS No.71751-41-2):.....2.0%*

OTHER INGREDIENTS:98.0%

TOTAL:100.0%

*Contains 0.15 pounds Abamectin per gallon.

Contains VOC \leq 20% EP; qualifies as "Low VOC" under California regulations

KEEP OUT OF REACH OF CHILDREN
CAUTION

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.

(If you **DO NOT** understand the label, find someone to explain it to you in detail.)

SEE INSIDE LABEL BOOKLET FOR ADDITIONAL FIRST AID AND PRECAUTIONARY STATEMENTS

For Chemical Spill, Leak, Fire, or Exposure, Call CHEMTREC (800) 424-9300

For Medical Emergencies Only, Call (877) 325-1840

EPA Reg. No. 228-734

EPA Est.No. 70815-GA-001

Net Contents

1 Gal. (3.78 L)

Manufactured for
Nufarm Americas Inc.
11901 S. Austin Avenue
Alsip, IL 60803



Grow a better tomorrow.

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
CAUTION

Harmful if swallowed. Causes moderate eye irritation. Avoid contact with eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Some materials that are chemical-resistant to this product are listed below. For more options, follow the instructions for Category B on an EPA chemical-resistant category selection chart.

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Waterproof or Chemical-resistant gloves, such as barrier laminate or butyl rubber \geq 14 mils
- Socks plus shoes

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. **DO NOT** reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

FIRST AID

If swallowed:	<ul style="list-style-type: none">• Call a poison control center or doctor immediately for treatment advice.• DO NOT give any liquid to the person.• DO NOT induce vomiting unless told to do so by the poison control center or doctor.• DO NOT give anything by mouth to an unconscious person.
If in eyes:	<ul style="list-style-type: none">• Hold eye open and rinse slowly and gently with water for 15-20 minutes.• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.• Call a poison control center or doctor for treatment advice.
If on skin or clothing:	<ul style="list-style-type: none">• Take off contaminated clothing.• Rinse skin immediately with plenty of water for 15-20 minutes.• Call a poison control center or doctor for treatment advice.
If inhaled:	<ul style="list-style-type: none">• Move person to fresh air.• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible.• Call a poison control center or doctor for further treatment advice.

(continued)

FIRST AID (continued)

HOT LINE NUMBER

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-877-325-1840 for emergency medical treatment information.

NOTE TO PHYSICIAN

Early signs of intoxication include dilation of pupils, muscular incoordination, and muscle tremors. Toxicity following accidental ingestion of this product can be minimized by early administration of chemical adsorbents (e.g., activated charcoal). If toxicity from exposure has progressed to cause severe vomiting, the extent of resultant fluid and electrolyte imbalance should be gauged. Appropriate supportive parenteral fluid replacement therapy should be given, along with other required supportive measures (such as maintenance of blood pressure levels and proper respiratory functionality) as indicated by clinical signs, symptoms, and measurements. In severe cases, observations should continue for at least several days until clinical condition is stable and normal. Since abamectin is believed to enhance GABA activity in animals, it is probably wise to avoid drugs that enhance GABA activity (barbiturates, benzodiazepines, valproic acid) in patients with potentially toxic abamectin exposure.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish and wildlife. **DO NOT** apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. **DO NOT** apply when weather conditions favor drift from target areas. **DO NOT** contaminate water when disposing of equipment washwater or rinsate. This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. **DO NOT** apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area.

This product may impact surface water quality due to runoff of rain water. This is especially true for poorly draining soils and soils with shallow ground water. This product is classified as having a medium potential for reaching both surface water and aquatic sediment via runoff for several weeks to months after application. A level, well maintained vegetative buffer strip between areas to which this product is applied and surface water features such as ponds, streams, and springs will reduce the potential loading of abamectin from runoff water and sediment. Runoff of this product will be reduced by avoiding applications when rainfall is forecast to occur within 48 hours.

Use of this product may pose a risk to threatened and endangered species of fish, amphibians, crustaceans (including fresh water shrimp), and insects. All use of this product in the state of California should comply with the recommendations of the California Endangered Species Project. Before using this product in California, consult with your county agriculture commissioner to determine use limitations that apply in your area.

ATTENTION: This product contains chemicals known to the state of California to cause birth defects or other reproductive harm.

DIRECTIONS FOR USE

RESTRICTED USE PESTICIDE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. **DO NOT** apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

DO NOT enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours. **Exception:** For grape girdling cane turning, and tying in grapes. **DO NOT** enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 4 days.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls
- Waterproof or Chemical-resistant gloves, such as barrier laminate or butyl rubber \geq 14 mils
- Shoes plus socks

FAILURE TO FOLLOW DIRECTIONS AND PRECAUTIONS ON THIS LABEL MAY RESULT IN CROP INJURY, POOR PEST CONTROL, AND/OR ILLEGAL RESIDUES.

PRODUCT INFORMATION

When applied following the **Directions for Use** on this label, this product controls listed mites and insects in various crops. Optimum control of target insects requires this product to be mixed with sufficient amounts of water (and horticultural oil or nonionic surfactants, when recommended) to provide a thorough coverage of foliage. Spray volume in gallons will vary with the size and number of trees per acre or with the density of target crop foliage. Optimum insect control is achieved only if the foliage receives a thorough coverage of this product. Calibrate all spray equipment to ensure delivery of sufficient spray solution to thoroughly cover foliage of target crops.

Horticultural Oils: When specified on this label, apply this product with horticultural oils (NOT dormant oils) which are approved for use on the target crop. This combination of products will provide residual control and/or improved efficacy. If using concentrations of oils at greater than 2% (v/v), test this rate on small plots for phytotoxicity before using this rate to treat larger plots. To prevent crop injury, observe all precautions and restrictions on the oil label. See precautions under **Phytotoxicity**, below.

Nonionic Surfactants: Nonionic surfactants applied in combination with this product can provide better insect control by allowing the spray mixture to distribute evenly on the leaf and enter the leaf cuticle. When using this product in combination with a nonionic surfactant, carefully follow the Directions for Use and Precautions on the nonionic surfactant label and in official spray guides. See precautions under **Phytotoxicity**, below.

Phytotoxicity: Although this product has been tested for phytotoxicity and compatibility and is safe to use on a variety of crops and as a tank mix in combination with many commonly used products, it is not possible to test all crop varieties grown under a variety of growing conditions or to test all possible combination of products. Before use, test potential mixtures (pesticides, crop oils, nonionic surfactants, and nutritional additives) with this product to ensure physical compatibility and lack of phytotoxic effects to the target crop.

New York State Restrictions: Aerial applications of this product are prohibited in the State of New York.
DO NOT apply this product through any type of irrigation system.

SPRAY DRIFT MANAGEMENT

Except as noted below for specific crops, do not apply with ground application equipment within 25 ft., or by aircraft within 150 ft. of lakes, reservoirs, rivers, permanent streams, marshes, potholes, or natural ponds; estuaries or commercial fish farm ponds. **DO NOT** cultivate within 25 ft. of the aquatic area so as to allow growth of a vegetative filter strip.

All aerial and ground application equipment must be properly calibrated and maintained using appropriate carriers. For aerial applications, the spray boom should be mounted on the aircraft so as to minimize drift caused by wing tip vortices. The minimum practical boom length should be used and must not exceed 75% of wing span or rotor diameter.

Use the largest droplet size consistent with good pest control. Formation of very small droplets may be minimized by appropriate nozzle selection, by orienting nozzles away from the air stream as much as possible, and by avoiding excessive spray boom pressure. Sprays should be released at the lowest height consistent with pest control and flight safety. Applications more than 10 ft. above the crop canopy should be avoided.

Make aerial or ground applications when the wind velocity favors on-target product deposition (approximately 3 to 10 mph). **DO NOT** apply when wind velocity exceeds 15 mph. Avoid applications when wind gusts approach 15 mph. Inclusion of a compatible drift reducing agent is encouraged. Risk of exposure to aquatic areas can be reduced by avoiding applications when wind direction is toward aquatic areas. **DO NOT** cultivate within 25 ft. of the aquatic area so as to allow growth of a vegetative filter strip.

Low humidity and high temperatures increase the evaporation rate of spray droplets and therefore the likelihood of increased spray drift to aquatic areas. Avoid spraying during conditions of low humidity and/or high temperature or, when making applications under these conditions, set up equipment to produce larger droplets to compensate for evaporation.

DO NOT make aerial or ground applications during temperature inversions or if heavy rainfall is expected during application. Inversions are characterized by stable air and increasing temperatures with height above the ground. Mist or fog may indicate the presence of an inversion in humid areas. The applicator may detect the presence of an inversion in humid areas. The applicator may detect the presence of an inversion by producing smoke and observing a smoke layer near the ground surface.

Observe the following precautions when using ground application equipment to spray tree crops (almonds, apples, avocado, citrus, plums, prunes, walnuts) or hops in the vicinity of aquatic areas such as lakes, reservoirs, permanent streams, marshes, potholes, natural ponds, estuaries, or commercial fish ponds.

- **DO NOT** apply this product when weather conditions favor drift to aquatic areas
- **DO NOT** apply within 110 ft. upwind of aquatic areas or when wind speed is above 8 mph.
- Spray last 3 rows upwind of aquatic areas using nozzles on one side only, with spray directed away from the aquatic areas.
- Avoid spray going over tops of trees by adjusting or turning off top nozzles. Shut off nozzles on the side away from the grove/orchard when spraying the outside row. Shut off nozzles when turning at ends of row and passing tree gaps in rows.

RESISTANCE MANAGEMENT

Because of the inherent risks of resistance development to any product, it is strongly advised that this product be used in a sound resistance management program which includes rotation with other products with different modes of action. Effective control of labeled pests will not be achieved if insect or mite-tolerant strains develop. When applying this product to plants that are hosts of labeled pests with multiple generations per crop per year, use resistance management practices such as the following.

- Rotate this product with other products that have different modes of action,
- **DO NOT** apply this product to successive insect generations,
- Follow the use rates and intervals between applications that are specified in this label,
- Use a non-chemical alternative (such as beneficial arthropods),
- Plant susceptible plants following non-susceptible plants, and
- Use other cultural practices

For additional information regarding the implementation of these or other resistance management practices, consult your local agricultural advisor or Nufarm representative.

CROP USE DIRECTIONS
Apple, Pears (including Oriental pear trees)

Target Insects	Application Rate of ABAMEX Miticide/Insecticide fl. oz./A (fl. oz./100 gal)	Application Instructions and Timing	Restrictions
<p>European red mites McDaniel spider mites Pear rust mites Pear psyllas Tentiform leafminers Two-spotted spider mites White apple leafhoppers Yellow mites</p>	<p>Concentrate Sprays: 10-20 fl. oz/A (Dilute Sprays: 2.5-5.0 fl. oz./100 gal) ADD a horticultural oil for best results</p>	<p>Apply using conventional dilute or concentrate ground equipment. Use sufficient volumes (greater than 40 gal of water/A) to ensure thorough foliage coverage. Spider Mites: Apply when mites first appear. For best results, apply before 5 spider mites per leaf is observed. Spray deposits of this product plus a horticultural spray oil on new foliage growth provides greater residual control. Apply this product for spider mite control only during petalfall and up to six weeks following petalfall. Tentiform Leafminers: For best results, apply to control new hatch (eggs) and early stages of first and second generation leafminers. DO NOT apply during bloom. White Apple Leafhopper (Not for use west of the Rocky Mountains): Apply this product only to the first generation soon after petalfall; use in combination with horticultural spray oil. Application without horticultural spray oil may result in reduced efficacy and/or residual control. However, fruit injury to certain apple varieties (on light-skinned varieties such as Golden Delicious) may occur with this combination of this product and spray oil if other products are applied after this spray mix application. Refer to the Directions for Use and Precautions on horticultural spray oil labels. If needed, a second application may be made but allow at least 21 days to elapse after the first application.</p>	<p>Pre-Harvest Interval – 28 days DO NOT exceed 20 fl. oz/A per application or 40 fl. oz/A per growing season. DO NOT apply this product to control white apple leafhopper west of the Rocky Mountains. DO NOT make more than 2 applications per season. DO NOT apply by air. Refer to additional directions on spray oils in the PRODUCT INFORMATION section. See specific restrictions for this crop in the SPRAY DRIFT MANAGEMENT section of this label. To avoid phytotoxicity or loss of crop, do not apply this product plus oil if Captan was applied 14 days before the application or if Captan will be applied 14 days after application. DO NOT graze livestock in treated orchards. Horticultural Spray Oil (DO NOT use a dormant oil) 1 gal./A (minimum) (0.25%, v/v)</p>
<p>NOTE: Concentrate Spray rates (ABAMEX Miticide/Insecticide/A) are determined by selecting the amount needed for a full cover dilute spray. The amount of ABAMEX Miticide/Insecticide/A as a concentrate spray is the same amount as required for dilute sprays to the orchard and could be less than 10 fl. oz. ABAMEX Miticide/ Insecticide/A for small trees. Rates of ABAMEX Miticide/Insecticide/100 gal are based on 400 gal/A of dilute spray.</p>			

AVOCADOS

Target Insects	Application Rate of ABAMEX Miticide/ Insecticide fl. oz./A	Application Instructions and Timing	Restrictions
<p>Avocado thrips (<i>Scirtothrips perseae</i>) Persea Mites (<i>Oligonychus perseae</i>)</p>	<p>DO NOT use less than 10 fl. oz. of this product per Acre</p> <p>Low to moderate insect infestations: 10.0-15.0 Heavy pest infestations: 15.0-20.0</p> <p>ADD a horticultural oil for best results</p>	<p>Apply using ground equipment in 100 or more gal water/A. If using 400 gal/A or more, apply 2.5 fl. oz. of this product/100 gal of water and 1.0-4.0% of NR (narrow range) 415 oil.</p> <p>Although aerial applications are permitted when this product is applied in 50 or more gal water/A, better control of target insects is obtained using ground applications. Users accept all liability for level and duration of thrip control from aerial applications of this product.</p> <p>Apply this product when thrips first appear but prior to finding 5 immature thrips per leaf or fruit. For low to moderate infestations, include 1.0-4.0% of NR 415 oil in combination with this product and the horticultural spray oil approved for use on avocados.</p> <p>If needed, a second application may be made but allow at least 30 days to elapse after the first application.</p>	<p>Pre-Harvest Interval – 14 days</p> <p>DO NOT make more than 2 applications per season.</p> <p>DO NOT exceed 20 fl. oz/A per application or more than 40 fl. oz./A per season.</p> <p>DO NOT allow livestock to graze in treated orchards.</p> <p>Refer to additional directions on spray oils in the PRODUCT INFORMATION section.</p> <p>Horticultural Spray Oil (DO NOT use a dormant oil), minimum 1 to 4%, v/v (minimum)</p>
<p>NOTE:</p> <ul style="list-style-type: none"> • Low infestations are defined as 1-2 immature thrips per leaf/fruit • Moderate infestations are defined as 2-4 thrips per leaf/fruit; • Heavy/severe infestations are defined as more than 5 thrips per leaf/fruit. <p>NOTE: To prevent crop injury, observe all precautions and restrictions on the oil label. When using oil concentrations above 2.0%, treat a small test area before making a large-scale application.</p>			

CELERIAC (*Apium graveolens*)

Target Insects	Application Rate of ABAMEX Miticide/Insecticide fl. oz./A	Application Instructions and Timing	Restrictions
Two-spotted spider mites	16.0	<p>Apply using ground equipment using conventional dilute or concentrate sprayers.</p> <p>Apply when mites first appear using sufficient volumes (greater than 20 gal of water per acre) to ensure thorough foliage coverage.</p> <p>If needed, a second application may be made but allow at least 7 days to elapse after the first application.</p> <p>For best results, use in combination with a non-ionic surfactant.</p>	<p>Pre-Harvest Interval – 7 days</p> <p>To avoid insect resistance to this product, do not apply more than two sequential applications.</p> <p>DO NOT exceed 48 fl. oz/A after transplanting.</p> <p>Refer to additional directions on nonionic surfactants in the PRODUCT INFORMATION section.</p>

CITRUS FRUIT CROP

Crops in this group include calamondin, citrus citron, citrus hybrids (includes chironja, tangelo, tangor), grapefruit, kumquat, lemon, lime, mandarin (tangerine), pummelo, Satsuma mandarin, sour orange, and sweet orange.

Target Insects	Application Rate of ABAMEX Miticide/Insecticide fl. oz./A (fl. oz./100 gal)	Application Instructions and Timing	Restrictions
Broad mites Citrus bud mites Two-spotted spider mites	Concentrate Sprays: 10-20 fl. oz/A (Dilute Sprays: 1.0-2.0 fl. oz./100 gal)	<p>Apply using conventional dilute or concentrate ground equipment.</p> <p>Use sufficient volumes (greater than 40 gal of water/A) to ensure thorough foliage coverage.</p>	<p>Pre-Harvest Interval – 7 days</p> <p>To avoid insect resistance to this product, do not apply to citrus grown in nurseries.</p> <p>DO NOT apply this product, or any other abamectin containing product within 30 days of the last treatment.</p> <p>DO NOT exceed 40 fl. oz/A (0.047 lb. a.i./A) of this product, or any other abamectin containing product per growing season.</p> <p>DO NOT make more than 3 applications of this product, or any other abamectin containing product per growing season.</p>
Citrus rust mites	Concentrate Sprays: 5-20 fl. oz/A (Dilute Sprays: 0.5-2.0 fl. oz./100 gal)	<p>ADD a horticultural oil for best results</p> <p>Asian citrus psylla: Apply to protect newly expanding foliage flush during the spring, summer or fall. ABAMEX Miticide/Insecticide will only control adults and nymphs present at the time of application.</p> <p>Mites: Apply when mites first appear during spring, summer, and/or fall.</p>	
Citrus leafminers	Concentrate Sprays: 5-20 fl. oz/A (Dilute Sprays: NR)		
Asian citrus psyllids Citrus thrips	Concentrate Sprays: 10-20 fl. oz/A (Dilute Sprays: NR)		

(continued)

CITRUS FRUIT CROP (continued)

Target Insects	Application Rate of ABAMEX Miticide/ Insecticide fl. oz./A (fl. oz./100 gal)	Application Instructions and Timing	Restrictions
Asian citrus psyllids Citrus thrips	Concentrate Sprays: 10-20 fl. oz/A (Dilute Sprays: NR)	<p>Citrus bud mite: For best results, time the spray at "bud swell".</p> <p>Citrus leafminer: Apply to protect new growth during spring, summer, or fall.</p> <p>Citrus thrips: Application targeted for citrus thrips will only control the current generation and must be correctly timed. Apply when economic thresholds have been reached (after egg hatch has begun - preferably early to mid-hatch).</p> <p>For Asian citrus psylla, broad mites, citrus leafminers and citrus rust mites optimum results are achieved if spray applications are made at 150-300 gals. Spray mixture/A plus a minimum of 3 gal./A of horticultural oil. Apply with a ground speed of 1 to 1.5 mph.</p> <p>To control citrus leafminer -Ground or aerial application is permitted; however, do not apply with aircraft to citrus in California. For Aerial application use a minimum of 10 gallons of finished spray volume per acre. Under conditions such as high pest populations, dense foliage, or adverse application conditions (such as high temperatures) use a greater volume of water to insure adequate coverage.</p> <p>To manage resistance, if pest infestation requires additional treatments, use other alternate insecticides with different modes of action which are labeled for use on these pests and crops.</p>	<p>DO NOT allow livestock to graze in treated orchards.</p> <p>Refer to additional directions on spray oils in the PRODUCT INFORMATION section. See specific restrictions for this crop in the SPRAY DRIFT MANAGEMENT section of this label.</p> <p>For all other pests besides to control citrus leafminer - DO NOT apply by air.</p> <p>Horticultural Spray Oil (DO NOT use a dormant oil) 1 gal./A (minimum) (0.20%, v/v); see specific rates for certain pests under Application Instructions and Timing</p>
<p>NR = Not recommended.</p> <p>NOTE: Concentrate Spray rates (ABAMEX Miticide/Insecticide /A) are adjusted to apply the amount/A equal to that used in full cover dilute spray. The amount of ABAMEX Miticide/Insecticide/A as a concentrate spray is the same amount as required for dilute sprays to the orchard and could be less than 10 fl oz. ABAMEX Miticide/Insecticide/A for small trees. Rates of ABAMEX Miticide/Insecticide/100 gal are based on 1,000 gal/A of dilute spray.</p>			

COTTON

Target Insects	Application Rate of ABAMEX Miticide/ Insecticide fl. oz./A	Application Instructions and Timing	Restrictions
Carmine spider mites Pacific spider mites Strawberry spider mites Two-spotted spider mites	Early season: 4.0-6.0	Apply this product when mites first appear. Rate will depend on the cotton plant size and thickness of foliage. Use the 4-6 fl. oz./A rate only early in season and for cotton plants 10 inches or less in height. Applications can be made by air or ground using a minimum of 5 gallons of water/A. West of the Rockies: DO NOT apply less than 4 fl. oz./A and can only be applied by ground equipment.	DO NOT apply more than 16 fl. oz/A per application. DO NOT make more than 2 applications per season with a 21-day interval between applications. DO NOT exceed 32 fl. oz/A per growing season. Pre-Harvest Interval – 20 days
	Mid-season/Lay-by: 8.0-16.0	Apply this product when mites first appear. Rate will depend on the cotton plant size and thickness of foliage. Applications can be made by air or ground using a minimum of 5 gallons of water/A.	DO NOT graze livestock in treated fields or feed cotton foliage to livestock.

CUCURBIT VEGETABLE CROP GROUP (CROP GROUP 9)

Crops in this group include chayote, Chinese waxgourd, citron melon, cucumbers, gherkin, edible gourds (includes hyotan, cucuzza, hechima, Chinese okra); *Momordica* spp., (includes balsam apple, balsam pear, bittermelon, Chinese cucumber); muskmelon, (hybrids and/or cultivars of *Cucumis melo* includes true cantaloupe, cantaloupe, casaba, Crenshaw melon, golden pershaw melon, honeydew melon, honey balls, mango melon, Persian melon, pineapple melon, Santa Claus melon and snake melon), pumpkin, summer squash (includes crookneck squash, scallop squash, straightneck squash, vegetable marrow, zucchini), winter squash, (includes butternut squash, calabaza, Hubbard squash, acorn squash, spaghetti squash), watermelon (includes hybrids and/or varieties of *Citrullus lanatus*).

Target Insects	Application Rate of ABAMEX Miticide/ Insecticide fl. oz./A	Application Instructions and Timing	Restrictions
Leafminers Spider mites	Low to moderate pest infestations: 8.0-12.0 Heavy pest infestations: 16.0	Use ground equipment in 20 or more gal water/A. Although aerial applications are permitted when this product is applied in 5 or more gal water/A, better control of target insects is obtained using ground applications. Users accept all liability for level and duration of mite control from aerial applications of this product.	Pre-Harvest Interval – 7 days DO NOT exceed 48 fl. oz/A per growing season. DO NOT make more than 2 sequential applications per season. DO NOT apply by air in the State of New York.

(continued)

CUCURBIT VEGETABLE CROP GROUP (CROP GROUP 9) (continued)

Target Insects	Application Rate of ABAMEX Miticide/ Insecticide fl. oz./A	Application Instructions and Timing	Restrictions
		Apply when insects first appear. Increased water volumes are required under certain situations (severe pest infestations, highly dense foliage, elevated temperatures) to ensure thorough foliage coverage. If needed, a second application may be made but allow at least 7 days to elapse after the first application.	

DRY BEAN

* For use on cowpeas that are grown only for dry seed. Do not allow livestock to graze cowpea forage and do not harvest cowpea forage or hay for use as livestock feed.

Target Insects	Application Rate of ABAMEX Miticide/ Insecticide fl. oz./A	Application Instructions and Timing	Restrictions
<i>Liriomyza</i> leafminers Spider mites	Low to moderate pest infestations: 8.0-12.0 Heavy pest infestations: 13.0-16.0	Do not apply in less than 10 gal. of water/A with ground application equipment. Do not apply in less than 5 gal. of water/A with aircraft. Under conditions such as high pest populations, dense foliage, or adverse application conditions (such as high temperatures), use a greater volume of water to ensure adequate coverage. Leafminers and Spider mites: Apply when adult leafminers or spider mites first appear, and repeat as needed to maintain control within constraints of a sound resistance management program. If needed, a second application may be made but allow at least 6 days to elapse after the first application.	Pre-Harvest Interval – 7 days DO NOT exceed 16 fl. oz/A per application. DO NOT exceed 48 fl. oz/A per growing season. DO NOT make more than 2 sequential applications. Refer to additional directions on nonionic surfactants in the PRODUCT INFORMATION section. DO NOT apply by air in the State of New York.

FRUITING VEGETABLES CROP GROUP (CROP GROUP 8)

Crops in this group include eggplant, groundcherry, pepino, peppers (including bell pepper, chili pepper, cooking pepper, pimento, sweet pepper); tomatillo and tomato.

Target Insects	Application Rate of ABAMEX Miticide/Insecticide fl. oz./A	Application Instructions and Timing	Restrictions
Broad mites Colorado potato beetles <i>Liriomyza</i> leafminers Potato psyllids Spider mites Thrips palmi Tomato psyllids Tomato Russet mites	Low to moderate pest infestations: 8.0-12.0 Heavy pest infestations: 16.0 ADD a nonionic surfactant for best results	Use ground equipment in 20 or more gal water/A. Although aerial applications are permitted when this product is applied in 5 or more gal water/A, better control of target insects is obtained using ground applications. Users accept all liability for level and duration of mite control from aerial applications of this product. Apply when insects first appear. Increased water volumes are required under certain situations (severe pest infestations, highly dense foliage, elevated temperatures) to ensure thorough foliage coverage. If needed, a second application may be made but allow at least 7 days to elapse after the first application. A non-ionic surfactant (not binder or sticker-types) used in combination with this product provides improved wetting of foliage and allows an even deposition of the spray. Mites: For best control apply with ground equipment. Broad, russet, and spider mites: Apply when mites first appear. Thrips palmi: Apply when thrips are first observed. Leafminers: Apply when adult flies are first observed Potato Psyllids: Sequential application 10-14 days apart may be needed for infestations with heavy egg and nymph populations.	Pre-Harvest Interval – 7 days To avoid insect resistance to this product, do not apply to vegetables grown for transplant. DO NOT exceed 48 fl. oz/A per growing season. DO NOT make more than 2 sequential applications. Refer to additional directions on nonionic surfactants in the PRODUCT INFORMATION section. DO NOT apply by air in the State of New York. For resistance management, DO NOT use on fruiting vegetables grown for transplanting.
Tomato pinworms	16.0	Tomato pinworm: Application may be made from the time moth activity is detected up to, but no later than, the time when newly emerged larvae are present.	

**BERRY AND SMALL FRUIT
GRAPES**

Target Insects	Application Rate of ABAMEX Miticide/Insecticide fl. oz./A	Application Instructions and Timing	Restrictions
<p>Pacific spider mites Two-spotted spider mites Variegated leafhoppers* Western grapeleaf skeletonizers Western grape leafhoppers* Willamette spider mites</p> <p>*Note: This product only provides Knock-down control of pest.</p>	<p>Low to moderate pest infestations: 8.0-12.0 Heavy pest infestations: 16.0 ADD a nonionic surfactant for best results</p>	<p>Use conventional ground application equipment in 50 gal or more of water/A to ensure thorough foliage coverage, or with electrostatic spray equipment in 5 gal or more of water/A.</p> <p>Spider mites: Apply when mites first appear (or before more than 5 motiles per leaf are observed).</p> <p>Western grapeleaf skeletonizer: Apply with a non-ionic surfactant when the larvae are initially seen. For best results, apply shortly after eggs hatch.</p> <p>Leafhoppers: For contact knock-down control of these pests, apply with a non-ionic surfactant when pest first appears.</p> <p>European Grapevine Moth: Apply to aid in control of first generation of pest on grapes.</p> <p>Maximum coverage is required to control these pests, and this product must be applied to both sides of the rows. DO NOT spray alternate rows.</p> <p>If needed, a second application may be made but allow at least 21 days to elapse after the first application.</p> <p>A non-ionic (not binder or sticker-type) surfactant must be used in combination with this product to provide improved wetting of foliage and an even deposition of the spray.</p> <p>It is impossible to test the safety of this product plus non-ionic surfactants on all grape varieties under the various conditions which lead to phytotoxicity. Therefore carefully follow the directions for use on the nonionic surfactant label.</p>	<p>DO NOT enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours.</p> <p>Exception: For grape girdling cane turning, and tying in grapes. DO NOT enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 4 days.</p> <p>Pre-Harvest Interval – 28 days</p> <p>DO NOT exceed 16 fl. oz/A per application or 32 fl. oz/A per growing season.</p> <p>DO NOT make more than 2 applications per season.</p> <p>DO NOT graze livestock in treated vineyards.</p> <p>Refer to additional directions on nonionic surfactants in the PRODUCT INFORMATION section.</p> <p>DO NOT apply by air.</p>
<p>European Grapevine Moth (<i>Lobesia botrana</i>)</p>	<p>10-16</p>	<p>(This cell is merged with the previous row's application instructions.)</p>	<p>(This cell is merged with the previous row's restrictions.)</p>

HERB CROP SUBGROUP (CROP SUBGROUP 19A)

Crops in this group include angelica, balm, basil, borage, burnet, camomile (or chamomile), catnip, chervil (dried), chives, chives (Chinese), cilantro (leaf), clary, coriander (leaf), costmary, culantro (leaf), curry (leaf), dill weed, horehound, hyssop, lavender, lemongrass, lovage (leaf), marigold, marjoram (*Origanum* species), nasturtium, parsley (dried), pennyroyal, rosemary, rue, sage, summer and winter savory, sweet bay, tansy, tarragon, thyme, wintergreen, woodruff, and wormwood.

Target Insects	Application Rate of ABAMEX Miticide/ Insecticide fl. oz./A	Application Instructions and Timing	Restrictions
<p><i>Liriomyza</i> leafminers Spider mites</p>	<p>Low to moderate pest infestations: 8.0 Heavy pest infestations: 16.0 ADD a nonionic surfactant for best results</p>	<p>Apply using ground equipment using conventional dilute or concentrate sprayers. Apply when adult leafminer flies first appear using sufficient volumes (greater than 30 gal of water per acre) to ensure thorough foliage coverage. If needed, a second application may be made but allow at least 7 days to elapse after the first application. A non-ionic surfactant (not binder or sticker-types) used in combination with this product provides improved wetting of foliage and allows an even deposition of the spray on the foliage.</p>	<p>Pre-Harvest Interval for listed herbs except chive– 14 days. Pre-Harvest Interval for chives - 7 days. DO NOT apply more than 6 applications of 8 fl. oz. of this product/A or 3 applications of 16 fl. oz. of this product/A per cropping season. DO NOT make more than 2 applications per single cutting (harvest). DO NOT apply by air. Refer to additional directions on nonionic surfactants in the PRODUCT INFORMATION section. DO NOT graze livestock in treated fields or feed treated foliage to livestock.</p>

HOPS (EXCLUDING CALIFORNIA)

Target Insects	Application Rate of ABAMEX Miticide/ Insecticide fl. oz./A	Application Instructions and Timing	Restrictions
Two-spotted spider mites	8.0 (minimum)-16.0 ADD a nonionic surfactant for best results	<p>Use conventional ground application as dilute or concentrate sprays. Use sufficient volumes (greater than 40 gal of water/A) to ensure thorough foliage coverage. If trellis height is 6-8 feet, use 40 gal of water/A at the 8-16 fl. oz/A rate; if height is greater than 6-8 feet, use 100 gal/A at the 16 fl. oz/A rate.</p> <p>Note that while concentrate sprays DO NOT require as much water, the amount of this product/A should be the same as for dilute sprays.</p> <p>Apply when mites first appear. If needed, a second application may be made but allow at least 21 days to elapse after the first application and repeat application only after using an alternative miticide with a different mode of action.</p> <p>A non-ionic surfactant (not binder or sticker-types) used in combination with this product provides improved wetting of foliage and allows an even deposition of the spray on the foliage.</p>	<p>Pre-Harvest Interval – 28 days DO NOT exceed 16 fl. oz/A per application or 32 fl. oz/A per growing season.</p> <p>DO NOT apply by air. DO NOT apply more than 2 applications per growing season. Refer to additional directions on nonionic surfactants in the PRODUCT INFORMATION section. DO NOT graze livestock in treated hop yards.</p>

LEAFY VEGETABLE (Except Brassica Vegetables) CROP GROUP (CROP GROUP 4)

Crops in this group include amaranth (leafy amaranth, Chinese spinach, tampala), arugula (rocket), cardoon, celery, celtuce, chervil, Chinese celery, chrysanthemum (edible leaved and garland), corn salad, cress (garden and upland (yellow rocket and winter cress)), dandelion, dock (sorrel), endive (escarole), fennel, Florence fennel, lettuce (head and leaf), New Zealand spinach, orach, parsley, purslane (garden and winter), radicchio (red chicory), rhubarb, spinach, Swiss chard and vine spinach.

Target Insects	Application Rate of ABAMEX Miticide/ Insecticide fl. oz./A	Application Instructions and Timing	Restrictions
Carmine spider mites <i>Liriomyza</i> leafminers Two-spotted spider mites	Low to moderate pest infestations: 8.0-12.0 Heavy pest infestations: 16.0 ADD a nonionic surfactant for best results	Apply this product when mites or adult leafminer flies are observed and use ground equipment in 20 or more gal water/A. Although aerial applications are permitted when this product is applied in 5 or more gal water/A, better control of target insects is obtained using ground applications. Users accept all liability for level and duration of insect control from aerial applications of this product. Spider mites: Apply when mites first appear, and repeat as needed to maintain control. Leafminers: Apply when adult flies are first observed and repeat applications as needed to maintain control. Increased water volumes are required under certain situations (severe pest infestations, highly dense foliage, elevated temperatures) to ensure thorough foliage coverage. If needed, a second application may be made but allow at least 7 days to elapse after the first application. A non-ionic surfactant (not binder or sticker-types) used in combination with this product provides improved wetting of foliage and allows an even deposition of the spray on the foliage.	Pre-Harvest Interval – 7 days DO NOT exceed 48 fl. oz/A per growing season. To avoid insect resistance to this product, do not apply to vegetables grown for transplants. DO NOT make more than 2 sequential applications. Refer to additional directions on nonionic surfactants in the PRODUCT INFORMATION section. DO NOT apply by air in the State of New York.

MINT (PEPPERMINT AND SPEARMINT)

Target Insects	Application Rate of ABAMEX Miticide/Insecticide fl. oz./A	Application Instructions and Timing	Restrictions
Two-spotted spider mites	8.0-12.0 ADD a nonionic surfactant for best results	<p>Apply this product when mites first appear and use ground equipment in 20 or more gal water/A. Although aerial applications are permitted when this product is applied in 5 or more gal water/A, better control of target insects is obtained using ground applications. Users accept all liability for level and duration of insect control from aerial applications of this product.</p> <p>Increased water volumes are required under certain situations (severe pest infestations, highly dense foliage, elevated temperatures) to ensure thorough foliage coverage.</p> <p>If needed, a second application may be made but allow at least 7 days to elapse after the first application.</p> <p>A non-ionic surfactant (not binder or sticker-types) used in combination with this product provides improved wetting of foliage and allows an even deposition of the spray on the foliage.</p>	<p>Pre-Harvest Interval – 28 days</p> <p>DO NOT exceed 12 fl. oz./A per application or 36 fl. oz./A per growing season.</p> <p>DO NOT make more than 2 sequential applications.</p> <p>DO NOT apply by air in the State of New York.</p> <p>Refer to additional directions on nonionic surfactants in the PRODUCT INFORMATION section.</p> <p>DO NOT graze livestock in treated fields or feed treated foliage to livestock.</p>

ONION, BULB (CROP SUBGROUP 3-07A)

Crops in this group are bulb onion, including bulb Chinese, bulb daylily, bulb fritillaria, bulb garlic, bulb great-headed garlic, bulb lily, bulb onion, pearl onion, potato onion, bulb serpent garlic, bulb shallot, cultivars, varieties, and/or hybrids of bulb onion, subgroup 3-07A.

Target Insects	Application Rate of ABAMEX Miticide/Insecticide fl. oz./A	Application Instructions and Timing	Restrictions
<p><i>Liriomyza</i> leafminers Thrips</p>	<p>Low to moderate pest infestations: 8.0-12.0 Heavy pest infestations: 13.0-16.0</p>	<p>Do not apply in less than 20 gal. of water/A with ground application equipment. Do not apply in less than 5 gal. of water/A with aircraft. Under conditions such as high pest populations, dense foliage, or adverse application conditions (such as high temperatures), use a greater volume of water to ensure adequate coverage.</p> <p>Leafminers: Apply when adult leafminers first appear, and repeat as needed to maintain control within constraints of a sound resistance management program.</p> <p>Thrips: Apply in a thrips management program and when thrips are at economic threshold. Repeat application, as needed, to maintain control within constraints of a sound resistance management program.</p> <p>Do not use this product as a rescue treatment for thrips control.</p> <p>If needed, a second application may be made but allow at least 7 days to elapse after the first application.</p> <p>Make 2 consecutive applications or then rotate to a chemistry with a different mode of action. Make at least 2 applications of a chemistry with a different mode of action before making additional applications of this product.</p>	<p>Pre-Harvest Interval – 30 days DO NOT exceed 16 fl. oz/A per application. DO NOT exceed 48 fl. oz/A per growing season. DO NOT make more than 2 sequential applications. Refer to additional directions on nonionic surfactants in the PRODUCT INFORMATION section. DO NOT apply by air in the State of New York.</p>
<p>Note: Add a non-ionic activator type wetting, spreading and/or penetrating adjuvant approved for use on bulb onion. Do not use binder sticker type adjuvants. Tank mixing with products that have binder sticker type formulations can reduce performance of this product.</p>			

STONE FRUIT CROP GROUP (CROP GROUP 12)

Crops in this group include apricot, cherry (sweet and tart), nectarine, peach, plum, plum (chickasaw, damson, and Japanese), plumcot and prune (fresh).

Target Insects	Application Rate of ABAMEX Miticide/Insecticide fl. oz./A (fl. oz./100 gal)	Application Instructions and Timing	Restrictions
European red mites Pacific spider mites Two-spotted spider mites	Concentrate Sprays: 10-20 fl. oz/A (Dilute Sprays: 2.5-5.0 fl. oz./100 gal) PLUS a horticultural oil or nonionic surfactant (required for optimum insect control)	Apply using conventional dilute or concentrate ground equipment. Make applications when mites first are seen. Use sufficient volumes (greater than 40 gal of water/A) to ensure thorough foliage coverage. If needed, a second application may be made but allow at least 21 days to elapse after the first application. Always apply this product in combination with either a nonionic surfactant, or with horticultural spray oil (not a dormant oil), but use caution since oil may cause phytotoxicity to foliage and fruit.	Pre-Harvest Interval – 21 days DO NOT exceed 20 fl. oz/A per application or 40 fl. oz/A per growing season. DO NOT make more than 2 applications per season. Refer to additional directions on spray oils and nonionic surfactants in the PRODUCT INFORMATION section. See specific restrictions for this crop in the SPRAY DRIFT MANAGEMENT section of this label. DO NOT apply by air. DO NOT graze livestock in treated orchards. Horticulture Spray Oil (NOT a dormant oil) 1 gal/A (minimum) (0.25%, v/v) Nonionic Surfactant – labeled rate

NOTE: Concentrate Spray rates (ABAMEX Miticide/Insecticide/A) are determined by selecting the amount needed for a full cover dilute spray. The amount of ABAMEX Miticide/Insecticide/A as a concentrate spray is the same amount as required for dilute sprays to the orchard and could be less than 10 fl. oz. ABAMEX Miticide/Insecticide/A for small trees. Rates of ABAMEX Miticide/Insecticide/100 gal are based on 400 gal/A of dilute spray.

NOTE:

- 1) A nonionic surfactant that spreads on and/or penetrates the leaf cuticle can improve insect control. It is impossible to test on all stone fruit varieties under the variety of conditions that may cause crop injury. Therefore, when using ABAMEX Miticide/Insecticide in combination with a non-ionic surfactant, carefully follow the **Directions for Use** and **Precautions** on the surfactant label and in official spray guides
- 2) A horticultural oil (not a dormant oil) may improve efficacy and may be used but it may increase the possibility of phytotoxicity to foliage and fruit.

**BERRY AND SMALL FRUIT
STRAWBERRIES**

Target Insects	Application Rate of ABAMEX Miticide/ Insecticide fl. oz./A	Application Instructions and Timing	Restrictions
<p>Aphids Carmine mites Cyclamen mites Strawberry spider mites Thrips Two-spotted spider mites Whiteflies</p>	<p align="center">16.0</p>	<p>Apply using conventional ground equipment or electrostatic sprayers.</p> <p>Optimum coverage of the leaves (tops and bottoms) can be achieved by adjusting the spray volumes and placement of the nozzles.</p> <p>Apply when mites first appear using sufficient volumes (greater than 50 gal of water/A for conventional sprayers, greater than 10 gal of water/A for electrostatic sprayers) to ensure thorough foliage coverage.</p> <p>Make a second application at 7-10 day intervals.</p> <p>Repeat the two step application process, if needed, but only 21-days after the second application is made.</p> <p>A non-ionic or silicone-based (not a sticker or binder type) surfactant used in combination with this product provides improved wetting of foliage and allows an even deposition of the spray.</p>	<p>Pre-Harvest Interval – 3 days</p> <p>To avoid insect resistance to this product, do not apply to strawberries grown in nurseries.</p> <p>DO NOT exceed 16 fl. oz/A per application or more than 64 fl. oz./A per growing season.</p> <p>Refer to additional directions on nonionic surfactants in the PRODUCT INFORMATION section.</p> <p>DO NOT apply by air.</p>

TREE NUTS CROP GROUP (CROP GROUP 14) AND PISTACHIO

Crops in this group include almond, beech nut, Brazil nut, butternut, cashew, chestnut, chinquapin, filbert (hazelnut), hickory nut, macadamia nut, pecan, and walnuts (black and English).

Target Insects	Application Rate of ABAMEX Miticide/Insecticide fl. oz./A (fl. oz./100 gal)	Application Instructions and Timing	Restrictions
<p>European red mites Pacific spider mites Strawberry spider mites Two-spotted spider mites</p>	<p>Concentrate Sprays: 10-20 fl. oz/A (Dilute Sprays: 2.5-5.0 fl. oz./100 gal)</p> <p>PLUS</p> <p>a horticultural spray oil (required for optimum insect control)</p>	<p>Apply using conventional dilute or concentrate ground equipment.</p> <p>Use sufficient volumes (greater than 40 gal of water/A) to ensure thorough foliage coverage.</p> <p>Apply this product when spider mites first appear. Greater residual spider mite control is achieved when spray is deposited on new foliage growth.</p> <p>If needed, a second application may be made but allow at least 21 days to elapse after the first application.</p> <p>To manage resistance, if pest infestation requires additional treatments, use other alternative insecticides with different modes of action which are labeled for use on these pests and crops.</p>	<p>Pre-Harvest Interval – 21 days</p> <p>DO NOT exceed 20 fl. oz/A per application or 40 fl. oz/A per growing season.</p> <p>DO NOT make more than 2 applications per season.</p> <p>DO NOT apply by air.</p> <p>Refer to additional directions on spray oils in the PRODUCT INFORMATION section.</p> <p>See specific restrictions for this crop in the SPRAY DRIFT MANAGEMENT section of this label.</p> <p>DO NOT graze livestock in treated orchards.</p> <p>Horticultural Spray Oil (DO NOT use a dormant oil) 1 gal./A (minimum) (0.25%, v/v)</p>
<p>NOTE: Concentrate Spray rates (ABAMEX Miticide/Insecticide/A) are determined by selecting the amount needed for a full cover dilute spray. The amount of ABAMEX Miticide/Insecticide/A as a concentrate spray is the same amount as required for dilute sprays to the orchard and could be less than 10 fl. oz. ABAMEX Miticide/Insecticide/A for small trees. Rates of ABAMEX Miticide/Insecticide/100 gal are based on 400 gal/A of dilute spray.</p>			

TUBEROUS AND CORM VEGETABLES CROP SUBGROUP (CROP SUBGROUP 1 C)

Crops in this group include arracacha, arrowroot, artichoke, Chinese artichoke, edible canna, Jerusalem artichoke, cassava (bitter and sweet), chayote (root), chufa, dasheen, ginger, leren, potato, sweet potato, tanager, turmeric, yam bean, and yam (true).

Target Insects	Application Rate of ABAMEX Miticide/Insecticide fl. oz./A	Application Instructions and Timing	Restrictions
<p>Colorado potato beetles <i>Liriomyza</i> leafminers Potato psyllids Spider mites</p>	<p>Low to moderate pest infestations: 8.0-12.0 Heavy pest infestations: 16.0</p>	<p>Apply using conventional ground or aerial application equipment.</p> <p>Apply when mites and adult leafminer flies first appear. Use sufficient volumes (greater than 20 gal of water/A for ground applications and greater than 5 gal of water/A for aerial applications) to ensure thorough foliage coverage.</p> <p>Colorado potato beetle: Begin applications when 50% of the egg masses have hatched or larvae are observed. If a second application is made, ensure it is to control a single Colorado potato beetle generation.</p> <p><i>Liriomyza</i> leafminers: Make the first application when adult flies are first observed. Repeat applications as needed to maintain control.</p> <p>Spider mites: Make the first application when mites first appear. Repeat application as needed to maintain control.</p> <p>If needed, a second application may be made but allow at least 7 days to elapse after the first application.</p> <p>A non-ionic or silicone-based (not a sticker or binder type) surfactant used in combination with this product provides improved wetting of foliage and allows an even deposition of the spray.</p>	<p>Pre-Harvest Interval – 14 days DO NOT exceed 32 fl. oz/A per crop season for control of spider mites or Colorado potato beetle.</p> <p>DO NOT exceed 48 fl. oz/A per crop season for control of leafminers.</p> <p>DO NOT make more than 2 applications to control Colorado potato beetle per crop season.</p> <p>DO NOT make more than two sequential applications.</p> <p>DO NOT apply by air in the State of New York.</p> <p>Refer to additional directions on nonionic surfactants in the PRODUCT INFORMATION section.</p> <p>DO NOT graze or feed treated foliage to livestock.</p>

STORAGE AND DISPOSAL

DO NOT contaminate water, food, or feed by storage or disposal.

PESTICIDE STORAGE: Store in a tightly closed container in a cool, dry place.

PESTICIDE DISPOSAL: Improper disposal of excess pesticide spray mixture or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL: Nonrefillable container. **DO NOT** reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. **Triple rinse as follows:** Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

SPILLS: For minor spills, leaks, etc., follow all precautions indicated on this label and clean up immediately. Take special care to avoid contamination of equipment and facilities during cleanup procedures and disposal of wastes. In the event of a major spill, fire or other emergency, call 1-800-424-9300, day or night.

WARRANTY DISCLAIMER

The directions for use of this product must be followed carefully. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, (1) THE GOODS DELIVERED TO YOU ARE FURNISHED "AS IS" BY MANUFACTURER OR SELLER AND (2) MANUFACTURER AND SELLER MAKE NO WARRANTIES, GUARANTEES, OR REPRESENTATIONS OF ANY KIND TO BUYER OR USER, EITHER EXPRESS OR IMPLIED, OR BY USAGE OF TRADE, STATUTORY OR OTHERWISE, WITH REGARD TO THE PRODUCT SOLD, INCLUDING, BUT NOT LIMITED TO MERCHANTABILITY, FITNESS FOR A PARTICULAR PURPOSE, USE, OR ELIGIBILITY OF THE PRODUCT FOR ANY PARTICULAR TRADE USAGE. UNINTENDED CONSEQUENCES, INCLUDING BUT NOT LIMITED TO INEFFECTIVENESS, MAY RESULT BECAUSE OF SUCH FACTORS AS THE PRESENCE OR ABSENCE OF OTHER MATERIALS USED IN COMBINATION WITH THE GOODS, OR THE MANNER OF USE OR APPLICATION, INCLUDING WEATHER, ALL OF WHICH ARE BEYOND THE CONTROL OF MANUFACTURER OR SELLER AND ASSUMED BY BUYER OR USER. THIS WRITING CONTAINS ALL OF THE REPRESENTATIONS AND AGREEMENTS BETWEEN BUYER, MANUFACTURER AND SELLER, AND NO PERSON OR AGENT OF MANUFACTURER OR SELLER HAS ANY AUTHORITY TO MAKE ANY REPRESENTATION OR WARRANTY OR AGREEMENT RELATING IN ANY WAY TO THESE GOODS.

LIMITATION OF LIABILITY

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO EVENT SHALL MANUFACTURER OR SELLER BE LIABLE FOR SPECIAL, INCIDENTAL, OR CONSEQUENTIAL DAMAGES, OR FOR DAMAGES IN THEIR NATURE OF PENALTIES RELATING TO THE GOODS SOLD, INCLUDING USE, APPLICATION, HANDLING, AND DISPOSAL. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, MANUFACTURER OR SELLER SHALL NOT BE LIABLE TO BUYER OR USER BY WAY OF INDEMNIFICATION TO BUYER OR TO CUSTOMERS OF BUYER, IF ANY, OR FOR ANY DAMAGES OR SUMS OF MONEY, CLAIMS OR DEMANDS WHATSOEVER, RESULTING FROM OR BY REASON OF, OR ARISING OUT OF THE MISUSE, OR FAILURE TO FOLLOW LABEL WARNINGS OR INSTRUCTIONS FOR USE, OF THE GOODS SOLD BY MANUFACTURER OR SELLER TO BUYER. ALL SUCH RISKS SHALL BE ASSUMED BY THE BUYER, USER, OR ITS CUSTOMERS. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BUYER'S OR USER'S EXCLUSIVE REMEDY, AND MANUFACTURER'S OR SELLER'S TOTAL LIABILITY SHALL BE FOR DAMAGES NOT EXCEEDING THE COST OF THE PRODUCT.

If you do not agree with or do not accept any of directions for use, the warranty disclaimers, or limitations on liability, do not use the product, and return it unopened to the Seller, and the purchase price will be refunded.

RV100115

NOTES

RESTRICTED USE PESTICIDE
TOXIC TO FISH, MAMMALS, AND AQUATIC ORGANISMS
FOR RETAIL SALE TO AND USE ONLY BY CERTIFIED APPLICATORS OR
PERSONS UNDER THEIR DIRECT SUPERVISION, AND ONLY FOR THOSE
USES COVERED BY THE CERTIFIED APPLICATOR'S CERTIFICATION.

GROUP 6 INSECTICIDE

Abamex™
Miticide/Insecticide

ACTIVE INGREDIENT:

Abamectin (CAS No.71751-41-2): 2.0%*

OTHER INGREDIENTS: 98.0%

TOTAL: 100.0%

*Contains 0.15 pounds Abamectin per gallon.

Contains VOC ≤20% EP; qualifies as "Low VOC" under California regulations

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

Si usted no entiende la etiqueta, busque a alguien para que se la
explique a usted en detalle. (If you do not understand the label, find
someone to explain it to you in detail.)

**SEE INSIDE BOOKLET FOR ADDITIONAL FIRST AID
AND PRECAUTIONARY STATEMENTS**

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

Harmful if swallowed. Causes moderate eye irritation. Avoid contact with
eyes or clothing. Wash thoroughly with soap and water after handling and
before eating, drinking, chewing gum, using tobacco or using the toilet.

FIRST AID

IF SWALLOWED: Call a poison control center or doctor immediately
for treatment advice. **DO NOT** give any liquid to the person. **DO NOT**
induce vomiting unless told to do so by the poison control center or
doctor. **DO NOT** give anything by mouth to an unconscious person. **IF IN
EYES:** Hold eye open and rinse slowly and gently with water for 15-20
minutes. Remove contact lenses, if present, after the first 5 minutes,
then continue rinsing eye. Call a poison control center or doctor for
treatment advice. **IF ON SKIN OR CLOTHING:** Take off contaminated
clothing. Rinse skin immediately with plenty of water for 15-20
minutes. Call a poison control center or doctor for treatment advice. **IF
INHALED:** Move person to fresh air. If person is not breathing, call 911
or an ambulance, then give artificial respiration, preferably by mouth-
to-mouth, if possible. Call a poison control center or doctor for further
treatment advice. **HOT LINE NUMBER:** Have the product container or

FIRST AID (continued)

label with you when calling a poison control center or doctor, or going
for treatment. You may also contact 1-877-325-1840 for emergency
medical treatment information. **NOTE TO PHYSICIAN:** Early signs of
intoxication include dilating of pupils, muscular incoordination, and
muscle tremors. Toxicity following accidental ingestion of this product
can be minimized by early administration of chemical adsorbents (e.g.,
activated charcoal). If toxicity from exposure has progressed to cause
severe vomiting, the extent of resultant fluid and electrolyte imbalance
should be gauged. Appropriate supportive parenteral fluid replacement
therapy should be given, along with other required supportive
measures (such as maintenance of blood pressure levels and proper
respiratory functionality) as indicated by clinical signs, symptoms,
and measurements. In severe cases, observations should continue
for at least several days until clinical condition is stable and normal.
Since abamectin is believed to enhance GABA activity in animals, it is
probably wise to avoid drugs that enhance GABA activity (barbiturates,
benzodiazepines, valproic acid) in patients with potentially toxic
abamectin exposure.

STORAGE AND DISPOSAL

DO NOT contaminate water, food, or feed by storage or disposal.
PESTICIDE STORAGE: Store in a tightly closed container in a cool,
dry place.

PESTICIDE DISPOSAL: Improper disposal of excess pesticide spray
mixture or rinsate is a violation of Federal law. If these wastes cannot
be disposed of by use according to label instructions, contact your
State Pesticide or Environmental Control Agency or the Hazardous
Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL: Nonrefillable container. DO NOT reuse or
refill this container. Triple rinse container (or equivalent) promptly after
emptying. **Triple rinse as follows:** Empty the remaining contents into
application equipment or a mix tank and drain for 10 seconds after
the flow begins to drip. Fill the container ¼ full with water and recap.
Shake for 10 seconds. Pour rinsate into application equipment or a mix
tank or store rinsate for later use or disposal. Drain for 10 seconds
after the flow begins to drip. Repeat this procedure two more times.
Then offer for recycling or reconditioning, or puncture and dispose of in
a sanitary landfill, or by other procedures approved by State and local
authorities. Plastic containers are also disposable by incineration, or, if
allowed by State and local authorities, by burning. If burned, stay out of
smoke. **SPILLS:** For minor spills, leaks, etc., follow all precautions
indicated on this label and clean up immediately. Take special care
to avoid contamination of equipment and facilities during cleanup
procedures and disposal of wastes. In the event of a major spill, fire
or other emergency, call 1-800-424-9300, day or night.

EPA REG. NO. 228-734
EPA EST. NO. 70815-GA-001

**Net Contents:
1 Gal. (3.78 L)**

Manufactured for
Nufarm Americas Inc.
11901 S. Austin Avenue
Alsip, IL 60803

PULL HERE TO OPEN
PRESS TO RESEAL

14930000

RV100115