



# Millennium ULTRA<sup>2</sup>

Department of Agriculture  
STATE OF HAWAII

**LICENSED**

For selective broadleaf weed control in ornamental lawns, turf grasses, and on sod farms

**CONTAINS 2,4-D, CLOPYRALID AND DICAMBA**

**CONTROLS:** Dandelion, Chickweed, Black medic, Knotweed, Plantain, Oxalis, Clover, Cocklebur, Thistle and many other species of broadleaf weeds; some of which are listed on this label.

PERIOD 2014-2016 LIC. NO.

**9805.32**

**ACTIVE INGREDIENTS:**

Dimethylamine Salt of 2,4-Dichlorophenoxyacetic Acid* .....	37.32%
Monoethanolamine Salt of 3,6-Dichloro-2-Pyridinecarboxylic Acid** .....	2.54%
Dimethylamine Salt of Dicamba (3,6-Dichloro-o-Anisic Acid)*** .....	4.65%

**OTHER INGREDIENTS:**..... 55.49%

**TOTAL** ..... 100.00%

Isomer Specific Method, Equivalent to:

*2,4-Dichlorophenoxyacetic Acid.....	31.00%, 3.000 lbs./gal.
**3,6-Dichloro-2-Pyridinecarboxylic Acid.....	1.93%, 0.183 lbs./gal.
***3,6-Dichloro-o-Anisic Acid .....	3.86%, 0.375 lbs./gal.

**NOT FOR SALE OR USE IN NASSAU AND SUFFOLK COUNTIES IN N.Y. STATE. IN CALIFORNIA AND WASHINGTON, TURFGRASS AND LAWN USES ARE RESTRICTED TO GOLF COURSES ONLY. IN OREGON, THE USE OF THIS PRODUCT IS LIMITED TO THE SITES STATED ON THIS LABEL WHICH ARE AGRICULTURAL, RIGHTS-OF-WAY, GOLF COURSE OR CEMETERY SITES.**

**KEEP OUT OF REACH OF CHILDREN**

**DANGER / PELIGRO**

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

SEE INSIDE BOOKLET FOR FIRST AID AND ADDITIONAL  
PRECAUTIONARY STATEMENTS

For Chemical Spill, Leak, Fire, or  
Exposure, Call CHEMTREC  
(800) 424-9300

For Medical Emergencies Only,  
Call (877) 325-1840

Manufactured for  
Nufarm Americas Inc.  
150 Harvester Drive  
Burr Ridge, IL 60527

EPA Reg. No. 228-332

EPA Est. No. 228-IL-1



7 3621176296 2

**Net Contents**  
**2.5 Gal.**  
**(9.46 L)**

**Nonrefillable Container**

**14300000**





**PRECAUTIONARY STATEMENTS  
HAZARDS TO HUMANS AND DOMESTIC ANIMALS  
DANGER / PELIGRO**

Corrosive. Causes irreversible eye damage. Harmful if swallowed or absorbed through skin. Do not get in eyes, on skin or on clothing.

**PERSONAL PROTECTIVE EQUIPMENT (PPE):**

Some materials that are chemical-resistant to this product are made of any waterproof material. If you want more options, follow the instructions for category A on an EPA chemical-resistance category selection chart.

**All mixers, loaders, applicators, and other handlers must wear:**

- long-sleeved shirt and long pants,
- shoes and socks, plus
- goggles or face shield \*

\* After this product has been diluted with water in accordance with Directions for Use, safety glasses for protective eyewear can be worn.

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

This product is not to be applied through any type of aerial application equipment.

**USER SAFETY RECOMMENDATIONS**

**Users Should:**

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. If pesticide gets on skin, wash immediately with soap and water.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

**FIRST AID**

**IF IN EYES**

- Hold eye open and rinse slowly and gently with water for 15 to 20 minutes.
- Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.
- Call a poison control center or doctor for treatment advice.

**IF SWALLOWED**

- Call a poison control center or doctor immediately for treatment advice.
- Have person sip a glass of water if able to swallow.
- Do not induce vomiting unless told to do so by the poison control center or doctor.
- Do not give anything by mouth to an unconscious person.

**IF ON SKIN  
OR CLOTHING**

- Take off contaminated clothing.
- Rinse skin immediately with plenty of water for 15 to 20 minutes.
- Call a poison control center or doctor for treatment advice.

**HOT LINE NUMBER**

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-877-325-1840 for emergency medical treatment information.

**NOTE TO PHYSICIAN**

Probable mucosal damage may contraindicate the use of gastric lavage.

**ENVIRONMENTAL HAZARDS**

This pesticide may be toxic to fish and aquatic invertebrates. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas. Do not contaminate water when disposing of equipment wash waters or rinsate.

This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistern or well may result in contamination of drinking water or groundwater.

Clopyralid, an active ingredient in this product, may leach through soil into groundwater under certain conditions of use. Use of this product where soils are permeable, particularly where the water table is shallow, may result in leaching to groundwater. Caution should be exercised when handling this product at mixing, loading, and disposal sites to prevent contamination of groundwater supplies. Use of closed systems for mixing and transferring this pesticide will reduce the probability of spills. Placement of the mixing/loading equipment on an impervious pad to contain spills will help prevent groundwater contamination.





## DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. READ ENTIRE LABEL BEFORE USING THIS PRODUCT. USE STRICTLY IN ACCORDANCE WITH LABEL PRECAUTIONARY STATEMENTS AND DIRECTIONS.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the State or Tribal agency responsible for pesticide regulation.

### AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about Personal Protective Equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 48 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls worn over short-sleeve shirt and short pants,
- Chemical-resistant footwear plus socks,
- Chemical-resistant gloves made of any waterproof material,
- Chemical-resistant headgear for overhead exposure, and
- Protective eyewear.

### NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Do not enter or allow people (or pets) to enter the treated area until sprays have dried.

## USE PRECAUTIONS

This product is for use on Non-Residential Ornamental Turf Lawns (Industrial and Institutional) Athletic/Recreational Fields (Parks, Playgrounds and Sports Fields) Cemeteries, Sod Farms, Golf Courses (Fairways, Aprons, Tees\* and Roughs), Industrial Sites, Non-Cropland, Rights-of-Way and Roadsides. \*Excluding Bentgrass Tees.

Do not use on residential turfgrass. Do not collect grass clippings for mulch or compost. Do not send clippings to a compost facility. Applicator must give notice to landowners/property managers to not use grass clippings for mulching or composting.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application.

## SPRAY DRIFT MANAGEMENT

A variety of factors including weather conditions (e.g., wind direction, wind speed, temperature, relative humidity) and method of application (e.g., ground, aerial, airst, chemigation) can influence pesticide drift. The applicator must evaluate all factors and make appropriate adjustments when applying this product.

### Droplet Size

When applying sprays that contain 2,4-D as the sole active ingredient, or when applying sprays that contain 2,4-D mixed with active ingredients that require a coarse or coarser spray, apply only as a coarse or coarser spray (ASAE standard 572) or a volume mean diameter of 385 microns or greater for spinning atomizer nozzles.

When applying sprays that contain 2,4-D mixed with other active ingredients that require a medium or more fine spray, apply only as a medium or coarser spray (ASAE standard 572) or a volume mean diameter of 300 microns or greater for spinning atomizer nozzles.

### Wind Speed

Do not apply at wind speeds greater than 15 mph. Only apply this product if the wind direction favors on-target deposition and there are no sensitive areas (including, but not limited to, residential areas, bodies of water, known habitat for nontarget species, nontarget crops) within 250 feet downwind. If applying a medium spray, leave one swath unsprayed at the downwind edge of the treated field.

### Temperature Inversions

If applying at wind speeds less than 3 mph, the applicator must determine if: a) conditions of temperature inversion exist, or b) stable atmospheric conditions exist at or below nozzle height. Do not make applications into areas of temperature inversions or stable atmospheric conditions.





### Susceptible Plants

Do not apply under circumstances where spray drift may occur to food, forage, or other plantings that might be damaged or crops thereof rendered unfit for sale, use or consumption. Susceptible crops include, but are not limited to, cotton, okra, flowers, grapes (in growing stage), fruit trees (foliage), soybeans (vegetative stage), ornamentals, sunflowers, tomatoes, beans, and other vegetables, or tobacco. Small amounts of spray drift that might not be visible may injure susceptible broadleaf plants.

### Other State and Local Requirements

Applicators must follow all state and local pesticide drift requirements regarding application of 2,4-D herbicides. Where states have more stringent regulations, they must be observed.

### Equipment

All aerial and ground application equipment must be properly maintained and calibrated using appropriate carriers or surrogates.

Do not apply with a nozzle height greater than 4 feet above the crop canopy.

Avoid drift of spray mist to vegetables, flowers, ornamental plants, shrubs, trees and other desirable plants. Do not pour spray solutions near desirable plants. Do not use on Centipedegrass, St. Augustinegrass, Dichondragrass, nor on lawns or turf where desirable clovers are present. Avoid fine mists. Use lawn type sprayer with coarse spray as wind drift is less likely. Avoid contact with exposed feeder roots of ornamentals and trees. Maximum control of weeds will be obtained from spring or early fall applications when weeds are actively growing. Use the higher rate for hard-to-control weeds. Do not exceed specified dosages for any area; be particularly careful within the dripline of tree and other ornamental species. Do not apply to newly seeded grasses until well established. Do not use grass clippings from turf treated with this product for composting and/or mulch. Avoid broadcast applications when air temperature exceeds 90°F. When using small, spot treatment applications in temperature over 90°F, turf injury may occur.

When treating Carpetgrass and St. Augustine grass, avoid broadcast applications when air temperature exceeds 80°F. When air temperatures exceed 80°F, limit application to spot treatment only.

The suitable use of this product on non-recommended turf species may be determined by treating a small area at any rate/acre which does not exceed 3 pints/acre. The treated area should be observed for any sign of turf injury for a period of 30 days of normal growing conditions to determine the phytotoxicity and efficacy to the treated area.

For optimum results: (1) avoid applying during excessively dry or hot periods unless irrigation is used; (2) turf should not be mowed 1 to 2 days before and following application; (3) reseed no sooner than 3 to 4 weeks after application of this product. Adding oil, wetting agent, or other surfactant to the spray may be used to increase effectiveness on weeds but doing so may reduce selectivity to turf resulting in turf damage. Clean and rinse spray equipment using soap or detergent and water, and rinse thoroughly before reuse for other sprays.

## COMPATIBILITY

This product can be mixed with some liquid fertilizers or liquid iron materials. Because liquid fertilizers and liquid iron differ in pH, free ammonia content, density, salt concentration and percentage of water, a compatibility test (given below) is recommended prior to mixing in the application equipment. All regulations, either State or Federal, relating to the application of liquid fertilizers or liquid iron and this product must be strictly followed. The following compatibility test should always be performed prior to full-scale mixing.

1. Pour 18 fluid ounces of water into a quart jar.
2. Add 1 fluid ounce of either the liquid fertilizer or liquid iron to be used.
3. Add 1 fluid ounce of this product.
4. Close jar and shake well.
5. Watch the mixture for several seconds after shaking and check again after 30 minutes.
6. If the mixture does not show signs of separating, the combination may be used. If the mixture foams excessively, gels, separates or gets very thick, do not combine for field application.
7. Compatibility may be improved by the use of a compatibility agent. Follow the previously outlined test procedures and add 1/6 fluid ounce of the compatibility agent between steps (the compatibility agent must be added to the fertilizer or iron before adding this product).
8. If the mixture does not separate, gel, foam or get very thick, it may be used for field application. Mix only the amount to be sprayed. Do not allow to stand overnight.





## WEEDS CONTROLLED

Alder	Clover, white	Henbit	Oxalis	Spiny amaranth
Artichoke	Cockle	Hoary cress	Pansy, field	Spurge
Aster	Cocklebur	Honeysuckle	Parsley-piert	Spurge, prostrate
Aster, wild	Cocklebur, spiny	Horsenettle	Parsnip	Spurge, spotted
Bedstraw	Coffebean	Horsetail	Parsnip, wild	Spurweed
Bedstraw, pinewoods	Coffeeweed	Ironweed	Pearlwort	St. Johnswort
Beggartick	Coreopsis, plants	Ivy, Poison	Pennycress	Stinkweed
Betony, Florida	Creeping Jenny	Jewelweed	Pennywort	Stitchwort
Biden	Croton	Jimsonweed	Peppergrass	Strawberry, wild
Bindweed	Croton, wooly	Knapweed, spotted	Pepperweed	Sumac
Bindweed, field	Cudweed	Knawel	Pepperweed, Virginia	Sunflower
Bindweed, hedge	Curly indigo	Knotweed	Pigweed	Sunflower, false
Bitter wintercress	Daisy, English	Knotweed, prostrate	Pigweed, prostrate	Sweet Potato, wild
Bittercress, hairy	Daisy, oxeye	Kochia	Pigweed, redroot	Tansy ragwort
Bitterweed	Dandelion	Lambsquarter	Pigweed, Russian	Tanweed
Black-eyed Susan	Dandelion, fall	Lespedeza	Pigweed, smooth	Tarweed
Bloodflower, milkweed	Dandelion, false	Lettuce, blue	Pigweed, tumble	Thistle
Boxelder	Dock	Lettuce, prickly	Scarlet pimpernel	Thistle, blessed
Brassbuttons	Dogbane	Lettuce, wild	Plantain	Thistle, bull
Bristly oxtongue	Dogfennel	Locoweed	Plantain, black-seed	Thistle, Canada
Broomweed	Elderberry	Lupine	Plantain, bracted	Thistle, musk
Buckhorn	Fiddleneck	Mallow	Plantain, broadleaf	Thistle, Russian
Buckwheat, wild	Fieldcress, Austrian	Mallow, Indiana	Plantain, buckhorn	Thistle, Scotch
Bur ragweed	Flaree, redstem	Mallow, Venice	Plantain, hoary	Toadflax
Burdock	Flax, false	Marigold, roundleafed	Plantain, narrowleaf	Trailing crownvetch
Burweed	Fleabane (daisy)	Marigold, wild	Plantain, slender	Trefoil, tick
Buttercup	Fleabane, hairy	Marshelder	Plantain, woolly	Tumbleweed
Buttercup, bulbous	Fleabane, rough	Matchweed	Pokeweed	Velvet leaf
Carpetweed	Flixweed	Medic, black	Poorjoe	Veronica
Carrot, wild	Florida pusley	Mexicanweed	Povertyweed	Vervain
Catchweed bedstraw	Four-o'clock, wild	Morningglory	Prickly sida	Vervain, blue
Catnip	Frenchweed	Morningglory, woolly	Primrose	Vervain, hoary
Catsear	Galinsoga	Mugwort	Puncture vine	Vervain, prostrate
Catsear, spotted	Galinsoga, smallflower	Mullein, common	Purslane	Vervain, tall
Chamomile, corn	Garlic, wild	Mustard	Radish, wild	Vetch
Chickweed	Geranium, Carolina	Mustard, black	Ragweed	Vetch, bird
Chickweed, common	Geranium, wild	Mustard, garlic	Rape, wild	Vetch, milk
Chickweed, mousear	Goathead	Mustard, hedge	Rose, Southern wild	Vetch, narrowleaf
Chicory	Goatsbeard	Mustard, tansy	Rush	Vetch, wild
Cinquefoil	Goldenrod	Mustard, tumble	Salsify, Western	Virginia buttonweed
Cinquefoil, rough	Ground-ivy	Mustard, white	Shepherdspurse	Virginia creeper
Clematis, Western	Groundsel	Mustard, wild	Smartweed	Wavyleaf bullthistle
Clover	Gumweed	Nettle	Sneezeweed	Willow
Clover, annual yellow	Hawkweed	Nettle, bull	Sorrel, red	Witchweed
Sweet	Hawkweed, orange	Nettle, burning	Sorrel, sheep	Woodsorrel
Clover, crimson	Healall	Nettle, dead	Sowthistle	Wormseed
Clover, hop	Heartleaf drymary	Nettle, stinging	Sowthistle, common	Yarrow
Clover, red	Heathaster	Nettle, tall	Spanishneedle	Yellow rocket
Clover, strawberry	Hemlock, poison	Oak, poison	Spatterdock	Yellowflower
Clover, sweet	Hemp	Onion, wild	Speedwell	pepperweed

Some of these species may require repeat spot applications and/or use of higher rate listed on this product label even under ideal conditions for application. Dilute only with water.



### ORNAMENTAL LAWNS, TURF, AND SOD FARMS

SITE	USE RATE		SPRAY VOLUME IN GALLONS		APPLICATION METHODS AND INSTRUCTIONS
	Fl. Oz. Per 1,000 Sq. Ft.	Pints per Acre	Per 1,000 Sq. Ft.	Per Acre	
<b>Cool Season Grasses</b> (Bluegrass Fescue spp. Ryegrass)	.73 to 1.10	2 to 3	0.02	1 to 4	<b>Controlled Droplet and Spinning Disc Applicators</b>
			up to 0.11	up to 5	<b>Low Volume Ride On, Walk Behind and Towable Applicators</b> (such as Perma Green™ and Turf Tamer®) <b>Note:</b> Use only equipment that is capable of delivering uniform coverage.
			0.11	5 to 10	<b>Low Volume Boom Sprayers</b> Limited coverage of the target leaf surface may reduce performance.
			0.23 to 1.84	10 to 80	<b>Conventional Boom Sprayers</b>
			0.46 to 4.6	20 to 200	<b>Professional Lawn Care Hand Held Gun</b> <b>Note:</b> For optimum results use spray volumes between 85 and 135 gallons per acre.
	—	per Gallon per 1,000 Sq. Ft.	—	<b>Portable Sprayers</b> (such as back packs, hand-held cans and atv units)	
<b>Bentgrass</b> (excluding Golf Course Greens and Tees)	0.55 to 0.73	1.5 to 2	.023 to 5.5	20 to 240	<b>Conventional Boom Sprayers</b> Avoid applications during periods when turf is under stress due to high heat, humidity and reduced moisture. Slight yellowing may occur but will disappear in about seven days.
<b>Warm Season Grasses</b> (Bahigrass, Common Bermudagrass, and Zoysiagrass)	0.73 to 1.10	2 to 3	0.02 to 0.09	1 to 4	<b>Controlled Droplet and Spinning Disc Applicators</b>
			up to 0.11	up to 5	<b>Low Volume Ride On, Walk Behind and Towable Applicators</b> (such as Perma Green™ and Turf Tamer®) <b>Note:</b> Use only equipment that is capable of delivering uniform coverage.
			0.23 to 0.55	5 to 10	<b>Low Volume Boom Sprayers</b> Limited coverage of the target leaf surface may reduce performance.
			0.23 to 1.84	10 to 80	<b>Conventional Boom Sprayers</b>
			0.46 to 4.6	20 to 200	<b>Professional Lawn Care Hand Held Gun</b> <b>Precautions:</b> Use lower rates when turf is under drought stress. Applications made during transition from dormancy to green up may delay green up or cause temporary discoloration.
	.73 to 1.10		per Gallon per 1,000 Sq. Ft.		<b>Portable Sprayers</b> (such as back packs, hand-held cans and atv units)



## USE RESTRICTIONS

### Postemergence:

Limited to two applications per year.

Maximum of 4 pints of product per acre per application.

The maximum seasonal rate is 8 pints per acre excluding spot treatments.

For sod farms, use a minimum of 21 days between applications.

Use of this product in certain portions of California, Oregon, and Washington is subject to the January 22, 2004 Order for injunctive relief in Washington Toxics Coalition, et. al. v. EP, C01-0132C, (W.D. WA). For further information, please refer to <http://www.epa.gov/espp/wtc/>.

## STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

**STORAGE:** This product should be stored in its original container in a cool, dry locked place out of the reach of children and out of direct sunlight. Do not use or store near heat or open flame.

**PESTICIDE DISPOSAL:** To avoid wastes, use all material in this container by application according to label directions. If waste cannot be avoided, offer remaining product to a waste disposal facility or pesticide disposal program (often such programs are run by state or local governments or by industry).

### CONTAINER DISPOSAL:

**Nonrefillable Containers 5 Gallons or Less:** Nonrefillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. **Triple rinse as follows:** Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by State and local authorities. Plastic containers are also disposable by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

**Nonrefillable Containers Larger than 5 Gallons:** Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available. Triple rinse or pressure rinse container (or equivalent) promptly after emptying. **Triple rinse as follows:** Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. **Pressure rinse as follows:** Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 psi for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

**Refillable Container Larger than 5 Gallons:** Refillable container. Refill this container with pesticide only. Do not reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from this container into application equipment or a mix tank. Fill the container about 10% full with water and, if possible, spray all sides while adding water. If practical, agitate vigorously or recirculate water with the pump for two minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times.





### WARRANTY DISCLAIMER

The directions for use of this product must be followed carefully. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, (1) THE GOODS DELIVERED TO YOU ARE FURNISHED "AS IS" BY MANUFACTURER OR SELLER AND (2) MANUFACTURER AND SELLER MAKE NO WARRANTIES, GUARANTEES, OR REPRESENTATIONS OF ANY KIND TO BUYER OR USER, EITHER EXPRESS OR IMPLIED, OR BY USAGE OF TRADE, STATUTORY OR OTHERWISE, WITH REGARD TO THE PRODUCT SOLD, INCLUDING, BUT NOT LIMITED TO MERCHANTABILITY, FITNESS FOR A PARTICULAR PURPOSE, USE, OR ELIGIBILITY OF THE PRODUCT FOR ANY PARTICULAR TRADE USAGE. UNINTENDED CONSEQUENCES, INCLUDING BUT NOT LIMITED TO INEFFECTIVENESS, MAY RESULT BECAUSE OF SUCH FACTORS AS THE PRESENCE OR ABSENCE OF OTHER MATERIALS USED IN COMBINATION WITH THE GOODS, OR THE MANNER OF USE OR APPLICATION, INCLUDING WEATHER, ALL OF WHICH ARE BEYOND THE CONTROL OF MANUFACTURER OR SELLER AND ASSUMED BY BUYER OR USER. THIS WRITING CONTAINS ALL OF THE REPRESENTATIONS AND AGREEMENTS BETWEEN BUYER, MANUFACTURER AND SELLER, AND NO PERSON OR AGENT OF MANUFACTURER OR SELLER HAS ANY AUTHORITY TO MAKE ANY REPRESENTATION OR WARRANTY OR AGREEMENT RELATING IN ANY WAY TO THESE GOODS.

### LIMITATION OF LIABILITY

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO EVENT SHALL MANUFACTURER OR SELLER BE LIABLE FOR SPECIAL, INCIDENTAL, OR CONSEQUENTIAL DAMAGES, OR FOR DAMAGES IN THEIR NATURE OF PENALTIES RELATING TO THE GOODS SOLD, INCLUDING USE, APPLICATION, HANDLING, AND DISPOSAL. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, MANUFACTURER OR SELLER SHALL NOT BE LIABLE TO BUYER OR USER BY WAY OF INDEMNIFICATION TO BUYER OR TO CUSTOMERS OF BUYER, IF ANY, OR FOR ANY DAMAGES OR SUMS OF MONEY, CLAIMS OR DEMANDS WHATSOEVER, RESULTING FROM OR BY REASON OF, OR ARISING OUT OF THE MISUSE, OR FAILURE TO FOLLOW LABEL WARNINGS OR INSTRUCTIONS FOR USE, OF THE GOODS SOLD BY MANUFACTURER OR SELLER TO BUYER. ALL SUCH RISKS SHALL BE ASSUMED BY THE BUYER, USER, OR ITS CUSTOMERS. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BUYER'S OR USER'S EXCLUSIVE REMEDY, AND MANUFACTURER'S OR SELLER'S TOTAL LIABILITY SHALL BE FOR DAMAGES NOT EXCEEDING THE COST OF THE PRODUCT.

**If you do not agree with or do not accept any of directions for use, the warranty disclaimers, or limitations on liability, do not use the product, and return it unopened to the Seller, and the purchase price will be refunded.**

(RV051311)

<sup>1</sup> The State Sale and Use Prohibitions of this product must be part of printed label text for any product which might possibly enter the channels of trade in the states of New York, California, Oregon and/or Washington. The label prohibitions are optional for all other states.

Millennium is a registered trademark of Nufarm Australia Limited

